

University of Tasmania Open Access Repository

Cover sheet

Title

Chemical compositions, feed intakes and digestibilities of crop residue based rations in non-lactating Red Sokoto goats in the subhumid zone of Nigeria

Author

Malau-Aduli, BS, Eduvie, LO, Lakpini, CAM, Malau-Aduli, AEO

Bibliographic citation

Malau-Aduli, BS; Eduvie, LO; Lakpini, CAM; Malau-Aduli, AEO (2003). Chemical compositions, feed intakes and digestibilities of crop residue based rations in non-lactating Red Sokoto goats in the subhumid zone of Nigeria. University Of Tasmania. Journal contribution.

https://figshare.utas.edu.au/articles/journal_contribution/Chemical_compositions_feed_intakes_and_digestibilitie lactating_Red_Sokoto_goats_in_the_subhumid_zone_of_Nigeria/22847636

Is published in: 10.1046/j.1344-3941.2003.00091.x

Copyright information

This version of work is made accessible in the repository with the permission of the copyright holder/s under the following,

Licence.

If you believe that this work infringes copyright, please email details to: oa.repository@utas.edu.au

Downloaded from University of Tasmania Open Access Repository

Please do not remove this coversheet as it contains citation and copyright information.

University of Tasmania Open Access Repository

Library and Cultural Collections University of Tasmania Private Bag 3 Hobart, TAS 7005 Australia E oa.repository@utas.edu.au Chemical compositions, feed intakes and digestibilities of crop -residue based rations in non-lactating Red Sokoto goats in the sub -humid zone of Nigeria

Bunmi Sherifat MALAU-ADULI¹, Lawrence EDUVIE², Clarence LAKPINI² and Aduli Enoch

Othniel MALAU-ADULI2*

1. Department of Animal Science, Ahmadu Bello University, PMB 1044 Zaria, Nigeria.

National Animal Production Research Institute, Ahmadu Bello University, PMB 1096 Zaria, Nigeria.

*Present address of corresponding author: Laboratory of Animal Breeding and Reproduction,

National Agricultural Research Centre for Western Region,

60 Yoshinaga, Kawai, Oda, Shimane 694-0013, Japan.

E-mail: aduli40@yahoo.co.uk, aduli@affrc.go.jp

Tel: +81-90-6837-1953, +81-854-82-1285

Fax: +81-854-82-2280

RUNNING TITLE: CROP-RESIDUES IN RED SOKOTO GOAT FEEDING

Abstract

This study was conducted to evaluate the chemical composition, intake and digestibility of crop-residue based rations by Red Sokoto goats maintained on natural pastures and Digitaria smutsii hay during the dry season of the year. Twenty eight non-lactating does were blocked for weight and assigned to 7 treatment groups comprising of 4 does each in a completely randomized design. Ration A the conventional concentrate ration, was used as the positi ve control, Rations B and C were the two crop residue based test rations, while Ration D the unsupplemented treatment, was used as the negative control. Each of the supplementation rations was fed at 1 and 2% of the doe's body weight. Ration A had the high est crude protein percentage of 17.19% while Rations B and C had 9.54 and 10.38% respectively. The naturally grazed pastures and Digitaria smutsii hay (Ration D) contained the least protein: 2.76 and 4.75% respectively. Ration D also had the highest percen tages of acid detergent fibre, neutral detergent fibre and lignin (49.14, 74.73 and 9.49% in hay and 50.29, 8.27 and 11.5% in grazed pastures, respectively). Ration A on the other hand, had the least percentages of acid detergent fibre (20.00%), neutral detergent fibre (40.01%) and lignin (4.64%). The results indicated that the supplemented group of does had significantly higher (P<0.05) dry matter and crude protein intakes as well as nutrient digestibilities than the unsupplemented groups. A comparison of the unsupplemented animals with all the other treatment groups revealed that dry matter digestibility improved by a range of 4.1 to 27.9%, while crude protein digestibility improved by 17.1 to 42.2%, the highest value being in does on Ration A. It was conc luded that goats were able to subsist and make appreciable gains in the long dry season on crop -based diets that compared favourably with the conventional concentrate rations. Of the two tested crop-residue based rations, Ration C is a better supplementation package than Ration B. Key words: Red Sokoto goats, crop residue, feed intake, digestibility, dry season

Introduction

The Northern regions of Nigeria are concentrated with most of the nation 's ruminant livestock. However, these areas are characterised by a long and pronounced dry season (6-9 months), and this often causes serious shortage of feed for the ruminants. The problem of dry season livestock feeding in particular, has directed research efforts towards harnessing and enhancing the utilization of abundant arable by-products and crop residues. The abundance of crop residues makes them cheap sources of nutrients for ruminants. Nevertheless, they are generally low in nutrients (Nicholson 1984). Various strategies have been adopted in improving their nutrients and utilisation. One of such is by judicious supplementation to provide the m ost limiting nutrients (Preston 1982; Alhassan 1988). Balancing the nutrients that provide the major building blocks for tissue synthesis and milk production should be the primary concern of the nutritionist (Leng 1990). This can be achieved by careful blending of the crop residues to meet the nutrient requirements of the ruminants.

Supplementation using residues such as maize offal (Fadugba 1990), groundnut haulms and shells (Adu & Lakpini 1983; Ikhatua & Adu 1984; Alawa & Umunna 1993) and cowpea vines and husks (Alhassan et al. 1984) have been documented. From these studies, the inclusion of supplements in small ruminant rations has often resulted in increased feed intake as a compensation for the reduced energy concentration of such diets. However, appropriate packages to guide small ruminant producers have not been developed. Also, relevant information on cheap, alternative sources has not been passed on to the producers. Therefore, this study was undertaken to determine the nutrient composition and digestibilities of some locally available and affordable crop residue feed resources in comparison with the conventional supplementation in Red Sokoto goats in the sub -humid zone of Nigeria.

Materials and methods

Location and management of experimental animals

The experiments were conducted in the Experimental Unit of the Small Ruminant Research Programme of the National Animal Production Research Institute, Shika, Zaria, Nigeria. Shika falls between latitudes 11 and 12°N and between longitudes 7 and 8°E, with an altitude of 640m above sea level. Shika is located within the Northern Guinea Savannah Zone with an average annual rainfall and temperature of 1,107 mm and 24.4 °C respectively. The seasonal distribution of the annual rainfall is approximately 0.1% (11.0 mm) in the late dry season (January-March), 25.8% (285.6 mm) in the early-wet season (April-June), 69.6% (770.4 mm) in the late-wet season (July-September) and 4.5% (49.8 mm) in the early dry season (October - December). The experiments were conducted during the dry seasons (between October and March). The animals were routinely dewormed with anthelmintic drugs and dipped in an acaricide (Asuntol) solution against ecto parasites. The animals were housed in well-ventilated pens during the night.

Digestibility trial: Twenty eight adult Red Sokoto does ranging between 24.6 and 26.4 kg were used for this experiment. The animals were balanced for weight and blocked into seve n groups with four animals per group. The component ingredients in the different rations are shown in Table 1. Each doe was individually offered its appropriate corresponding ration to evaluate the digestibility of the diets. The study comprised a two -week preliminary period of realimentation and adjustment, and one week of sample collection. The animals were housed in individual metabolism cages with facilities for separate collection of faeces and urine. The animals were weighed at the beginning and end of the study. Feces were collected each morning just before feeding. 10% of each daily faecal output was collected for chemical analyses. Samples of the different rations fed were taken daily and bulked, from which sub -

samples were taken for laboratory analysis. Also, samples of the individual feed ingredients were analysed in the laboratory. Water was made available to the animals ad libitum. The inventory, abundance and palatability of the plant species in the grazed paddock was conducted as described by Lakpini et al. (1997).

Laboratory analyses: Proximate analyses of feed and faecal samples were carried out by the AOAC (1980) methods. Dry matter of samples was determined in an oven at 105 °C for 48 hours. Nitrogen determination was by the Micro Kjedahl met hod, while the Soxhlet extraction procedure was used for ether extraction. Crude fibre was determined by alternate refluxing with weak solutions of H_2SO_4 and KOH. The detergent fibre fractions (Neutral detergent fibre, acid detergent fibre and lignin) were determined according to Goering and Van Soest (1970). Dry matter intake (DMI) was determined using the following equation:

DMI (g/day) = %DM/100 x feed intake.

Dry matter digestibility (DMD (%)) was calculated as:

100 x [DM intake (g) – DM output (g)] /DM intake (g)].

The other digestibilities were calculated as above.

Statistical analysis: Differences in intake and digestibility were analysed using the Generalised Linear Models Procedure (PROC GLM) of SAS (1987) in a one -way analysis of variance. The effect of treatment was tested and significant differences between treatment means established by Duncan's Multiple Range Test.

Results

The chemical composition of the individual feed ingredients and the experimental diets are shown in Tables 2 and 3 respectively. Table 3 shows that all the rations had high dry matter (DM) contents with a mean value of about 95%. Ration A had the highest crude protein (CP) value of 17.19% followed by Rations B and C with 9.54 and 10.38%, respectively. The CP

value of the grazed pastures (dry season) was the lowest (2.76%), while that of Digitaria hay was 4.75%. The least ash content value was obtained in Ration D with the hay having 8.47% and the grazed pasture having 7.02%. The highest ash content value of 13.85% was obtained in Ration A. Ration A also had the highest ether extract (EE) value of 14.08% and Ration D, the least (0.78 and 2.40% for grazed pasture and Digitaria hay, respectively). Acid detergent fibre (ADF), neutral detergent fibre (NDF) and lignin contents were I east in Ration A with values of 20.00%, 40.01% and 4.64% respectively. Ration D on the other hand, contained the highest ADF, NDF and lignin values of 49.14%, 74.73% and 9.49% for hay and 50.29, 80.27 and 11.5% for grazed pasture respectively. The calculated chemical analysis of the experimental rations (Table 3) reveals that Ration A had a CP content of 17.05% while the test rations B and C had 9.82 and 10.85%, respectively. Ration A also had a metabolisable energy (ME) of 11.17 MJ/kg DM, while Rations B and C had 10.29 and 10.17 MJ/kg DM respectively. Table 4 shows the DM and CP intakes and digestibilities of the nutrients. The Table shows that generally, the supplemented groups had significantly higher (P<0.05) DM and CP intakes and digestibilities than the unsupplemented group except animals on Ration B that had similar values to the unsupplemented group. It was also evident that increasing the level of supplementation also resulted in increased DM and CP intakes of all the experimental rations, with thes e increases being significant (P<0.05) and similar for Rations A and C. It was also observed that supplementation increased the digestibility of all the nutrients. However, animals on Ration B recorded very poor digestibility values and their counterparts in the unsupplemented group had the least. Even though the digestibility of nutrients decreased with increasing levels of supplementation, these decreases were not significant (P>0.05). Ration 1A (the conventional concentrate at 1% of body weight) gave the highest digestibility values. A comparison of the unsupplemented animals with all the other treatment groups reveals that DM digestibility improved by a range of 4.1 to 27.9% and CP digestibility by 17.1 to 42.2%, the highest being in

animals on Ration A at 1% level. Similar improvements trends were also noticeable for neutral detergent fibre (NDF) and acid detergent fibre (ADF). A simple economic analysis (Table 4) revealed that the conventional concentrate feed was the most expensive for supplementation particularly, at the 2% level (4.42 naira per animal per day). Of the two tested crop -residue rations, Ration 1B was significantly cheaper (P<0.05) than Rations 2B and 2C, but similar to Ration 1C.

Discussion

The quantity and type of ingredients used in formulating the rations influenced their chemical compositions. In Ration A, the inclusion of wheat offal and cottonseed cake boosted the protein level of the ration giving it a value of 17.19%. This value is higher than the recommended CP level of 15% for optimum maintenance of production by Nuru (1985) and 8.9 – 16.0% by NRC (1975). The metabolisable energy (ME) value of Ration A (11.17 MJ/kg DM) is also higher than the 9.5 MJ/kg DM recommended for maintenance by INRA (1988), but lower than the latter's recommended value for pregnant and lactating 65 -kg Saanen goats yielding 4 Kg of milk per day. Rations B and C had similar CP and ME values indicating that they are isocaloric and isonitrogenous rations. The preponderance of crop residues in Rations B and C was responsible for their high crude fibre and lignin levels. The current study showed that inspite of Rations B and C being isocaloric and isonitrogenous, animals on Ration C had better intakes and digestibilities than those on Ration B, possibly due to t he low palatability, hence low voluntary intake, and poor digestibility of Ration B.

The observed higher digestibilities of DM, CP, NDF and ADF at 1% level in comparison to 2% level can be attributed to the higher feed intake at the 2% level of inclusion. It has been established that higher feed intake results in a faster rate of passage of digesta from the reticulo-rumen (Swan & Lamming 1967). This does not allow for effective degradation,

hence lowering the digestibility of feed. Increasing the level of crop residue inclusion in the diet also increases the amount of lignin, which depresses the digestibility of the ration (McDonald et al. 1988), because the rate of microbial colonisation of a feed with high fibre content is comparatively lower (Silva & Orskov 1988). The poor intake and digestibility values obtained for the unsupplemented animals is due to the fact that Ration D had crude protein levels that are below the recommended minimum values for maintenance. This shows that there is the need for dry season supplementation in goats because the available feeds at that tim e are limiting in crude protein. It should also be clearly stated at this point that because of cumulative effects derived from residuals, there was a difference in chemical composition between Tables 2 and 3. It was obvious that these amounts of residuals were dependent on feeding level and feed quality, for instance, it was possible to obtain differences between Rations 1B and 2B as clearly demonstrated in Table 4 where DMI significant ly differed between Rations 1B (0.21 kg/day) and 2B (0.30 kg/day).

Of the two tested rations, Ration C seemed to have produced better intakes and digestibilities in the animals, possibly due to the composition of the rations. It contained maize offal which has very low fibre content (Alawa & Umunna 1993), groundnut haulms which have been demonstrated to be better quality roughages than Digitaria smutsii hay and contain adequate protein to maintain ruminants without any form of supplementation during the pe riods of feed scarcity (Ikhatua & Adu 1984). The groundnut shells fed to the animals were also crushed before inclusion into the ration as suggested by Alawa and Umunna (1993). This must have aided their consumption and digestibility. Even though Ration B contained groundnut haulms, the combination of Guinea corn bran and cowpea husk which had low crude protein percentages, must have reduced the intake and digestibility of the ration. Alhassan et al. (1984) observed lower digestibility values in sheep and g oats (48.8 and 56.3% respectively) compared with cattle (73.6%) when they fed them cowpea vines. This might imply that cattle

do better on cowpea residues than small ruminants. From the economic analysis, the high cost of the conventional concentrate ration shows that it is beyond the reach of a typical smallholder goat farmer, whereas the crop-residue based rations seem quite affordable. Even though Ration B had the least cost, it was glaring that it had lower intake and digestibility compared to Ration C, indicating in essence, that Ration C had a better efficiency of utilisation

In conclusion, goats are able to subsist and make appreciable gains even in the long dry seasons of the sub-humid zone of Nigeria on crop-residue based diets that compare favourably with conventional concentrate rations. Ration C is a better package than Ration B and is therefore recommended and at 1% level of inclusion, to small ruminant farmers due to its high intake and digestibility as well as its affordability than at 2% leve I of inclusion.

Acknowledgement

The authors gratefully acknowledge the support of the International Atomic Energy Agency (IAEA) Vienna, Austria in funding this project, and to the Director of the National Animal Production Research Institute, Ahmadu Bello University Shika-Zaria for permission to publish this work

References

- Adu IF, Lakpini CAM. 1983. Effect of feeding chopped and unchopped groundnut haulms ("Harawa") on nutrient utilisation and the production of rumen metabolites in Yankasa lambs. Nigerian Journal of Animal Production 10, 110-113.
- Alawa JP, Umunna NN. 1993. Alternative feed formulation in the developing countries: Prospects for utilisation of agro-industrial by-products. Journal of Animal Production Research 13, 63-98.
- Alderman G. 1985. Prediction of the energy value of compound feeds. In: Recent advances in animal nutrition. W. Haresign and D.J.A. Cole (E ditors), Butterworths, London, UK.
- Alhassan WS. 1988. Crop residue utilisation with special reference to pastoral production: Past, present and future. Proceedings, National Conference on Pastoralism in Nigeria, 26-27th June, 1988, 70-87.
- Alhassan WS, Ehoche OW, Adu IF, Obilara TA, Kallah MS. 1984. Crop residue potential of agricultural development projects: Nutritive value and residue management. NAPRI Annual Report, National Animal Production Research Institute, Shika, Nigeria, 35-45.

- AOAC. 1980. Association of Official Analytical Chemists. Official Methods of Analysis 13th edn., Washington D.C., USA.
- Fadugba AA. 1990. Evaluation of the nutritive value of maize offal in poultry diets. M.Sc Thesis, Ahmadu Bello University Zaria, Nigeria.
- Goering HK, Van Soest PJ. 1970. Forage fibre analysis (apparatus, reagents, procedures and some applications). Agricultural Handbook 379, ARS, USDA, Washington D.C.
- Ikhatua UI, Adu IF. 1984. A comparative evaluation of the utilization of groundnut haulms and Digitaria hay by Red Sokoto goats. Journal of Animal Production Research 4, 145-152.
- INRA. 1988. Institut National de la Recherché Agronomique. Alimentation des bovins, ovins et caprins, INRA, Paris, France.
- Lakpini CAM, Balogun BI, Alawa JP, Onifade OS, Otaru SM. 1997. Effects of graded levels sun-dried cassava peels in supplement diets fed to Red Sokoto goats in the first trimester of pregnancy. Animal Feed Science Technology 67, 197-204.
- Leng RA. 1990. Factors affecting the utilisation of poor quality forages by ruminants particularly under tropical conditions. Nutrition Research Review 3, 277-303.
- McDonald P, Edwards RA, Greenhalgh JFD. 1988. Animal Nutrition (4th Edition). Longman Scientific and Technical, England, UK.
- Nicholson JWG. 1984. Digestibility, nutritive values and intake. In: Sundstol F, Owen E (eds), Straw and other fibrous feeds, pp. 85-90, Elsevier Publishers, Amsterdam, Netherlands.
- NRC. 1975. National Research Council. Nutrient requirements of domestic animals 5 th Edition), Nutrient requirements of sheep, NRC, Washington DC, USA.
- Nuru S. 1985. Strategies for improved production of fresh foods in Nigeria. Paper presented at the Nigerian Institute of Food Science and Technology Symposium, Federal Industrial Research Organisation (FIRO), Oshodi, Lagos, Nigeria, 13 th July, 1985.
- Preston TR. 1982. Nutritional limitations associated with the feeding of tropical forages. Journal of Animal Science 54, 877-884.
- SAS 1987. Statistical Analysis System, SAS Institute, Cary, North Carolina, USA.
- Silva AT, Orskov ER. 1988. The effect of five different supplements on the degradation of straw in sheep given untreated barley straw. Animal Feed Science and Technology 19, 289-298.
- Swan H, Lamming GE. 1967. Studies on the nutrition of ruminants: II. The effect of crude fibre in maize-based rations on the carcass composition of Friesian steers. Animal Production 9, 203-208.

Ration	Ingredients	% inclusion	Remarks		
1A + Basal diet	Maize	40	Positive control		
	Wheat offal	35	(conventional concentrate)		
	Cottonseed cake	20	offered at 1% of body		
	Bone meal	3	weight		
	Salt	2	J. J		
2A + Basal diet			Positive control		
	~	✓	(conventional concentrate)		
			offered at 2% of body		
			weight		
1B + Basal diet	Guinea-corn bran	39.5	Test Ration 1 offered at 1%		
	Cowpea husk	30	of body weight		
	G/Nut haulms	30	,		
	Salt	0.5			
2B + Basal diet			Test Ration 1 offered at 2%		
	✓	✓	of body weight		
1C + Basal diet	Maize offal	49.5	Test Ration 2 offered at 1%		
	G/Nut shells	20	of body weight		
	G/Nut haulms	30	5 5		
	Salt	0.5			
2C + Basal diet			Test Ration 2 offered at 2%		
	~	✓	of body weight		
D	Digitaria hay and natural	Ad libitum	Negative control		
(Basal Diet)	grazed pasture		(unsupplemented)		

Table 1 Component ingredients in the different rations

Feedstuff	DM	СР	CF	Ash	EE	NFE
Maize	90.73	9.56	2.20	9.67	4.05	74.52
Wheat offal	87.60	16.90	11.30	6.40	3.80	61.60
Cottonseed cake	93.60	29.94	23.50	5.16	5.76	35.64
Bone meal	75.00	36.00	3.00	49.00	4.00	8.00
Guinea corn bran	93.33	7.60	24.80	6.95	3.01	59.90
Cowpea husks	91.41	7.10	33.40	7.14	0.65	58.91
Groundnut haulms	93.65	15.63	23.26	8.00	2.43	51.00
Maize offal	89.07	10.08	1.50	0.80	1.70	60.30
Groundnut shells	96.05	5.90	31.8	8.50	1.31	50.30

Table 2 Chemical composition of the major feed ingredients (DM basis) (%)

Ration	DM	СР	Ash	EE	ADF	NDF	LIGNIN		
Ration A	93.87	17.19	13.85	14.08	20.00	40.01	4.64		
Ration B	94.97	9.54	10.55	10.43	38.10	68.42	8.94		
Ration C	95.94	10.38	11.97	12.45	36.65	54.74	8.23		
Ration D Hay	94.78	4.75	8.47	2.40	49.14	74.73	9.49		
Ration D Natural pastures	96.26	2.76	7.02	0.78	50.29	80.27	11.5		
Calculated analysis of the experimental r ations									
R	Ration A		Ration	3 Ra		ation C			
CP (%) 1	17.05		9.82	2		10.85			
ME (MJ /kg DM) 1	11.17		10.29	1		10.17			

Table 3 Chemical composition of the experimental diets (dry matter basis) (%)

The ME values of the experimental rations were calculated as per Alderman (1985) as follows: ME (MJ/kg DM) = 11.78 + 0.00654CP + (0.000665EE)² – CF(0.00414EE) – 0.0118A where CP = Crude Protein, EE = Ether Extract, CF = Crude Fibre, A = Ash

Ration	1A	2A	1B	2B	1C	2C	D	SEM
Nutrient intake	(kg/day)							
DMI	0.24 ^b	0.47ª	0.21 ^{bc}	0.30 ^{ab}	0.23 ^b	0.42ª	0.15 ^c	± 0.02
CPI	0.044ª	0.087ª	0.012 ^b	0.017 ^b	0.032ª	0.072ª	0.009 ^b	± 0.01
Apparent diges	tibility of nut	rients (%)						
DM	84.3ª	83.0ª	62.5 ^d	60.5 ^e	75.8 ^b	67.8 ^c	56.4 ^f	± 2.84
СР	90.6ª	89.2ª	69.5 ^d	65.5 ^e	82.7 ^b	78.1 ^c	48.4 ^f	± 3.07
NDF	69 .5ª	66.6 ^b	62.1c ^d	61.9 ^d	65.9 ^b	63.7c	60.1 ^e	± 3.23
ADF	51.7ª	49 .8ª	43.9 ^{bc}	42.8 ^c	46.1 ^b	44.4 ^{bc}	42.3 ^{bc}	± 5.01
Economic analy	<u>sis of the fe</u>	<u>eds (Naira)*</u>						
Cost of feed consumed per animal per day	2.19 ^b	4.42ª	0.50 ^e	1.06 ^d	0.83d ^e	1.55°	-	± 0.15

Table 4 Mean nutrient intake, apparent digestibility coefficients and cost of the experimental diets

a,b,c,d,e,f means within the same row bearing different superscript letters differ significantly (P<0.05) Naira = Nigerian currency (100 kobo make 1 naira and current exchange rate is 1US\$ = 140 Naira)