

University of Tasmania Open Access Repository

Cover sheet

Title

Factors affecting egg production and layer bird mortality in private poultry farms in the subhumid zone of Nigeria

Author

Malau-Aduli, AEO, Bawa, GS, Joel, MK

Bibliographic citation

Malau-Aduli, AEO; Bawa, GS; Joel, MK (2003). Factors affecting egg production and layer bird mortality in private poultry farms in the subhumid zone of Nigeria. University Of Tasmania. Journal contribution.
https://figshare.utas.edu.au/articles/journal_contribution/Factors_affecting_egg_production_and_layer_bird_mor

Is published in: [10.1046/j.1344-3941.2003.00111.x](#)

Copyright information

This version of work is made accessible in the repository with the permission of the copyright holder/s under the following,

Licence.

If you believe that this work infringes copyright, please email details to: oa.repository@utas.edu.au

Downloaded from [University of Tasmania Open Access Repository](#)

Please do not remove this coversheet as it contains citation and copyright information.

University of Tasmania Open Access Repository

Library and Cultural Collections

University of Tasmania

Private Bag 3

Hobart, TAS 7005 Australia

E oa.repository@utas.edu.au

CRICOS Provider Code 00586B | ABN 30 764 374 782

utas.edu.au

ORIGINAL ARTICLE

Factors affecting egg production and layer bird mortality in private poultry farms in the subhumid zone of Nigeria

Aduli Enoch Othniel MALAU-ADULI,¹ Gideon Shaibu BAWA² and Kayode JOEL²

¹National Animal Production Research Institute and ²Department of Animal Science, Ahmadu Bello University, Zaria, Nigeria

ABSTRACT

Egg production and layer bird mortality data obtained from five different private farms in Zaria within the subhumid zone of Nigeria, over a 6-year period (1994–1999) were subjected to least squares analysis to determine the effects of age, season and year. It was demonstrated that age had a highly significant influence ($P < 0.01$) on egg production in that birds falling within the age group 30–39 weeks produced the highest number of eggs (3255 ± 109), while birds over 100 weeks of age produced the lowest number of eggs (1206 ± 412). Similarly, seasonal variation in egg production was also significant ($P < 0.05$); the highest egg production (2926 ± 90) was obtained during the early dry season, and the lowest (2423 ± 95) during the late wet season. Mortality was generally low (0.0–0.9%) and not significantly different from 20 to 49 weeks of age ($P > 0.05$). However, from 50 to 100 weeks of age, highly significant differences ($P < 0.01$) in mortality were observed, with the 90–100 weeks age group recording the highest mortality of $15.7 \pm 1.3\%$. Furthermore, yearly variations in egg production and mortality were highly significant ($P < 0.01$) with 1999 producing the most eggs (4410 ± 102) and recording the lowest mortality (1.4%). It was concluded that due to the significant seasonal, age and yearly variations, data on egg production and mortality need to be adjusted for these effects to enable unbiased comparison between and within farms. Furthermore, farmers can make adequate feeding preparations commensurate with each season for optimal egg production.

KEYWORDS: age, egg production, mortality, season, subhumid zone.

INTRODUCTION

Two major indices for measuring the profitability of raising layer birds in a poultry enterprise are total number of good quality eggs produced and mortality. In a comparative analysis of egg production and mortality between farms, it is important to adjust for all possible sources of variation in order to ensure that such a comparison is unbiased. Kekeocha (1978) and Dafwang (1987) reported that age and disease, among others, are factors that affect the productivity of laying hens. Conclusions drawn from data emanating from government and research institute farms may not necessarily reflect the true situation of production in private farms because the management, personnel and infrastructural facilities are not the same. Therefore, the objective of this study was to investigate the effects of age, season and year on egg production and mor-

tality in five private farms in Zaria located within the subhumid zone of Nigeria.

MATERIALS AND METHODS

Description of the study area and data collection

The data used for this analysis consisted of 6672 records on egg production and mortality from 5 private farms (Farms 1, 2, 3, 4 and 5) in Zaria, Nigeria.

Correspondence: Aduli Enoch Othniel Malau-Aduli, Laboratory of Animal Breeding and Reproduction, National Agricultural Research Center for Western Region, 60 Yoshinaga, Kawai, Oda-shi, Shimane-ken 694–0013 Japan. (Email: aduli40@yahoo.co.uk, aduli@affrc.go.jp)

Received 17 May 2002; accepted for publication 9 January 2003.

Farms 1, 2 and 4 were backyard poultry enterprises with an approximate flock size of 500–800 layers, while Farms 3 and 5 were commercial poultry farms with approximate flock sizes of 2000 and 3000 layers, respectively. All the farms kept birds of the Shika Brown breed. This breed of hen was developed at the National Animal Production Research Institute (NAPRI) Shika, Nigeria, and is known for its persistency of egg production, hardiness and total number of eggs produced. The study area is located in the Northern Guinea Savannah with an average annual rainfall of about 1100 mm, distributed monthly between May and October and a monthly mean temperature of 14–30°C and 21–30°C during the wet and hot dry seasons, respectively.

Management of the birds

The birds in Farms 3 and 5 were raised under an intensive management system. They were housed in three-tier wire battery cages fitted with feeders and waterers, located in tropical type open-sided poultry buildings. Feeds were served once a day at 07.00 hours and fresh clean water provided *ad libitum*. Eggs were picked twice a day in the mornings and in evenings. Laying hens were observed daily and egg production and mortality records were adequately kept. Birds in Farms 1, 2 and 4 were raised under a deep litter system. Feeds were served every morning and clean fresh water was provided *ad libitum*. Egg production and mortality records were taken on a daily basis as in Farms 3 and 5. In all the farms, strict health management routines of vaccination, deworming and delousing were observed.

Data editing and statistical analysis

Variables analyzed included farm, age of birds, season and year. The years were from 1994 to 1999, while season was classified into early dry (November–January), late dry (February–April), early wet (May–July) and late wet (August–October). The general linear models procedure (PROC GLM) of SAS (1986) was utilized in running least squares analysis of variance to test for fixed effects and their interactions on total number of eggs produced and percentage mortality. The analytical model used was:

$$Y_{ijkl} = \mu + F_i + A_j + S_k + YR_l + (FS)_{ik} + (FYR)_{il} + (AS)_{jk} + (AYR)_{jl} + (SYR)_{kl} + e_{ijklm}$$

where Y_{ijklm} = observation on the i^{th} farm of the j^{th} age-group within the k^{th} season of the l^{th} year of production; μ = overall mean; F_i = fixed effect of the i^{th} farm;

A_j = fixed effect of the j^{th} age-group; S_k = fixed effect of the k^{th} season; YR_l = fixed effect of the l^{th} year; $(FS)_{ik} + (FYR)_{il} + (AS)_{jk} + (AYR)_{jl} + (SYR)_{kl}$ = interactions; and e_{ijklm} = random error.

Since all interactions were not significant, these were dropped from the model. Furthermore, means based on farms were not tabulated in the present study because the private farms did not want their individual productivity reflected, to guard against possible administrative backlash in a situation where laxity seemed to be implied by the results. However, there were significant differences between farms.

RESULTS AND DISCUSSION

Effect of season

Seasonal fluctuations in the subhumid zone of Nigeria are very crucial to overall egg production by layer birds, mortality, sales and feed costs. It is typical for temperatures and relative humidity in this zone during the late dry season of February to April to reach 36°C and 21%, respectively. At this period, egg production is at its lowest because the birds drastically reduce their feed intake. On the market scene, egg sales also drop as few consumers buy because of storage problems during the hot season. In contrast, during the late wet season of August to October, when temperature and relative humidity average 25°C and 72%, respectively, the feed intake of the birds increases and egg production also goes up. Therefore, season had a highly significant effect ($P < 0.01$) on egg production and mortality (Table 1). The number of eggs produced was highest in the early dry season (2936 ± 90) and lowest during the late wet season (2423 ± 95). Generally, the hot season does not favor egg production because of the heat stress experienced during this period (Abdu *et al.* 1982). Feed intake decreases with increase in temperature (Payne 1966a). Dafwang (1987) opined that pullets should be reared in such a way that they do not come into peak production during the hottest months of the year. The temperature of Zaria where this work was carried out ranges from 21 to 36°C during the hot season. The thermoneutral zone for optimal production by laying birds has been reported to be 12.6–20.0°C. The temperature attained in the early wet season and late wet season is far higher than 26°C which may account for the reduction in eggs produced during this period. Ogbogu (1988) noted that the comfort zone for layers is between 13 and 21°C and that at 30–40°C, the egg

Table 1 Seasonal variation in egg production and mortality

Season	Total no. days	No. observations	No. birds	No. eggs produced	Mortality (%)
Early dry (Nov–Jan)	90	1162	722 ± 10 ^a	2936 ± 90 ^a	2.4 ± 1.4 ^b
Late dry (Feb–April)	90	2155	710 ± 10 ^a	2600 ± 107 ^b	6.0 ± 1.72 ^a
Early wet (May–July)	90	1152	709 ± 12 ^b	2567 ± 105 ^b	1.9 ± 1.68 ^b
Late wet (Aug–Oct)	90	2203	711 ± 11 ^b	2423 ± 95 ^b	1.2 ± 1.55 ^b

Column means bearing different superscripts are significantly different ($P < 0.01$).

Table 2 Yearly variation in egg production and mortality (Least square means ± SE)

Year	Total no. days	No. observations	No. birds	No. eggs produced	Mortality (%)
1994	365	1102	607 ± 7	2037 ± 186 ^c	10 ± 2.9 ^c
1995	365	1105	628 ± 7	2102 ± 106 ^c	4.8 ± 1.7 ^c
1996	366	1107	488 ± 4	1529 ± 107 ^d	6.3 ± 1.7 ^b
1997	365	1115	530 ± 5	2251 ± 109 ^c	2.5 ± 1.7 ^d
1998	365	1103	848 ± 14	3127 ± 90 ^b	4.7 ± 1.4 ^c
1999	365	1140	909 ± 15	4410 ± 102 ^a	1.4 ± 1.6 ^b

Column means bearing different superscripts are significantly different ($P < 0.01$).

size and number produced dropped drastically due to reduced feed intake. The shell and internal quality of eggs were also poor since the protein, minerals (calcium) and vitamins consumed were not enough to meet the demands of normal egg production. Season had a significant effect ($P < 0.05$) on mortality rate and the late dry season recorded the highest mortality of 6.0%. However, there were no seasonal variations ($P > 0.05$) in mortality during the early dry, early wet and late wet seasons. Payne (1966a) observed a low egg production and mortality in leghorn pullets when raised at a constant temperature of 32°C when compared with those raised in a temperature range of 13–32°C. Furthermore, Payne (1966b) demonstrated that at 37°C or higher for about 8 h, hens started to die of heat prostration while reduced egg size and shell thinning were common.

Effect of year

Year had a significant effect ($P < 0.05$) on egg production (Table 2). The number of eggs produced increased from 1994 to 1999, with 1999 recording the highest number of eggs produced (4410 ± 102). However, the number of eggs produced dropped in 1996 (1529 ± 107) before rising again between 1997 and 1999. In different years, the climatic and weather situations change and this exerts an influence on rainfall, temperature, relative humidity and disease outbreaks. This is the major reason why there were significant differences in the number of eggs produced in different years.

Effect of age

Table 3 shows that age had a significant effect ($P < 0.05$) on egg production. The peak production was between 30 and 39 weeks of age. There was no significant difference ($P > 0.05$) in eggs produced between 30–39 weeks and 40–49 weeks (3255 ± 109 and 3104 ± 110, respectively). Thereafter, there was a decrease in the number of eggs produced till the end of the laying period, from 50–59 weeks, 60–69 weeks, 70–79 weeks, 80–89 weeks, 90–100 weeks and 100 and above (2878 ± 110.83, 2664 ± 113, 2420 ± 112, 2119 ± 130, 2059 ± 182 and 1206 ± 413, respectively). This observation is in agreement with Oluyemi and Robert (1979), who described the laying pattern of the laying hen as having 4 phases: gradual rise, peak, plateau, and gradual decline to the point that it would not be economically justifiable to retain the hen in the flock. A major difference in this work, however, is that the decline in egg production was determined more by festive seasons like Christmas and Sallah when poultry farmers dispose of their flocks because of the high demand for chickens for which consumers are willing to pay up to four times the normal price for a spent layer bird. Therefore, from the result in Table 3, the highest mortality was recorded at 90–100 weeks of age (15.7 ± 3.1) which was statistically the same as the mortality recorded for 100 weeks and above (13.94 ± 7.0). The high mortality rate at this stage may be due to the large number of birds culled as a result of the low number of eggs produced. This is also in agreement with Oluyemi and Robert (1979) who reported

Table 3 Age-group variation in egg production and mortality (Least square means \pm SE)

Age group (weeks)	Total no. days	No. birds	No. eggs produced	Mortality (%)
20–29	70	768 \pm 13	2436 \pm 112 ^c	0.0 \pm 0.1 ^d
30–39	70	734 \pm 12	3255 \pm 109 ^a	0.2 \pm 1.8 ^d
40–49	70	729 \pm 12	3104 \pm 110 ^a	0.9 \pm 1.8 ^d
50–59	70	711 \pm 11	2878 \pm 110 ^b	2.7 \pm 1.9 ^c
60–69	70	690 \pm 9	2664 \pm 113 ^b	1.9 \pm 1.9 ^c
70–79	70	676 \pm 9	2420 \pm 130 ^c	2.6 \pm 2.0 ^c
80–89	70	685 \pm 8	2119 \pm 130 ^d	6.7 \pm 2.2 ^b
90–100	70	690 \pm 9	2059 \pm 182 ^d	15.7 \pm 3.1 ^a
>100	>70	659 \pm 7	1206 \pm 412 ^c	13.9 \pm 7.0 ^a

Column means bearing different superscripts are significantly different ($P < 0.01$).

that from the 62nd week of age, the number of eggs produced begins to drop gradually to practically nil.

CONCLUSION

It can be concluded that season, age and year have significant effects on egg production and mortality, hence there is a need to adjust for these factors in making comparisons between and within farms. For maximum egg production, the best time to start rearing pullets in the study area is the months of January or February, so that their peak production will fall between the months of June and August. For profit maximization, laying hens should not be kept in the flock beyond 62 weeks of age before disposing of them as spent layers.

ACKNOWLEDGMENTS

The authors express their gratitude to the owners of the three backyard poultry enterprises and the two commercial poultry farms in Zaria, Nigeria, for permit-

ting access to their egg production and mortality data without which this work could not have been published.

REFERENCES

- Abdu PA, Bishu G, Adesiyun AA, Adeboye DS. 1982. Survey for *Mycoplasma Gallisepticum* and *Mycoplasma soynoviae* antibodies in chickens in Zaria. *Nigerian Journal of Animal Production* **3**, 63–69.
- Dafwang II. 1987. Hot weather management tips for poultry farmers. *Nigerian Livestock Journal* **7**, 14–18.
- Kekeocha CC. 1978. Improving rural poultry production in Nigeria. *Poultry Science* **79**, 413–437.
- Ogbogu DA. 1988. Coping with ups and downs in poultry production. *Nigerian Livestock Farmer* **8**, 6–9.
- Oluyemi JA, Roberts FA. 1979. *Poultry Production in Warm Wet Climates*. Macmillan Publishers, London.
- Payne CG. 1966a. Environmental temperature and performance of light breed pullets. *Proceedings of the 13th World Poultry Congress, Kiev*, pp. 480–485.
- Payne CG. 1966b. Developments in the use of artificial heating for controlling animal environment. *Proceedings of the Rural Electrification Conference, London*, pp. 23–30.
- SAS. 1986. *Statistical Analysis System*. SAS, Cary, NC.