Epiphytic bryophytes and habitat microclimate variation in lower montane rainforest, Peru.

by

Jerry Romanski BSc. (University of Tasmania)

A thesis submitted in partial fulfilment of the requirements for an Honours Degree at the School of Geography and Environmental Studies, University of Tasmania (November, 2007).

Declaration

This thesis contains no material which has been accepted for the award of any other degree or diploma in any tertiary institution, and to the best of my knowledge and belief, contains no material previously published or written by another person, except where due reference is made in the text of the thesis.

Signed

Jerry Romanski 18 November 2007

Abstract

A survey of epiphytic bryophytes and a study of microclimatic variation within a tree crown were conducted in lower montane rainforest at 2400 m in the Peruvian Yungas. A total of 129 species (87 hepatics, 1 hornwort and 41 mosses) were collected on 3 trees, using different methodologies. A single Weinmannia sp. host supported 110 species (77 hepatics, 1 hornwort and 32 mosses). Species with a wide distribution on the hosts made up 47% of the epiphytic assemblage, 22% were limited to the trunks and 31% occurred only in the crowns. The overall hepatic to moss ratio was 2.15:1. Mosses exhibited a narrower distributional range than hepatics. The greatest species richness and abundance was found on the large branches in the mid-crown. Species found in the mid-crown generally had a broader distribution on hosts than those found on the lower trunk or the outer crown. Classification and ordination analyses of the species and environmental data indicated the presence of 4 communities: an outer crown, a mid to mid-outer crown, an upper trunk and a lower trunk community. Species distribution on hosts in the present study appeared to be influenced by the intensity of radiation, particularly diffuse radiation, and relative humidity related to a moisture availability gradient. Temperature appeared least important. Six species were selected as potential microclimate change indicators.

Variation in temperature, relative humidity, visible sky fraction, leaf area index and radiation intensity were measured throughout a single canopy a co-dominant *Weinmannia* sp. The opposing gradients of temperature and relative humidity displayed similar fluctuation patterns as found in lowland rainforest, although the range of the gradients was smaller, possibly due to greater atmospheric mixing facilitated by the more permeable canopy at the montane rainforest study site. Microclimate stability decreased with distance above the ground. The lower trunk micro-habitat was the most humid, coolest and least illuminated. The highest temperature and lowest relative humidity were recorded in the mid-outer crown, closely followed by the most irradiated crown periphery. The above canopy temperature remained cooler during the day than the tree crown. The lowest mean hourly temperature was recorded in the mid-crown, induced by rapid

radiative cooling in the evening. The air above the canopy was warmer at night than any micro-habitat on the tree host.

The study found 27 bryophytes species (5 mosses, 22 hepatics) not previously recorded for Peru. Recommendations for methodology in future bryophyte surveys in Peruvian lower montane rainforest are presented. Future studies must parallel more concerted efforts to raise the awareness of the local population about the ecological importance of bryophytes, particularly in cloud forests.

Resumen

Un estudio de briofitas epifitas y de la variación microclimática dentro de la copa de un árbol fueron conducidos en los bosques mas bajos de montaña a 2400 m. en las Yungas peruanas. Un total de 129 especies (87 hepáticas, 1 antocerota y 41 musgos) fueron recolectadas en 3 árboles, usando diferentes metodologías. En un solo Weinmannia sp. se encontró 110 especies (77 hepáticas, 1 antocerota y 32 musgos). Especies con una amplia distribución en los árboles contribuyeron 47% de epifitas, 22% fueron limitados a los troncos y 31% ocurrieron solo en las copas. La totalidad de tasa de hepáticas a musgos fue 2.15:1. Los musgos mostraron un rango de distribución mas estrecho que las hepáticas. La más grande riqueza y abundancia de especies fue encontrada en las ramas grandes de la media copa. Las especies encontradas en la media copa generalmente tuvieron una distribución mas ancha en los árboles que aquellos encontrados en el tronco bajo o en la copa exterior. Los análisis de clasificación y ordenación de las especies y data de medio ambiente indicaron la presencia de 4 comunidades: Copa exterior, de media copa a media copa exterior, parte superior del tronco y la comunidad de la parte baja del tronco. La distribución en los árboles del presente estudio pareció ser influenciada por la intensidad de radiación, particularmente radiación difusa y relativa humedad relacionada a la escala de disponibilidad de agua. La temperatura pareció menos importante. Seis especies fueron seleccionadas como indicadores potenciales del cambio de microclima.

La variación de la temperatura, humedad relativa, fracción visible de cielo, índice de la área foliar e intensidad de radiación fueron medidos en todas partes de un co-dominante *Weinmannia* sp. Las escalas opuestas de temperatura y humedad relativa mostraron similar fluctuación como los encontrados en Selva Baja; sin embargo, el rango de las escalas fue más pequeño, posiblemente a causa de mejor combinación atmosférica facilitada por el dosel más abierto en bosque de montaña elegido para el presente estudio. La estabilidad microclimática disminuyo con la distancia del suelo. El micro hábitat en la parte baja del tronco fue el más húmedo, más fresco y menos iluminado. La temperatura más alta y humedad relativa mas baja fueron registradas en la media copa exterior, cercanamente seguido por la periferia de la copa mas irradiada. La temperatura sobre el dosel permaneció más fresco durante el día que la misma copa del árbol. El promedio de horario de la temperatura mas bajo fue registrado en la media copa, causado por el rápido enfriamiento radiativo en la noche. El aire sobre el dosel fue mas caliente en la noche como en cualquier micro-hábitat en el árbol.

En el estudio se encontró 27 especies de briofitas (5 musgos, 22 hepáticas) que no eran citados en el Perú. Las recomendaciones para metodología en estudios futuros de briofitas en los bosques de montaña baja peruana, son incluidas en el presente informe. Futuros estudios deben suceder paralelos a los esfuerzos concertados para la concientización de la población sobre la importancia ecológica de las briofitas, particularmente en los bosques nublados.

Acknowledgments

This project would not have become reality without the assistance and support of many, both in Tasmania and Peru. This unusual honours project was made possible by the overwhelming support and encouragement of Professor Jamie Kirkpatrick and Dr. Emma Pharo at the School of Geography and Environmental Studies, University of Tasmania.

The present study was inspired by Damien Catchpole and his study of vascular epiphytes in a *Ficus* sp. on the slopes of the Yanachaga-Chemillén Range, Peru. He provided much advice and facilitated many aspects of the study and home life in Oxapampa. Also, a thank you to Karen Richards, whose love of nature has always been and will continue to be an inspiration.

Much logistical support was provided by the Instituto Nacional de Recursos Naturales (INRENA). Co-operation of the staff at the Oxapampa office headed by Eduardo De la Cadena is much appreciated. Access to the facilities of the Missouri Botanical Garden Selva Central branch was invaluable. The staff, particularly Rodolfo Vasquez Martinez, his partner Rocio Rojas and Abel Monteagudo made me feel very welcome. I would also like to thank Percy Summers at Pro-Pachitea and the Fundación Peruana para la Conservación de la Naturaleza (Pro-Naturaleza) for their logistical support. A special thanks to Assoc. Prof Carlos Llerena Pinto who assisted with student visa requirements.

The volume of taxonomic literature that was necessary to complete the present study is staggering. Majority of it had to arrive in electronic format to this corner of the Amazon basin. I would like to thank the University and the library and document delivery staff, for their assistance. Prof. Rob Gradstein provided some key literature, as did Rod Seppelt, who also donated his time to demonstrate some laboratory techniques.

Bryologist are rare anywhere and especially so in Peru. A special thank you to Jasmin Opisso Mejia at the Museo de la Historia Natural in Lima, who gave up much time and provided the expertise necessary to clarify many doubts and verify the moss identifications.

Lupe Huayllacahua Falcon assisted with the abstract translation and provided much needed support during the write up of the present study. And finally, I thank the people of Peru, and especially the residents of Oxapampa, for their warmth and for making this a truly memorable experience.

Table of Contents

Declaration	ii
Abstract	iii
Resumen	iv
Acknowledgments	vi
Table of Contents	vii
List of figures	ix
List of tables	xii
Chapter 1. Introduction	14
1.1 History	16
1.2 Objectives	
1.3 Structure of the thesis	
Chapter 2. Methods	20
2.1 Study area	20
2.2 Methodology review	23
2.3 Minimum subsample study	25
2.3.1 Methods	25
2.3.2 Results	27
2.3.3 Discussion	29
2.3.4. Conclusion	
2.4 Single tree study design and methods	
2.4.1 The host	
2.4.2 Crown stratification and access	
2.4.3 Floristic sampling and identification.	
2.4.4 Temperature and relative humidity	
2.4.5 PAR	35
2.4.6 Statistical analyses	
Chapter 3. Microclimate	
3.1 Background	
3.2 Results	
3.2.1 Temperature and relative humidity	
3.2.2 Radiation	41
3.2.3 Associations	47
3.2.4 Diurnal fluctuation	
3.3 Discussion	53
3.3.1 Habitat stability near the ground	53
3.3.2 The two faces of the trunk	54
3.3.3 Impact of crown architecture and epiphytes on microclimate	54
3.4 Conclusion	56
Chapter 4. Epiphytic bryophytes on Weinmannia sp	
4.1 Background	58
4.2 Results	59
4.2.1 Floristics	59
4.2.2 Species capture and richness estimation	67
4.2.3 Species abundance	68

4.2.4 Microhabitat communities	69
4.2.5 Classification of sites	71
4.2.6 Species classification and ordination	72
4.2.7 Epiphytic communities	76
4.3 Discussion	79
4.3.1 Species richness	79
4.3.2 Hepatic to moss ratio	82
4.3.3 Abundance, competition and disturbance	83
4.3.4 Epiphytic communities	85
4.4 Conclusion	88
Chapter 5. Environmental variables and species distribution	90
5.2 Results	93
5.2.1 Correlation between variables	93
5.2.2 Principal components analysis	94
5.2.3 Non-metric multi-dimensional scaling	95
5.2.4 Species - microhabitat associations	98
5.3 Discussion	100
5.3.1 Height	100
5.3.2 Diameter	103
5.3.3 Why is the exposed side of the trunk so different?	104
5.4 Conclusion	105
Chapter 6. Conclusion	106
6.1 Floristics and community structure	106
6.2 Species richness and sampling methodology	110
6.3 Microclimate and other environmental parameters	112
6.3.1 Microclimate	112
6.3.2 Environmental parameters	114
6.4 Future research needs	115
References	117
Appendix I Taxonomic literature used in species identification (by genus)	128
Appendix II Hourly wet and dry day temperature and humidity fluctuations	136
Appendix III Bryophyte species found on the Weinmannia sp. host	139
Appendix IV Species area curves for Weinmannia zones.	142
Appendix V Species code numbers used in classification and ordination plots	143
Appendix VI Species List	144