

Effects of Shell Abrasion and Aerial Exposure on  
the Performance of Pacific Oysters *Crassostrea gigas*  
(Thunberg, 1793) Cultured in Tasmania,  
Australia.

Colleen Maree O'Meley  
B.Sc. [Hons. (Chem.)]

Submitted in fulfilment of the requirements for the degree of  
Master of Applied Science in Aquaculture by Research

University of Tasmania, November 1995

# Declaration

I declare that this thesis contains no material which has been accepted for the award of any other degree or diploma in any tertiary institution and that to the best of my knowledge and belief, it contains no material previously published or written by any other person, except where due reference is made in the text of the thesis.

This thesis may be made available for loan and limited copying in accordance with the *Copyright Act* 1968.

A handwritten signature in black ink, reading "C. O'Meley". The signature is written in a cursive style with a large, looped 'C' and a stylized 'O'.

Colleen Maree O'Meley  
November, 1995

# Abstract

Two of the major management strategies used by Tasmanian oyster farmers for grow-out of unattached (single-seed) Pacific oysters (*Crassostrea gigas*) are shell abrasion, occurring either deliberately or inadvertently during mechanised grading, and manipulation of intertidal growing height (degree of aerial exposure). Some farmers assert that these strategies can promote faster meat growth, and hence higher condition indices [meat weight relative either to shell cavity volume (CI<sub>vol</sub>), or to shell weight (CI<sub>shell</sub>)]. These reports, however, are anecdotal and have not been substantiated in the literature. The present study was undertaken to evaluate the effects of shell abrasion and aerial exposure on the performance (growth, condition index, shell shape, glycogen content and gonad development) of Pacific oysters cultured in mesh baskets, in two separate experiments, on two commercial leases in Tasmania.

Pacific oysters in Experiment 1 were subjected to the following shell abrasion treatments; one-third were machine-graded twice (MM group), another third were machine-graded once (M group) and the last group were not machine-graded (C group, control). Their performance was measured over 81 d and in 87% of the 77 data sets (eleven performance indices measured on seven sample dates), shell abrasion was not a significant factor ( $P>0.05$ ). It is suggested that this was because the available oysters had little 'shell frill' (fragile shell extensions on the outer margins).

Pacific oysters used in Experiment 2 had large shell frill extensions, prior to being treated as follows; two-thirds of the oysters were initially machine-graded (M group) and one-third was not (C group), and then half of the M group oysters were shaken in their baskets after six weeks, and twelve weeks (MB group) into the experiment. Oysters machine-graded once (M group) lost a mean of  $3.3 \pm 0.4$  mm in shell height and  $5.9 \pm 1.0$  mm in shell length (mean  $\pm$  s.e.;  $n=13$ ). Additional shell frill was removed when oysters were shaken in their baskets (MB group). On the first occasion, the mean shell height and shell length were reduced by  $3.4 \pm 0.5$  mm and  $2.5 \pm 0.4$  mm (mean  $\pm$  s.e.;  $n=29$ ), respectively. When the baskets were shaken again, six weeks later, reductions of  $4.5 \pm 0.6$  mm in shell height and  $2.9 \pm 0.7$  mm in shell length (mean  $\pm$  s.e.;  $n=29$ ) were recorded.

The C group grew faster than the MB group, whilst results for the M group were usually intermediate. By the final sample (124 d) in Experiment 2, the

results ( $P < 0.05$ ) were, for; whole weight ( $\text{g oyster}^{-1}$ )  $C > M, MB$ , shell height  $C > M > MB$ , shell length  $C, M > MB$ , shell depth  $C > M, MB$ , dry shell weight ( $\text{g oyster}^{-1}$ )  $C > M, MB$ , and for dry meat weight ( $\text{g oyster}^{-1}$ )  $C > M$  but the MB group was not significantly different ( $P > 0.05$ ) to the other two. Reduced shell growth relative to meat growth, is one of the major factors influencing condition index; the MB group had a higher mean CIvol than both M and C groups ( $P < 0.05$ ), while the CIvol of the last two groups were not significantly different ( $P > 0.05$ ) by the final sample. The trends in CIshell values were similar, but less pronounced, and by the final sample the mean CIshell values were similar ( $P > 0.05$ ). Shell shape was significantly altered such that the MB group had a higher ( $P < 0.05$ ) mean cup index [= (shell height  $\times$  shell length) $^{0.5}$  / shell depth] but lower ( $P < 0.05$ ) mean roundness index (= shell length / shell height) compared to the other two groups. Throughout the experiment the mean glycogen content did not differ significantly ( $P > 0.05$ ) amongst groups.

The range, for average daily aerial exposure treatments, was much greater in Experiment 1 (0-26% exposure  $\text{d}^{-1}$ ) than in Experiment 2 (0-7% exposure  $\text{d}^{-1}$ ). By the final sample in Experiment 1, the mean whole weight, shell height, shell length, shell depth and dry shell weight of subtidal (0% exposure  $\text{d}^{-1}$ ; L group) oysters were higher ( $P < 0.05$ ) than those held at 26% exposure  $\text{d}^{-1}$  (H group). Because their dry meat weights were similar ( $P > 0.05$ ), the H group developed a higher ( $P < 0.05$ ) mean CIvol and CIshell than the L group. The H group had a higher ( $P < 0.05$ ) mean cup index but lower ( $P < 0.05$ ) mean roundness index compared to the L group, and the mean glycogen content of the H group was higher ( $P < 0.05$ ) than in the L group.

Aerial exposure levels of 0% exposure  $\text{d}^{-1}$  (L group) compared 7% exposure  $\text{d}^{-1}$  (H group) did not significantly affect ( $P > 0.05$ ) the mean whole weight, shell height, shell depth, dry shell weight or dry meat weight indices, although the shell length of the H group was higher ( $P < 0.05$ ) than that of the L group by the final sample in Experiment 2. Compared to the L group, the H group had higher, but not significantly different ( $P > 0.05$ ), mean CIvol and CIshell indices, and were slightly rounder but less cupped in shape. The H group did have a significantly higher ( $P < 0.05$ ) mean glycogen content by the last sample.

results ( $P < 0.05$ ) were, for; whole weight ( $\text{g oyster}^{-1}$ )  $C > M, MB$ , shell height  $C > M > MB$ , shell length  $C, M > MB$ , shell depth  $C > M, MB$ , dry shell weight ( $\text{g oyster}^{-1}$ )  $C > M, MB$ , and for dry meat weight ( $\text{g oyster}^{-1}$ )  $C > M$  but the MB group was not significantly different ( $P > 0.05$ ) to the other two. Reduced shell growth relative to meat growth, is one of the major factors influencing condition index; the MB group had a higher mean CIvol than both M and C groups ( $P < 0.05$ ), while the CIvol of the last two groups were not significantly different ( $P > 0.05$ ) by the final sample. The trends in CIshell values were similar, but less pronounced, and by the final sample the mean CIshell values were similar ( $P > 0.05$ ). Shell shape was significantly altered such that the MB group had a higher ( $P < 0.05$ ) mean cup index [= (shell height  $\times$  shell length) $^{0.5}$ /shell depth] but lower ( $P < 0.05$ ) mean roundness index (= shell length/shell height) compared to the other two groups. Throughout the experiment the mean glycogen content did not differ significantly ( $P > 0.05$ ) amongst groups.

The range, for average daily aerial exposure treatments, was much greater in Experiment 1 (0-26% exposure  $\text{d}^{-1}$ ) than in Experiment 2 (0-7% exposure  $\text{d}^{-1}$ ). By the final sample in Experiment 1, the mean whole weight, shell height, shell length, shell depth and dry shell weight of subtidal (0% exposure  $\text{d}^{-1}$ ; L group) oysters were higher ( $P < 0.05$ ) than those held at 26% exposure  $\text{d}^{-1}$  (H group). Because their dry meat weights were similar ( $P > 0.05$ ), the H group developed a higher ( $P < 0.05$ ) mean CIvol and CIshell than the L group. The H group had a higher ( $P < 0.05$ ) mean cup index but lower ( $P < 0.05$ ) mean roundness index compared to the L group, and the mean glycogen content of the H group was higher ( $P < 0.05$ ) than in the L group.

Aerial exposure levels of 0% exposure  $\text{d}^{-1}$  (L group) compared 7% exposure  $\text{d}^{-1}$  (H group) did not significantly affect ( $P > 0.05$ ) the mean whole weight, shell height, shell depth, dry shell weight or dry meat weight indices, although the shell length of the H group was higher ( $P < 0.05$ ) than that of the L group by the final sample in Experiment 2. Compared to the L group, the H group had higher, but not significantly different ( $P > 0.05$ ), mean CIvol and CIshell indices, and were slightly rounder but less cupped in shape. The H group did have a significantly higher ( $P < 0.05$ ) mean glycogen content by the last sample.

In neither experiment did shell abrasion or aerial exposure have a consistent effect on gonad development, or sex group ratio (male: female: indeterminate).

This study has shown that shell abrasion can retard shell growth, but improve the CIvol, CIshell and cup index for Pacific oysters which have substantial shell frill prior to abrasion. The roundness index and glycogen content, however, were not improved. Increased levels of aerial exposure led to an improved glycogen content, compared to subtidal oysters. Increased levels of exposure will also retard shell growth, but will improve the CIvol, CIshell and cup index, but not the roundness index. As such they are useful management tools but they do not promote faster meat growth.

## Acknowledgements

I wish to thank my supervisor Dr Greg Maguire for his critical guidance, encouragement and patience. My greatest thanks go to him, and Christine Maguire.

I am very grateful to Colin and Sue Dyke, and Peter and Jan Chew who gave me numerous oysters for this study, the use of their leases and equipment, and provided their time and expertise. This study would not have been possible without your support. Thanks also to Hayden Dyke and Tim Mc Mannus for building a second rack height at each lease, the hard work in getting the experiments set up, and the regular temperature monitoring and collection of water samples.

I was awarded two 'starter' grants from the University of Tasmania, and further support was provided by the National Key Centre for Teaching and Research in Aquaculture. Colin and Sue Dyke, and Roger Calvert also provided monetary donations; support greatly appreciated. Colin Sumner, along with the farmers above mentioned also provided critical input.

Special thanks to Adele and Greg Kent, who helped open and measure those thousands of oysters, and also helped me to stay sane while doing it. Stephen Hindrum did the glycogen analyses. Thanks also to Jane Griffith for critical review of the Introduction section of this manuscript.

This thesis is dedicated to my mother Engeliena O'Meley, my sister Serena who has always been by my side, my good friend Deb, and my other friends who luckily are too numerous to mention. Thank you all for your support and encouragement.

# Contents

Abstract	i
Acknowledgements	iv
Contents	v
List of Figures	viii
List of Tables	xi
1. Introduction	1
1.1 Review	1
1.1.1 Production and value of Pacific oysters	1
1.1.2 Taxonomy and distribution	1
1.1.3 Single-seed	3
1.1.4 Site selection	5
1.1.5 Factors affecting marketability of Pacific oysters	9
1.1.6 Effects of culture techniques on the performance of Pacific oysters .	12
1.1.6.1 Stocking density	12
1.1.6.2 Degree of aerial exposure	13
1.1.6.3 Shell abrasion	15
1.2 Summary and aims of this study .	19
2. Materials and Methods	21
2.1 Duration of experiments, sites, culture equipment and experimental oysters	21
2.2 Treatments .	25
2.2.1 Shell abrasion procedures	25
2.2.2 Stocking densities and aerial exposure levels	28
2.3 Sampling regime .	28
2.4 Abiotic measurements	29
2.4.1 Water temperature and salinity	29
2.4.2 Aerial exposure	29
2.5 Biotic measurements	30
2.5.1 Survival .	30
2.5.2 Whole weight and shell growth	30
2.5.3 Meat growth	32
2.5.4 Condition indices	32
2.5.5 Shape indices	33
2.5.6 Glycogen content	33
2.5.7 Gametogenesis	33



<b>2.</b>	<b>Materials and Methods (continued)</b>	
2.5.8	Shelf life . . . . .	34
2.6	Statistical analyses . . . . .	35
<b>3.</b>	<b>Results . . . . .</b>	<b>37</b>
	<b>Experiment 1</b>	
3.1	Abiotic measurements . . . . .	37
3.1.1	Water temperature and salinity . . . . .	37
3.1.2	Aerial exposure . . . . .	37
3.2	Biotic measurements . . . . .	37
3.2.1	Survival . . . . .	37
3.2.2	Growth . . . . .	38
	3.2.2.1 Whole weight and shell growth . . . . .	38
	3.2.2.2 Meat weight . . . . .	39
3.2.3	Condition indices . . . . .	39
	3.2.3.1 Volume condition index . . . . .	39
	3.2.3.2 Shell condition index . . . . .	39
3.2.4	Shape indices . . . . .	40
	3.2.4.1 Roundness index . . . . .	40
	3.2.4.2 Cup index . . . . .	40
3.2.5	Glycogen content . . . . .	40
3.2.6	Gametogenesis . . . . .	41
	<b>Experiment 2</b>	
3.3	Abiotic measurements . . . . .	64
3.3.1	Water temperature and salinity . . . . .	64
3.3.2	Aerial exposure . . . . .	64
3.4	Biotic measurements . . . . .	64
3.4.1	Survival . . . . .	64
3.4.2	Growth . . . . .	65
	3.4.2.1 Whole weight and shell growth . . . . .	65
	3.4.2.2 Meat growth . . . . .	66
3.4.3	Condition indices . . . . .	67
	3.4.3.1 Volume condition index . . . . .	67
	3.4.3.2 Shell condition index . . . . .	67
3.4.4	Shape indices . . . . .	67
	3.4.4.1 Roundness index . . . . .	68
	3.4.4.2 Cup index . . . . .	68
3.4.5	Glycogen content . . . . .	68
3.4.6	Gametogenesis . . . . .	68
3.4.7	Shelf life . . . . .	69

4.1	Discussion	97
4.1	Survival	97
4.2	Whole weight and shell growth	102
4.3	Meat growth	112
4.4	Condition index	114
4.5	Shape index	118
4.6	Glycogen content	122
4.7	Gametogenesis	124
4.8	Shelf life	126
5.	Conclusions	130
	Appendices	133
A	A comparative analysis of the volume condition index (CIvol) and the shell condition index (CIshell)	133
B	Normality of initial and final data in Experiments 1 and 2	141
C	Correlation analyses of ten performance indices measured in Pacific oysters	145
D	Growth in shell length and shell depth of Pacific oysters in Experiments 1 and 2	149
E	Additional results showing the shell dimensions and shape indices of Experiment 2 oysters subjected to shell abrasion treatments	159
F	The effect of shaking baskets of Pacific oysters on their whole weight, shell dimensions and shape indices	163
	References	166

## List of Figures

1.	Location of Experiment 1 and 2 study sites	22
2.	Basket enclosures and racks used in Experiments 1 and 2	23
3.	Pacific oyster with and without shell-frill extensions	24
4.	Grading machine used in Experiment 2	27
5.	External linear shell dimensions	31
6.	Surface water temperature and salinity - Experiment 1	42
7.	Whole weight - Experiment 1	
a.	Shell abrasion and aerial exposure	43
b.	Shell abrasion	44
c.	Aerial exposure	44
8.	Shell height - Experiment 1	
a.	Shell abrasion and aerial exposure	45
b.	Shell abrasion	46
c.	Aerial exposure	46
9.	Dry shell weight - Experiment 1	
a.	Shell abrasion and aerial exposure	47
b.	Shell abrasion	48
c.	Aerial exposure	48
10.	Dry meat weight - Experiment 1	
a.	Shell abrasion and aerial exposure	49
b.	Shell abrasion	50
c.	Aerial exposure	50
11.	Volume condition index (CIvol) - Experiment 1	
a.	Shell abrasion and aerial exposure	51
b.	Shell abrasion	52
c.	Aerial exposure	52
12.	Shell condition index (CIshell) - Experiment 1	
a.	Shell abrasion and aerial exposure	53
b.	Shell abrasion	54
c.	Aerial exposure	54
13.	Roundness index - Experiment 1	
a.	Shell abrasion and aerial exposure	55
b.	Shell abrasion	56
c.	Aerial exposure	56
14.	Cup index - Experiment 1	
a.	Shell abrasion and aerial exposure	57
b.	Shell abrasion	58
c.	Aerial exposure	58

## List of Figures (continued)

15.	Glycogen content - Experiment 1	
a.	Shell abrasion and aerial exposure . . . . .	59
b.	Shell abrasion . . . . .	60
c.	Aerial exposure . . . . .	60
16.	Sex group (i) and gonad stage (ii) - Experiment 1	
a.	Shell abrasion and aerial exposure . . . . .	61
b.	Shell abrasion . . . . .	62
c.	Aerial exposure . . . . .	63
17.	Surface water temperature and salinity - Experiment 2 .	71
18.	Whole weight - Experiment 2	
a.	Shell abrasion and aerial exposure . . . . .	72
b.	Shell abrasion . . . . .	73
c.	Aerial exposure . . . . .	73
19.	Shell height - Experiment 2	
a.	Shell abrasion and aerial exposure . . . . .	74
b.	Shell abrasion . . . . .	75
c.	Aerial exposure . . . . .	75
20.	Dry shell weight - Experiment 2	
a.	Shell abrasion and aerial exposure . . . . .	76
b.	Shell abrasion . . . . .	77
c.	Aerial exposure . . . . .	77
21.	Dry meat weight - Experiment 2	
a.	Shell abrasion and aerial exposure . . . . .	78
b.	Shell abrasion . . . . .	79
c.	Aerial exposure . . . . .	79
22.	Volume condition index (CIvol) - Experiment 2	
a.	Shell abrasion and aerial exposure . . . . .	80
b.	Shell abrasion . . . . .	81
c.	Aerial exposure . . . . .	81
23.	Shell condition index (CIshell) - Experiment 2	
a.	Shell abrasion and aerial exposure . . . . .	82
b.	Shell abrasion . . . . .	83
c.	Aerial exposure . . . . .	83
24.	Roundness index - Experiment 2	
a.	Shell abrasion and aerial exposure . . . . .	84
b.	Shell abrasion . . . . .	85
c.	Aerial exposure . . . . .	85

## List of Figures (continued)

25.	Cup index - Experiment 2		
a.	Shell abrasion and aerial exposure	. . .	86
b.	Shell abrasion	. . .	87
c.	Aerial exposure	. . .	87
26.	Glycogen content - Experiment 2		
a.	Shell abrasion and aerial exposure	. . .	88
b.	Shell abrasion	. . .	89
c.	Aerial exposure	. . .	89
27.	Sex group (i) and gonad stage (ii) - Experiment 2		
a.	Shell abrasion and aerial exposure	. . .	90
b.	Shell abrasion	. . .	91
c.	Aerial exposure	. . .	92
28.	Shelf-life experiment- Cumulative gape - Experiment 2		
a.	Shell abrasion and aerial exposure	. . .	93
b.	Shell abrasion	. . .	94
c.	Aerial exposure	. . .	94
29.	Shelf life experiment - Cumulative mortality - Experiment 2		
a.	Shell abrasion and aerial exposure	. . .	95
b.	Shell abrasion	. . .	96
c.	Aerial exposure	. . .	96

## List of Tables

1.	Summary of Chi-square analysis of gonad stage versus sex of Pacific oysters ( <i>C. gigas</i> ) in Experiment 1 . . . . .	41
2.	Summary of Chi-square analysis of gonad stage versus sex of Pacific oysters ( <i>C. gigas</i> ) in Experiment 2 . . . . .	70
3.	Rough-handling treatments used on Pacific oysters ( <i>C. gigas</i> ) by Spencer et al. (1992) . . . . .	98
4.	Percentage survival after 20 months, of Pacific oysters ( <i>C. gigas</i> ) subjected to agitation in air or water, for 1 or 2 min, dropped onto concrete from 0.5 or 1 m above, and no treatment (control), in the Spencer et al. (1992) study . . . . .	99
5.	Growth rates of Pacific oysters ( <i>C. gigas</i> ) in this study compared to those cultured by Maguire et al. (1994b) . . . . .	104
6.	Comparison of final shell shape indices of Pacific oysters ( <i>C. gigas</i> ) subjected to shell abrasion treatments in this study, to shape indices calculated from data in Robert et al. (1993) and Smith (1994) . . . . .	120