

Physical activity and depression symptom profiles in young men and women with major depression

Charlotte McKercher PhD¹, George C. Patton PhD^{2,3}, Michael D. Schmidt PhD^{1,4},
Alison J. Venn PhD¹, Terence Dwyer MD², Kristy Sanderson PhD¹

1. Menzies Research Institute Tasmania, University of Tasmania, Hobart, Australia

2. Murdoch Childrens Research Institute, Melbourne, Australia

3. University of Melbourne (Dept of Paediatrics)

4. Department of Kinesiology, University of Georgia, Athens, USA

Address correspondence to: Charlotte McKercher

Menzies Research Institute Tasmania

University of Tasmania

Private Bag 23

Hobart TAS 7000

Australia

W: +61 3 6222 7700

F: +61 3 6226 7755

E: Charlotte.McKercher@utas.edu.au

Declaration of interest: None

Acknowledgments

The Childhood Determinants of Adult Health study was funded by the National Health and Medical Research Council (211316), the National Heart Foundation (GOOH 0578), the Tasmanian Community Fund (D0013808), and Veolia Environmental Services. Dr Kristy Sanderson was supported by an Australian Research Council Future Fellowship (FT0991524). We gratefully acknowledge the study sponsors (Sanitarium, ASICS, and Target), and the contributions of the study's project manager, Ms Marita Dalton, all other project staff, volunteers, and study participants.

ABSTRACT

Objective: This study explored whether young adults with major depression who are physically active differ in their depression symptom profile from those physically inactive.

Methods: Analyses included data from 950 (47.6%) men and 1,045 women, (mean [standard deviation] age = 31.5 [2.6] years, participating in a national study.

Participants reported leisure physical activity (International Physical Activity Questionnaire) and ambulatory activity (pedometer steps/day). Diagnosis and symptoms of major depression were assessed using the Composite International Diagnostic Interview.

Results: Prevalence of major depression was 5.5% ($n = 52$) for men and 11.6% ($n = 121$) for women. Interactions between physical activity and sex were observed for depressed mood, appetite changes, vacillating thoughts and suicidality (all, $p < .050$). Among those with major depression, physically active men were significantly less likely to endorse the presence of insomnia (prevalence ratio [PR] = 0.78, 95% confidence interval [CI] = 0.63-0.96), fatigue (PR = 0.82, 95% CI = 0.69-0.99) and suicidality (PR = 0.69, 95% CI = 0.49-0.96) compared with inactive men. Physically active women were significantly less likely to endorse hypersomnia (PR = 0.50, 95% CI = 0.27-0.95), excessive/irrational guilt (PR = 0.76, 95% CI = 0.59-0.97), vacillating thoughts (PR = 0.74, 95% CI = 0.58-0.95) and suicidality (PR = 0.43, 95% CI = 0.20-0.89).

compared with inactive women. Associations were adjusted for age, physical health, educational attainment, depression severity and other depressive symptoms.

Conclusions: Among adults with major depression, those physically active seem to differ in their depression symptom profile from those physically inactive.

Keywords: depression, depressive symptoms, physical activity, population-based

BMI = body mass index; CDAH = Childhood Determinants of Adult Health; DSM-IV = Diagnostic and Statistical Manual of Mental Disorder, 4th edition; IPAQ = International Physical Activity Questionnaire; Short-Form Survey = SF-12v2.

INTRODUCTION

Observational research indicates that regular physical activity is associated with reduced risk of depression and depressive symptoms (1). Intervention studies also indicate beneficial effects of physical activity on depressive symptoms comparable to antidepressants (2). However, previous studies have examined the association between physical activity and global depressive symptoms only. Physical inactivity and depression are both accompanied by changes in psychological and physiological functioning including anhedonia, fatigue, and psychomotor activity. Observed associations may therefore be due to the inclusion of physical inactivity-related characteristics in depression diagnosis. This has implications for an individual's likelihood of endorsing certain diagnostic criterion in order to reach a diagnostic threshold and attain a clinical diagnosis. It is important to clarify therefore whether associations are due to the inclusion of physical inactivity-related characteristics in the diagnostic system or whether those who are physically active are in fact different in their depression symptom profile from those who are physically inactive.

Research examining associations between physical activity and specific depressive symptoms is sparse. Intervention studies indicate that increased exercise may be beneficial in improving depressed mood (3, 4), sleep problems, negative self-worth, and concentration difficulties (4). Further, exercise withdrawal (i.e. induced physical inactivity) has been found to precipitate altered sleep patterns as well as increases in

depressed mood (5, 6) and fatigue (5). Finally, observational studies indicate inverse associations between sports participation (7, 8), physical activity (9, 10), and suicidal behaviour. It remains unclear therefore whether physical inactivity is associated with all depressive symptoms, or whether there is specificity to type of symptom. Clarifying which symptoms co-vary with physical activity may enhance our understanding of the protective mechanisms involved and inform the use of physical activity interventions for depression.

To our knowledge no studies have examined associations between physical activity and individual Diagnostic and Statistical Manual of Mental Disorders (*DSM-IV*) (11) depressive symptoms. Using a population-based cohort we have previously demonstrated inverse associations between physical activity and prevalence of depression in young adults (12). Current analyses examine the prevalence of individual depressive symptoms in physically active and physically inactive young adults diagnosed with major depression. We hypothesized that depressive symptoms corresponding to characteristics of physical inactivity, such as anhedonia, fatigue and psychomotor changes, would differentiate those physically active from those physically inactive.

METHOD

Participants

The Childhood Determinants of Adult Health (CDAH) Study is a 20-year follow-up of 8,498 children (7-15 years) who participated in the 1985 Australian Schools Health and Fitness Survey (13). Sampling procedures and data collection protocols have been described previously (14). Briefly, a nationally representative sample of 109 schools was selected with probability proportional to enrolment. Approximately 20-years later, 6,840 (81%) of the baseline sample were traced and 5,170 (61%) enrolled and provided follow-up data. Of these, 2,410 attended a study clinic and 2,070 completed the main exposure and outcome measures used in the current analysis. Those currently pregnant ($n = 75$) were excluded leaving data from 1,995 clinic attendees. Figure 1 illustrates the recruitment and retention of cohort participants. A subsample of 1,681 clinic attendees (84.3%) additionally completed a 7-day pedometer diary.

Insert Figure 1 here

There were no significant differences at baseline (1985) between those who did and those who didn't provide a pedometer diary at follow-up in depressed mood (8.9% versus 8.2%, $p = .64$) and total past-week physical activity (424 versus 459 minutes/week, $p = .10$). The study was approved by the Southern Tasmania Health

and Medical Human Research Ethics Committee and all participants provided written informed consent.

Measures

Participants were sent questionnaires prior to attending a clinic where height and weight were measured, the Composite International Diagnostic Interview (CIDI) was completed and pedometers were issued.

Physical Activity

Participants completed the International Physical Activity Questionnaire (IPAQ) (15) which assesses the frequency and duration of moderate- and vigorous-intensity leisure physical activity in the previous 7-days. The IPAQ demonstrates very good levels of repeatability and fair to moderate validity when compared to data from accelerometers (16).

Following detailed instructions participants wore a pedometer (Yamax Digiwalker SW-200) for 7-days and kept a pedometer diary. Ambulatory activity (mean steps/day) over the wear period was calculated for each participant with at least four valid days of measurement. Pedometers have shown evidence of reliability (17) and convergent and discriminative validity (18).

Diagnostic assessment

Major depression and depressive symptoms based on *DSM-IV* diagnostic criteria for the previous 12-months were assessed using the CIDI-Auto (version 2.1) (19). The CIDI-Auto includes a list of dichotomous symptom items that separately operationalize the nine *DSM-IV* criteria for major depression. For compound diagnostic criterion, individual symptom components are also assessed. *DSM-IV* organic exclusion and diagnostic hierarchy rules were used in making diagnoses. Diagnosis of major depression requires endorsement of at least one of the two core symptoms of depression (depressed mood, anhedonia), together with at least three associated criteria to give a total of five or more diagnostic criteria. Due to the skip logic embedded in the computerized CIDI-Auto analyses were by necessity restricted to those who endorsed at least one of these core symptoms. These participants were then assessed for the presence of the other seven diagnostic criteria. This subsample who met the diagnostic criteria for a diagnosis of major depression ($n = 173$) comprised the analysis sample.

Sociodemographic and health information

Age, gender, marital status, educational attainment, main source of income, current occupation, smoking status and number of live births (women only) were collected by self-report. Physical health status was assessed using the physical health component summary (PCS) scale of the Short-Form Survey (SF-12v2). Symptom severity was assessed using the mental health component summary (MCS) scale of

the SF-12v2 as a proxy measure, with lower scores indicating increased severity (20).

Body mass index (BMI) (kg/m^2) using objectively measured height and weight was calculated and categorized according to standard BMI cut-points (21).

Statistical analysis

To examine potential reporting bias due to depression, Spearman's correlations were used to assess associations between leisure physical activity and ambulatory activity (steps/day). To identify gender differences in the association between physical activity and depressive symptoms, we calculated interaction terms for participants endorsing each symptom. Similarly, Pearson correlations were used to examine associations between symptom severity (SF-12 MCS) and physical activity.

Differences between physical activity groups (inactive versus active) for each depressive symptom were assessed using χ^2 or Fisher's exact tests as appropriate.

Associations between physical activity and each individual depressive symptom were examined using a two-step approach. Firstly, to compare the prevalence of depressive symptoms between physical activity groups, physical activity was categorized using approximate gender-specific tertile cut-points. Given the skewness of the leisure physical activity variable, categories of leisure physical activity were defined as inactive (0 minutes/week) and active (>0 minutes/week), with inactivity corresponding to the lowest tertile of physical activity. To retain comparability between physical activity variables, categories of ambulatory activity (steps/day)

were similarly defined as inactive (<7,500 steps/day) and active (≥7,500 steps/day) for men, and inactive (<6,800 steps/day) and active (≥6,800 steps/day) for women, with inactivity corresponding to the lowest tertile. For brevity, 'inactive' refers to those with nil leisure physical activity and low ambulatory activity. Using those 'inactive' as the reference group, prevalence ratios (PR) and 95% confidence intervals (CI) for each individual depressive symptom from each physical activity domain were then estimated using Poisson regression. Second, to examine associations between increasing durations of physical activity and each symptom, we calculated β -values and 95% CI using leisure physical activity and ambulatory activity (both log-transformed) as continuous variables. Analyses were conducted by gender due to the known differences in physical activity (22) and depressive aetiology (23) between men and women, as well as overall. Potential covariates were based on a bivariate association ($p < .25$) and retained in the model if the exposure coefficient varied by ≥10% (24). Variables identified *a priori* as potential confounders of the association between physical activity and depression are described previously. Covariates included in the final models are displayed in table footnotes. As individual depressive symptoms may be influenced by the presence of other depressive symptoms (e.g., anhedonia may lead to a loss of appetite), models were additionally adjusted for symptoms identified in univariable analyses as being significantly different between physical activity groups, per a previous analytical approach (25). As different symptoms tend to be endorsed at different levels of depression severity

(26) all models were additionally adjusted for depression severity (number of diagnostic criteria met). Overweight/obesity in women was hypothesized to be a mediating (27) rather than a confounding factor and was examined separately.

Analyses were performed using Stata/IC version 12.0 (Statacorp, 2011) with statistical comparisons treated as significant at $\alpha = .05$ (two-tailed). Due to the exploratory nature of the analysis no adjustments were made for multiple testing (28).

Results

In total, 1,995 participants (950 men, 1,045 women) completed the IPAQ and the depression module of the CIDI. Overall, 5.5% of men ($n = 52$) and 11.6% of women ($n = 121$) met the diagnostic criteria for major depression. For the sub-sample who returned a pedometer diary ($n = 1,681$), prevalence of major depression was 5.9% for men ($n = 47$) and 10.9% for women ($n = 97$). Leisure physical activity and ambulatory activity were significantly associated for both cases ($n = 149$; $r = .18$, $p = .03$) and non-cases ($n = 1,532$; $r = .10$, $p < .001$) suggesting that physical activity reported via the IPAQ was not substantially biased by depression. There were no significant differences in prevalence of major depression between those who did and those who didn't return a pedometer diary for both men ($n = 52$; 5.9% versus 3.2%, $p = .12$) and women ($n = 121$; 10.9% versus 15.2%, $p = .12$).

Characteristics of participants with major depression ($n = 173$) are displayed in Table

1. Mean symptom severity (SF-12 MCS) score was 37.7 (SD = 11.0), which is substantially lower than population norms (20). Regarding depression severity, around half of participants (48.1% men, 47.1% women) met the recommended criterion for severe major depression (i.e., endorsed ≥ 7 diagnostic criteria) (29).

Insert Table 1 here

Table 2 displays interaction terms, Pearson correlations and the proportion of those diagnosed with major depression who endorsed each depressive symptom by leisure physical activity group.

Insert Table 2 here

There were no significant interactions between gender and leisure physical activity however there was non-significant trend for a suicide plan ($\beta = -0.35$, $p = .060$).

Higher levels of symptom severity (SF-12 MCS) were associated with lower activity for depressed mood, fatigue, feelings of worthlessness and suicidality overall and

with appetite increase in men and thoughts of death in women (all $p < .050$). For those reporting no past-week leisure physical activity, a lower proportion of men endorsed anhedonia while a higher proportion of women reported a suicide plan (both $p < .050$). No participants endorsed *attempting* suicide in the past 12-months or worthlessness/guilt due to being impaired by depression.

Table 3 displays interaction terms, Pearson correlations and the proportion of those diagnosed with major depression who endorsed each depressive symptom by ambulatory activity group.

Insert Table 3 here

Significant interactions between gender and ambulatory activity were observed for depressed mood, appetite changes, vacillating thoughts and suicidality (all $p < .050$). Higher symptom severity (SF-12 MCS) was associated with lower ambulatory activity for psychomotor changes overall including agitation in men ($p < .01$) as well as retardation and suicidality in women (both $p < .050$). For participants reporting lower ambulatory activity, a higher proportion endorsed depressed mood, loss of appetite, vacillating thoughts, indecisiveness and suicidality (all $p < .050$). Of note, a

significantly higher proportion of both inactive men and women endorsed having a suicide plan.

Using those physically inactive as the reference group, adjusted PR and standardized β -values with corresponding 95% CI between physical activity domains and individual depressive symptoms were estimated for the analysis sample overall (Table 4) and for men (Table 5) and women (Table 6) separately.

Insert Table 4 here

For all participants with major depression, those reporting any leisure physical activity in the past week were less likely to report concentration difficulties/indecision ($p = .048$) compared with those reporting nil leisure activity. Non-significant trends were observed for objective psychomotor agitation ($p = .060$) and excessive/irrational guilt ($p = .050$). Inverse associations were observed between increasing durations of leisure physical activity and excessive/irrational guilt ($p = .025$) and death/suicidal thoughts ($p = .029$). Those reporting higher ambulatory activity were less likely to endorse fatigue, concentration difficulties/indecision (both $p < .050$) and a suicide plan ($p < .01$) compared with the inactive group. Increasing

durations of ambulatory activity were inversely associated with weight/appetite changes, objective psychomotor agitation (all $p < .050$) and a suicide plan ($p < .001$).

Insert Table 5 here

For men with major depression, those reporting any leisure physical activity were less likely to report insomnia, fatigability, death/suicidal thoughts and thoughts of death compared with those reporting nil leisure activity (all $p < .050$). An inverse association was observed between increasing durations of leisure physical activity and fatigue ($p = .02$) with non-significant trends observed for insomnia ($p = .070$), death/suicidal thoughts ($p = .060$) and thoughts of death ($p = .060$). Those reporting higher ambulatory activity were less likely to endorse fatigability and more likely to endorse worthlessness/guilt than men reporting lower ambulatory activity (both $p < .050$). While non-significant, higher active men were around 80% less likely to endorse a suicide plan compared with low active men ($p = .070$). Likewise, increasing durations of ambulatory activity were inversely associated with having a suicide plan ($p < .01$). Increasing activity was also associated with weight/appetite changes including loss of appetite (both $p < .050$).

Insert Table 6 here

For women with major depression, those reporting participation in some past-week leisure physical activity were less likely to endorse excessive/irrational guilt and a suicide plan compared with those reporting nil leisure activity (both $p < .050$).

Increasing durations of leisure activity were associated with anhedonia ($p = .020$), excessive/irrational guilt ($p = .008$) and having a suicide plan ($p = .015$) Those reporting higher ambulatory activity were significantly less likely to endorse hypersomnia, vacillating thoughts, indecisiveness (all $p < .050$), and a suicide plan ($p < .001$) compared with women reporting lower ambulatory activity. Increasing durations of ambulatory activity were associated with hypersomnia ($p = .020$), objective psychomotor agitation ($p < .050$) and a suicide plan ($p = .001$) with a non-significant trend for vacillating thoughts ($p = .060$).

The addition of overweight/obesity had minimal impact on associations between physical activity domains and depressive symptoms for women (data not shown).

Discussion

Using two different measures of physical activity, we compared the depression symptom profiles of physically active and inactive young men and women with

major depression. Key strengths include multiple physical activity assessment, analysis of depressive symptoms using a validated diagnostic instrument, measured height and weight, and examination of a range of potential confounders. Among those with major depression, active men were less likely to endorse the presence of insomnia, fatigue and suicidality, and more likely to endorse worthlessness/guilt compared with inactive men. Active women were less likely to endorse hypersomnia, excessive/irrational guilt, vacillating thoughts, indecisiveness, and suicidality compared with inactive women. Associations remained significant after adjustment for sociodemographic and health characteristics, depression severity and the influence of other depressive symptoms. Of note, inverse associations between physical activity and suicidality were consistent across gender and physical activity assessment. Increasing durations of physical activity were associated with decreased symptom severity over a range of symptoms including fatigue, sleep disturbance, psychomotor changes, worthlessness/guilt and suicidality.

Due to the inclusion of characteristics related to physical inactivity in the diagnostic system we expected certain depressive symptoms (i.e., anhedonia, fatigue and psychomotor difficulties) to discriminate between inactive and active participants. High endorsement of certain symptoms (e.g., anhedonia, depressed mood), limited our ability to detect significant differences between physical activity groups. Significant associations between physical activity and fatigue were consistently observed across physical activity domains in men however no significant differences

were observed between physical activity and fatigue in women. In addition, no significant associations were observed between physical activity and psychomotor difficulties in men. It may be that a minimum threshold of physical activity intensity is required for physical activity to influence psychomotor changes in young men. Observed gender differences may be reflective of symptom profiles for major depression differing between men and women (26). Further, the physical activity measures utilized may have been insufficiently sensitive to detect significant differences between physical activity groups.

The current analysis suggests that the correspondence in characteristics between physical activity and depression does not fully account for the well-documented association observed in the literature. Indeed, a striking finding was that leisure and ambulatory activity were characterized by decreased endorsement of suicidal symptoms in both men and women with associations remaining significant after adjustment for overweight/obesity in women and depression severity. These findings are consistent with and extend several population-based studies indicating significant inverse associations between sports participation and suicidal ideation in adolescents (8, 30), and physical activity and suicidal behaviour in adolescents (9) and college students (7, 10). A novel finding was that increasing durations of both leisure and ambulatory activity were inversely associated with having a suicide plan in young women. This is in contrast to previous research indicating that moderate to high levels of physical activity were positively associated with risk of suicide

ideation/plans in female college students (7). This finding did not appear to be accounted for by BMI and weight perception leading the authors to speculate that female undergraduate college students (aged 18 years and older) at risk of suicidal behaviour may have been using physical activity as a coping strategy. The older age and community-based nature of the current cohort may account for the different finding.

Several limitations should be considered when interpreting these results. The use of questionnaires may have contributed to imprecision in assessment of leisure physical activity and depressive symptoms. However, the IPAQ has demonstrated acceptable measurement properties (16) and diagnostic interviews for depression, including the CIDI, have been shown to improve standardization of diagnosis and eliminate clinician bias (31). Further, differences in depression prevalence at follow-up were observed in both men and women between those who did and didn't return a pedometer diary. While not statistically significant, the magnitude of these differences may be clinically important and as the direction varied by gender may constitute an important source of bias.

Due to the skip logic embedded in the computerized CIDI-Auto analyses were restricted to those who met the diagnostic threshold for major depression. Future studies should examine differences in depressive symptoms across the population as a whole, not just those meeting the diagnostic threshold. No adjustment was made

for multiple statistical testing, thereby increasing the risk of type I errors (28).

However, correction for multiple testing increases the risk of type II errors (32), and the novelty of this analysis provides some justification. Current findings require replication using a larger cohort with capacity to adjust for multiple comparisons. Future studies should also include a validated depression rating scale to enable a more in-depth examination of associations between physical activity and symptom severity. Finally, the cross-sectional design precludes any conclusions regarding causality.

In summary, young men and women with clinical depression who are physically active appear to have a different depressive symptom profile from those who are physically inactive. The effects of physical activity were apparent across a wide range of depressive symptoms suggesting that observed associations between physical inactivity and depression are not merely due to the epiphenomenon of shared characteristics. Associations were generally consistent across the two analytical approaches lending added assurance to these inferences. Indeed results suggest that physical activity may have particular benefits in reducing levels of sleep disturbance and suicidality. Given that suicide is one of the significant causes of death in young adults worldwide (33) clarifying the efficacy of physical activity in suicide prevention during this critical life stage would be an important advance in public health. Further examination of associations between physical activity and

depressive symptoms within a larger population-based cohort may assist in clarifying these associations.

REFERENCES

1. Teychenne M, Ball K, Salmon J: Associations between physical activity and depressive symptoms in women. *Int J Behav Nutr Phys Act* 2008; 5.
2. Dinas P, Koutedakis Y, Flouris A: Effects of exercise and physical activity on depression. *Ir J Med Sci* 2010;1-7.
3. Nabkasorn C, Miyai N, Sootmongkol A, Junprasert S, Yamamoto H, Arita M, Miyashita K: Effects of physical exercise on depression, neuroendocrine stress hormones and physiological fitness in adolescent females with depressive symptoms. *European Journal of Public Health* 2006; 16:179-184.
4. de Zeeuw EL, Tak EC, Dusseldorp E, Hendriksen IJ: Workplace exercise intervention to prevent depression: A pilot randomized controlled trial. *Mental Health and Physical Activity* 2010; 3:72-77.
5. Glass J, Lyden A, Petzke F, Stein P, Whalen G, Ambrose K, Chrousos G, Clauw D: The effect of brief exercise cessation on pain, fatigue, and mood symptom development in healthy, fit individuals. *J Psychosom Res* 2004; 57:391-398.
6. Berlin A, Kop W, Deuster P: Depressive mood symptoms and fatigue after exercise withdrawal: the potential role of decreased fitness. *Psychosom Med* 2006; 68:224-230.

7. Brown D, Blanton C: Physical activity, sports participation, and suicidal behavior among college students. *Med Sci Sports Exerc* 2002; 34:1087-1096.
8. Taliaferro L, Rienzo B, Miller M, Pigg Jr R, Dodd V: High school youth and suicide risk: Exploring protection afforded through physical activity and sport participation. *J Sch Health* 2008; 78:545-553.
9. Brosnahan J, Steffen LM, Lytle L, Patterson J, Boostrom A: The relation between physical activity and mental health among Hispanic and non-Hispanic white adolescents. *Arch Pediatr Adolesc Med* 2004; 158:818-823.
10. Taliaferro L, Rienzo B, Pigg Jr R, Miller M, Dodd V: Associations between physical activity and reduced rates of hopelessness, depression, and suicidal behavior among college students. *J Am Coll Health* 2009; 57:427-435.
11. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (4th edn) (DSM-IV)*. Washington DC: APA, 1994
12. McKercher C, Schmidt M, Sanderson K, Patton G, Dwyer T, Venn A: Physical activity and depression in young adults. *Am J Prev Med* 2009; 36:161-164.
13. Venn A, Thomson R, Schmidt M, Cleland V, Curry B, Gennat H, Dwyer T: Overweight and obesity from childhood to adulthood: a follow-up of participants in the 1985 Australian Schools Health and Fitness Survey. *Med J Aust* 2007; 186:458-460.

14. Pyke J. Australian Health and Fitness Survey 1985. Adelaide: The Australian Council for Health, Physical Education and Recreation, 1985
15. International Physical Activity Questionnaire. **International Physical Activity Questionnaire** [<http://www.ipaq.ki.se/ipaq.htm>]. 2005 [October 16, 2011].
16. Craig C, Marshall A, Sjöström M, Bauman A, Booth M, Ainsworth B, Pratt M, Ekelund U, Yngve A, Sallis J, Oja P: International Physical Activity Questionnaire: 12-country reliability and validity. Med Sci Sports Exerc 2003; 35:1381-1395.
17. Tryon W, Pinto L, Morrison D: Reliability assessment of pedometer activity measurements. Journal of Psychopathology and Behavioral Assessment 1991; 13:27-44.
18. Tudor-Locke C, Myers A: Methodological considerations for researchers and practitioners using pedometers to measure physical (ambulatory) activity. Res Q Exerc Sport 2001; 72:1-12.
19. World Health Organization. Composite International Diagnostic Interview (CIDI). Version 2.1. Geneva: WHO, 1997
20. Ware J, Kosinski M, Keller S: A 12-item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. Med Care 1996; 34:220-233.

21. National Heart Lung and Blood Institute. Clinical Guidelines for the Identification, Evaluation and Treatment of Overweight and Obesity in Adults: The Evidence Report. Bethesda, MD: National Heart, Lung, and Blood Institute, 1998
22. Caspersen C, Pereira M, Curran K: Changes in physical activity patterns in the United States, by sex and cross-sectional age. *Med Sci Sports Exerc* 2000; 32:1601-1609.
23. Kuehner C: Gender differences in unipolar depression: an update of epidemiological findings and possible explanations. *Acta Psychiatr Scand* 2003; 108:163-174.
24. Hosmer D, Lemeshow S. *Applied logistic regression*, 2nd edition. New York: Wiley & Sons, 2000.
25. Gaudiano B, Young D, Chelminski I, Zimmerman M: Depressive symptom profiles and severity patterns in outpatients with psychotic vs nonpsychotic major depression. *Compr Psychiatry* 2008; 49:421-429.
26. Carragher N, Mewton L, Slade T, Teesson M: An item response analysis of the DSM-IV criteria for major depression: Findings from the Australian National Survey of Mental Health and Wellbeing. *J Affect Disord* 2011; 130:92-98.

27. Sanderson K, Patton G, McKercher C, Dwyer T, Venn A: Overweight and obesity in childhood and risk of mental disorder: a 20-year cohort study. *Aust N Z J Psychiatry* 2011; 45:384-392.
28. Rothman KJ: No adjustments are needed for multiple comparisons. *Epidemiology* 1990; 1:43-46.
29. Kessler R, Zhao S, Blazer D, Swartz M: Prevalence, correlates, and course of minor depression and major depression in the National Comorbidity Survey. *J Affect Disord* 1997; 45:19-30.
30. Brown D, Galuska D, Zhang J, Eaton D, Fulton J, Lowry R, Maynard L: Physical activity, sport participation, and suicidal behavior: U.S. high school students. *Med Sci Sports Exerc* 2007; 39:2248-2257.
31. Erdman H, Klein M, Greist J: Direct patient computer interviewing. *J Consult Clin Psychol* 1985; 53:760-773.
32. Perneger T: What's wrong with Bonferroni adjustments. *Br Med J (Clin Res Ed)* 1998; 316:1236-1238.
33. Murray C, Lopez A: Mortality by cause for eight regions of the world: Global Burden of Disease Study. *Lancet* 1997; 349:1269-1276.

Figure 1. Flowchart of recruitment and retention of cohort participants

TABLE 1. Characteristics of participants with major depression

	Overall	Men	Women
	<i>n</i> = 173	<i>n</i> = 52	<i>n</i> = 121
Age, M (SD), y	31.4 (2.6)	31.5 (2.9)	31.4 (2.5)
Educational attainment, <i>n</i> (%)			
University	67 (38.7)	17 (32.7)	50 (41.3)
Diploma/vocational	51 (29.5)	20 (38.5)	31 (25.6)
Year 12 or less	55 (31.8)	15 (28.8)	40 (33.1)
Short-Form Survey, version 2, M (SD)			
Mental Health Component Summary	37.7 (11.0)	38.2 (11.8)	37.6 (10.6)
Physical Health Component Summary	54.4 (9.3)	55.6 (8.2)	53.8 (9.7)
Missing, <i>n</i>	5	2	3
Overweight/obese, <i>n</i> (%)	35 (20.2)	10 (19.2)	25 (20.7)
Depression severity ^a , M (SD)	6.54 (1.24)	6.56 (1.24)	6.54 (1.25)
Leisure physical activity, M (SD), minutes/week	130 (199)	163 (269)	115 (158)
Inactive, <i>n</i> (%)	53 (30.6)	39 (32.2)	14 (26.9)
Ambulatory activity, M (SD), steps/day	8223 (2983)	8843 (3344)	7923 (2761)
Inactive, <i>n</i> (%)	54 (37.5)	36 (37.1)	18 (38.3)
Missing, <i>n</i>	29	5	24

M, mean; SE, standard deviation

^aNumber of diagnostic criteria met.

TABLE 2. Descriptive statistics for depressive symptoms and leisure physical activity

Symptoms	Overall				Men			Women		
			Inactive	Active			Inactive	Active	Inactive	Active
	<i>n</i> = 173	<i>n</i> = 173	<i>n</i> = 53	<i>n</i> = 120	<i>n</i> = 52	<i>n</i> = 14	<i>n</i> = 38	<i>n</i> = 121	<i>n</i> = 39	<i>n</i> = 82
	β -value ^a	<i>r</i> ^b	%	%	<i>r</i> ^b	%	%	<i>r</i> ^b	%	%
Depressed mood	0.02	0.19*	88.7	90.8	0.22	92.9	89.5	0.17	87.2	91.5
Loss of interest	-0.02	0.15	86.8	87.5	0.28	71.4	94.7*	0.09	92.3	84.2
Weight/appetite changes	-0.02	0.13	81.1	76.7	0.11	71.4	73.7	0.13	84.6	78.1
Loss of appetite	-0.13	0.08	56.6	50.0	-0.18	50.0	63.2	0.21	59.0	43.9
Increase in appetite	0.13	0.21	35.9	34.2	0.64*	28.6	15.8	0.10	38.5	42.7
Weight loss	0.16	-0.01	24.5	30.0	-0.13	14.3	23.7	0.04	28.2	32.9
Weight gain	0.14	0.31*	24.5	25.8	0.66	14.3	15.8	0.20	28.2	30.5
Sleep disturbance	-0.05	0.12	94.3	87.5	0.18	100.0	92.1	0.08	92.3	85.4
Insomnia	-0.06	0.09	88.7	79.2	0.17	100.0	81.6	0.04	84.6	78.1
Hypersomnia	-0.01	0.17	37.7	39.2	0.01	21.4	42.1	0.22	43.6	37.8
Psychomotor difficulties	0.03	0.01	26.4	31.7	-0.12	21.4	36.8	-0.03	28.2	29.3
Psychomotor retardation	0.05	0.18	32.1	38.3	0.18	21.4	42.1	0.14	35.9	36.6
Objective psychomotor retardation	-0.05	0.13	20.8	22.5	0.02	14.3	29.0	0.03	23.1	19.5
Psychomotor agitation	0.14	-0.05	18.9	28.3	0.27	21.4	31.6	-0.25	18.0	26.8

Objective psychomotor agitation	0.38	-0.17	9.4	19.2	-0.10	7.1	15.8	-0.22	10.3	20.7
Fatigability	-0.01	0.18*	90.6	87.5	0.18	100.0	86.8	0.18	87.2	87.8
Worthlessness/guilt	0.15	0.1	37.7	46.7	0.15	28.6	39.5	0.09	41.0	50.0
Feelings of worthlessness	-0.02	0.21*	66.0	64.2	0.33	57.1	65.8	0.14	69.2	63.4
Excessive/irrational guilt	-0.08	0.11	73.6	61.7	0.21	64.3	60.5	0.06	76.9	62.2
Concentration difficulties/indecision	-0.01	0.14	100.0	95.8	0.17	100.0	92.1	0.12	100.0	97.6
Vacillating thoughts	0.03	0.09	73.6	77.5	0.08	85.7	76.3	0.09	69.2	78.1
Indecisiveness	-0.04	0.08	84.9	75.0	0.08	85.7	68.4	0.07	84.6	78.1
Death/suicidal thoughts	-0.08	0.24*	54.7	47.5	0.22	71.4	50.0	0.25	48.7	46.3
Thoughts of death	0.02	0.26*	50.9	50.0	0.19	71.4	44.7	0.31*	43.6	52.4
Suicidal ideation	-0.11	0.24	32.1	27.5	0.49	21.4	31.6	0.17	35.9	25.6
Suicide plan	-0.35	0.18	22.6	14.2	0.43	14.3	21.1	0.12	25.6	11.0*

^aβ-values for interaction between gender (men = 0, women = 1) and leisure physical activity (inactive = 0, active = 1).

^bPearson correlations between symptom severity (SF-12 Mental Component Summary) and leisure physical activity.

* $p < 0.05$

Leisure physical activity; inactive (0 minutes/week), active (>0 minutes/week).

TABLE 3. Descriptive statistics for depressive symptoms and ambulatory activity

Symptoms	Overall				Men			Women		
			Inactive	Active			Inactive	Active	Inactive	Active
	<i>n</i> = 144	<i>n</i> = 144	<i>n</i> = 54	<i>n</i> = 90	<i>n</i> = 47	<i>n</i> = 18	<i>n</i> = 29	<i>n</i> = 97	<i>n</i> = 36	<i>n</i> = 61
	β -value ^a	<i>r</i> ^b	%	%	<i>r</i> ^b	%	%	<i>r</i> ^b	%	%
Depressed mood	-0.08*	0.11	96.3	84.4*	0.09	94.4	89.7	0.10	97.2	82.0*
Loss of interest	0.05	0.17	83.3	90.0	0.17	88.9	86.2	0.18	80.6	91.8
Weight/appetite changes	-0.01	0.08	81.5	75.6	0.03	88.9	65.5	0.11	77.8	80.3
Loss of appetite	-0.22*	0.11	61.1	43.3*	-0.10	72.2	51.7	0.31	55.6	39.3
Increase in appetite	0.29*	-0.03	29.6	38.9	0.23	22.2	17.2	-0.09	33.3	49.2
Weight loss	0.05	0.28	24.1	22.2	0.08	27.8	13.8	0.38	22.2	26.2
Weight gain	0.19	0.10	22.2	25.6	-0.05	16.7	13.8	0.13	25.0	31.2
Sleep disturbance	-0.002	0.14	87.0	88.9	0.10	94.4	93.1	0.16	83.3	86.9
Insomnia	-0.01	0.12	81.5	80.0	0.07	94.4	79.3	0.15	75.0	80.3
Hypersomnia	-0.11	0.19	40.7	33.3	0.11	38.9	34.5	0.21	41.7	32.8
Psychomotor difficulties	-0.09	0.36*	31.5	27.8	0.32	38.9	31.0	0.35	27.8	26.2
Psychomotor retardation	-0.11	0.39**	44.4	34.4	0.46	50.0	31.0	0.36*	41.7	36.1
Objective psychomotor retardation	-0.18	0.39*	27.8	18.9	0.42	38.9	17.2	0.42	22.2	19.7
Psychomotor agitation	-0.13	0.43**	29.6	24.4	0.69**	38.9	27.6	0.20	25.0	23.0

Objective psychomotor agitation	0.11	0.50*	13.0	16.7	0.71	11.1	17.2	0.37	13.9	16.4
Fatigability	-0.05	0.12	94.4	84.4	0.14	100.0	82.8	0.11	91.7	85.3
Worthlessness/guilt	-0.001	0.06	42.6	43.3	-0.08	22.2	44.8	0.13	52.8	42.6
Feelings of worthlessness	-0.05	0.10	68.5	61.1	0.08	77.8	58.6	0.12	63.9	62.3
Excessive/irrational guilt	0.00	0.17	64.8	67.8	0.11	50.0	72.4	0.19	72.2	65.6
Concentration difficulties/indecision	-0.02	0.13	100.0	94.4	0.10	100.0	89.7	0.14	100.0	96.7
Vacillating thoughts	-0.13*	0.08	85.2	68.9*	0.12	77.8	75.9	0.03	88.9	65.6*
Indecisiveness	-0.08	0.13	87.0	72.2*	0.04	77.8	69.0	0.17	91.7	73.8*
Death/suicidal thoughts	-0.25*	0.12	59.3	37.8*	-0.08	72.2	41.4*	0.33*	52.8	36.1
Thoughts of death	-0.14	0.14	55.6	41.1	-0.09	66.7	37.9	0.35*	50.0	42.6
Suicidal ideation	-0.34*	0.07	37.0	18.9*	0.03	38.9	17.2	0.13	36.1	19.7
Suicide plan	-1.21**	0.04	31.5	4.4***	-0.07	33.3	6.9*	0.26	30.6	3.3***

^aβ-values for interaction between gender (men = 0, women = 1) and ambulatory activity (steps/day) (inactive = 0, active = 1).

^bPearson correlations between symptom severity (SF-12 Mental Component Summary) and ambulatory activity.

* $p < .05$, ** $p < .01$, *** $p < .001$

Ambulatory activity; physically inactive (<7,500 steps/day) and physically active (≥7,500 steps/day) for men, physically inactive (<6,800 steps/day) and physically active (≥6,800 steps/day) for women.

TABLE 4. Associations between physical activity and depressive symptoms for participants with major depression^a

Symptoms	Leisure physical activity (n = 168)			Ambulatory activity (n = 139)		
	%	PR (95% CI) ^b	β-value (95% CI) ^b	%	PR (95% CI) ^b	β-value (95% CI) ^b
Depressed mood	90.5	1.05 (0.94-1.17)	0.01 (-0.01-0.03)	89.2	0.92 (0.84-1.01)	-0.10 (-0.21-0.002)
Loss of interest	88.1	1.00 (0.90-1.12)	-0.01 (-0.03-0.01)	88.5	1.09 (0.96-1.24)	0.10 (-0.10-0.26)
Weight/appetite changes	77.4	0.95 (0.81-1.11)	-0.01 (-0.04-0.02)	77.0	0.99 (0.82-1.18)	-0.23 (-0.42--0.04)*
Loss of appetite	50.6	0.86 (0.65-1.15)	-0.03 (-0.09-0.02)	48.2	0.76 (0.53-1.08)	-0.49 (-0.90--0.08)*
Increase in appetite	35.7	0.98 (0.64-1.51)	0.004 (-0.08-0.09)	36.7	1.59 (0.99-2.54)	0.25 (-0.41-0.92)
Weight loss	28.6	1.22 (0.73-2.10)	0.04 (-0.06-0.14)	23.0	1.00 (0.55-1.82)	-0.05 (-0.81-0.72)
Weight gain	26.2	1.16 (0.68-1.98)	0.04 (-0.07-0.14)	22.2	1.60 (0.89-2.85)	0.24 (-0.60-1.08)
Sleep disturbance	89.3	0.95 (0.87-1.04)	-0.01 (-0.03-0.01)	87.8	1.04 (0.92-1.18)	-0.06 (-0.17-0.05)
Insomnia	81.6	0.90 (0.79-1.03)	-0.02 (-0.05-0.01)	79.9	1.03 (0.86-1.22)	-0.04 (-0.21-0.13)
Hypersomnia	38.1	1.13 (0.75-1.72)	0.03 (-0.05-0.11)	35.3	0.74 (0.45-1.23)	-0.44 (-1.01-0.13)
Psychomotor difficulties	29.8	1.34 (0.86-2.10)	0.10 (-0.02-0.14)	28.8	1.18 (0.71-1.97)	0.47 (-0.17-1.12)
Psychomotor retardation	35.7	1.25 (0.86-1.83)	0.03 (-0.04-0.10)	37.4	1.05 (0.70-1.56)	0.13 (-0.43-0.68)
Objective psychomotor retardation	21.4	1.17 (0.69-1.97)	0.04 (-0.06-0.14)	21.6	0.82 (0.42-1.58)	-0.06 (-0.75-0.64)
Psychomotor agitation	25.0	1.57 (0.83-2.99)	0.06 (-0.05-0.18)	25.9	0.95 (0.52-1.73)	0.71 (-0.02-1.44)
Objective psychomotor agitation	16.1	2.36 (0.97-5.76)	0.12 (-0.02-0.26)	15.1	2.03 (0.83-4.96)	1.47 (0.22 to 2.72)*
Fatigability	88.1	0.97 (0.86-1.08)	-0.01 (-0.03-0.01)	87.8	0.87 (0.78-0.98)*	-0.10 (-0.23-0.03)

Worthlessness/guilt	44.1	1.26 (0.87-1.82)	0.05 (-0.02-0.12)	43.2	0.97 (0.64-1.46)	0.11 (-0.41-0.63)
Feelings of worthlessness	64.3	1.00 (0.80-1.28)	-0.02 (-0.06-0.03)	63.3	0.99 (0.78-1.29)	-0.07 (-0.37-0.22)
Excessive/irrational guilt	64.9	0.81 (0.66-1.00)	-0.05 (-0.09--0.01)*	66.2	1.03 (0.80-1.32)	0.07 (-0.24-0.38)
Concentration difficulties/indecision	97.6	0.96 (0.92-0.99)*	-0.003 (-0.008-0.002)	97.1	0.94 (0.89-0.99)*	-0.06 (-0.13-0.02)
Vacillating thoughts	76.2	1.07 (0.89-1.29)	0.01 (-0.02-0.05)	74.8	0.86 (0.72-1.04)	-0.10 (-0.30-0.11)
Indecisiveness	78.0	0.90 (0.77-1.05)	-0.02 (-0.05-0.01)	77.7	0.87 (0.73-1.03)	-0.16 (-0.35-0.03)
Death/Suicidal thoughts	49.4	0.81 (0.63-1.05)	-0.06 (-0.11--0.01)*	45.3	0.83 (0.61-1.14)	-0.13 (-0.55-0.28)
Thoughts of death	50.0	0.92 (0.72-1.19)	-0.03 (-0.08-0.02)	46.0	0.95 (0.70-1.29)	0.01 (-0.40-0.42)
Suicidal ideation	29.2	0.92 (0.58-1.45)	-0.04 (-0.12-0.05)	25.9	0.73 (0.42-1.27)	-0.19 (-0.90-0.52)
Suicide plan	16.7	0.60 (0.32-1.14)	-0.11 (-0.24-0.02)	14.4	0.16 (0.05-0.51)**	-1.21 (-1.80--0.62)***

^aPrevalence ratios (PR) and 95% confidence intervals (CI) calculated using those inactive as the reference group. β -values and 95% CI calculated using physical activity (log-transformed) as a continuous variable.

^bPR and β -values with corresponding 95% CI for each symptom calculated using Poisson regression models adjusted for age, physical health, educational attainment, depression severity and other symptoms.

* $p < .05$, ** $p < .01$, *** $p < .001$

TABLE 5. Associations between physical activity and depressive symptoms for men with major depression^a

Symptoms	Leisure physical activity (n = 50)			Ambulatory activity (n = 45)		
	%	PR (95% CI) ^b	β-value (95% CI) ^b	%	PR (95% CI) ^b	β-value (95% CI) ^b
Depressed mood	92.0	1.06 (0.88-1.27)	0.006 (-0.03-0.04)	93.3	1.04 (0.86-1.26)	-0.04 (-0.20-0.12)
Loss of interest	88.0	1.30 (0.94-1.80)	0.03 (-0.03-0.09)	86.7	1.08 (0.82-1.42)	0.15 (-0.14-0.45)
Weight/appetite changes	72.0	1.09 (0.78-1.51)	0.01 (-0.05-0.07)	73.3	0.79 (0.57-1.10)	-0.49 (-0.86--0.10)*
Loss of appetite	58.0	1.28 (0.74-2.20)	0.004 (-0.09-0.10)	57.8	0.75 (0.44-1.27)	-0.64 (-1.14--0.14)*
Increase in appetite	20.0	0.54 (0.15-1.98)	-0.001 (-0.29-0.29)	20.0	1.18 (0.34-4.11)	-0.05 (-1.22-1.12)
Weight loss	20.0	0.83 (0.20-3.40)	-0.06 (-0.29-0.17)	17.8	0.64 (0.10-3.94)	0.38 (-1.37-2.13)
Weight gain	16.0	1.02 (0.25-4.16)	0.12 (-0.23-0.47)	15.6	1.63 (0.39-6.69)	0.09 (-1.21-1.39)
Sleep disturbance	94.0	0.88 (0.75-1.03)	-0.02 (-0.05-0.01)	93.3	1.03 (0.83-1.27)	0.06 (-0.11-0.22)
Insomnia	86.0	0.78 (0.63-0.96)*	-0.04 (-0.08-0.002)	84.4	0.85 (0.64-1.14)	-0.12 (-0.38-0.15)
Hypersomnia	38.0	1.90 (0.59-6.11)	0.09 (-0.09-0.27)	37.8	1.16 (0.51-2.65)	0.45 (-0.59-1.49)
Psychomotor difficulties	32.0	1.45 (0.47-4.48)	0.05 (-0.13-0.24)	33.3	0.74 (0.30-1.83)	0.63 (-0.77-2.02)
Psychomotor retardation	34.0	1.79 (0.63-5.10)	0.08 (-0.08-0.25)	35.6	0.65 (0.27-1.55)	0.28 (-1.35-0.78)
Objective psychomotor retardation	24.0	2.03 (0.58-7.11)	0.15 (-0.06-0.37)	24.4	0.42 (0.09-1.90)	-0.08 (-1.70-1.53)
Psychomotor agitation	30.0	1.03 (0.34-3.14)	-0.01 (-0.21-0.02)	33.3	0.64 (0.27-1.51)	0.48 (-0.80-1.76)
Objective psychomotor agitation	14.0	1.20 (0.14-10.70)	-0.05 (-0.34-0.24)	15.6	0.56 (0.11-2.98)	0.49 (-2.14-3.12)
Fatigability	90.0	0.82 (0.69-0.99)*	-0.05 (-0.10--0.01)*	88.9	0.83 (0.69-0.99)*	-0.20 (-0.43-0.04)

Worthlessness/guilt	36.0	1.86 (0.74-4.67)	0.12 (-0.03-0.27)	35.6	2.66 (1.11-6.35)*	0.81 (-0.67-2.29)
Feelings of worthlessness	64.0	1.17 (0.70-1.95)	0.03 (-0.06-0.12)	66.7	0.86 (0.54-1.35)	-0.29 (-0.70-0.12)
Excessive/irrational guilt	60.0	0.87 (0.53-1.42)	-0.02 (-0.12-0.07)	62.2	1.68 (0.95-2.98)	0.42 (-0.30-1.14)
Concentration difficulties/indecision	96.0	0.97 (0.91-1.03)	0.002 (-0.007-0.01)	95.6	0.93 (0.85-1.03)	-0.08 (-0.22-0.05)
Vacillating thoughts	80.0	0.99 (0.74-1.33)	0.01 (-0.03-0.06)	77.8	0.99 (0.71,1.37)	-0.002 (-0.34-0.34)
Indecisiveness	74.0	0.82 (0.58-1.16)	-0.02 (-0.08-0.04)	73.3	0.96 (0.64,1.43)	-0.21 (-0.58-0.17)
Death/Suicidal thoughts	56.0	0.69 (0.49-0.96)*	-0.07 (-0.15-0.004)	53.3	0.87 (0.56,1.35)	-0.23 (-0.72-0.27)
Thoughts of death	52.0	0.63 (0.43-0.92)*	-0.08 (-0.16-0.002)	48.9	0.86 (0.53-1.40)	-0.34 (-0.83-0.15)
Suicidal ideation	28.0	1.04 (0.35-3.10)	-0.02 (-0.21-0.17)	24.4	0.63 (0.20-2.00)	-0.31 (-1.79-1.16)
Suicide plan	18.0	0.90 (0.43-0.92)	-0.05 (-0.27-0.17)	15.6	0.16 (0.02-1.16)	-1.35 (-2.25--0.46)**

^aPrevalence ratios (PR) and 95% confidence intervals (CI) calculated using those inactive as the reference group. β -values and 95% CI calculated using physical activity (log-transformed) as a continuous variable.

^bPR and β -values with corresponding 95% CI for each symptom calculated using Poisson regression models adjusted for age, physical health, depression severity and other symptoms.

* $p < .05$, ** $p < .01$,

TABLE 6. Associations between physical activity and depressive symptoms for women with major depression^a

Symptoms	Leisure physical activity (n = 118)			Ambulatory activity (n = 94)		
	%	PR (95% CI) ^b	β-value (95% CI) ^b	%	PR (95% CI) ^b	β-value ^b
Depressed mood	89.8	1.10 (0.96-1.26)	0.03 (-0.001-0.05)	87.2	0.91 (0.78-1.06)	-0.08 (-0.21-0.06)
Loss of interest	88.1	0.89 (0.78-1.01)	-0.03 (-0.07--0.01)*	89.4	1.15 (0.93-1.42)	0.12 (-0.10-0.34)
Weight/appetite changes	79.7	0.92 (0.77-1.11)	-0.01 (-0.05 to 0.03)	78.7	1.07 (0.93-1.42)	-0.12 (-0.40-0.17)
Loss of appetite	32.2	0.73 (0.51-1.06)	-0.06 (-0.14-0.02)	43.6	0.82 (0.46-1.47)	-0.39 (-1.07-0.28)
Increase in appetite	42.4	1.12 (0.71-1.77)	0.01 (-0.08-0.10)	44.7	1.67 (0.91-3.07)	0.42 (-0.38-1.21)
Weight loss	47.5	1.25 (0.70-2.24)	0.04 (-0.07-0.15)	25.5	2.47 (0.87-6.96)	0.40 (-0.57-1.37)
Weight gain	30.5	1.14 (0.65-2.01)	0.02 (-0.09-0.12)	29.8	1.18 (0.60-2.34)	0.10 (-0.70-0.91)
Sleep disturbance	87.3	0.96 (0.85-1.09)	-0.01 (-0.03-0.01)	85.1	1.06 (0.85-1.32)	-0.16 (-0.34-0.02)
Insomnia	79.7	0.95 (0.78-1.15)	-0.02 (-0.06-0.02)	77.7	1.27 (0.92-1.75)	0.04 (-0.23-0.32)
Hypersomnia	38.1	0.90 (0.56-1.44)	-0.02 (-0.12-0.08)	34.0	0.50 (0.27-0.95)*	-0.89 (-1.66--0.13)*
Psychomotor difficulties	28.8	1.12 (0.66-1.89)	0.03 (-0.08-0.14)	26.6	1.47 (0.62-3.50)	0.25 (0.44-0.95)
Psychomotor retardation	36.4	1.10 (0.71-1.71)	0.01 (-0.08-0.10)	38.3	1.01 (0.55-1.85)	0.13 (-0.50-0.77)
Objective psychomotor retardation	20.3	0.92 (0.48-1.75)	-0.02 (-0.16-0.11)	20.2	1.03 (0.42-2.53)	-0.08 (0.91 to 0.75)
Psychomotor agitation	22.9	1.63 (0.73-3.62)	0.09 (-0.07-0.24)	22.3	1.89 (0.76-4.75)	1.03 (-0.08-2.13)
Objective psychomotor agitation	16.9	2.31 (0.83-6.43)	0.17 (-0.02-0.36)	14.9	3.95 (0.88-17.75)	1.11 (0.01 to 2.21)*
Fatigability	87.3	0.99 (0.86-1.16)	0.02 (-0.03-0.03)	87.2	0.87 (0.74-1.01)	-0.03 (-0.16-0.11)

Worthlessness/guilt	47.5	1.30 (0.85-2.01)	0.05 (-0.04-0.13)	46.8	0.75 (0.45-1.26)	0.09 (-0.50-0.68)
Feelings of worthlessness	64.4	1.03 (0.78-1.36)	-0.02 (-0.08-0.04)	61.7	1.35 (0.88-2.07)	0.26 (-0.23-0.75)
Excessive/irrational guilt	66.9	0.76 (0.59-0.97)*	-0.07 (-0.12--0.02)**	68.1	0.80 (0.59-1.08)	-0.04 (-0.36-0.28)
Concentration difficulties/indecision	98.3	0.97 (0.93-1.01)	-0.001 (-0.01-0.003)	97.9	0.96 (0.90-1.02)	-0.02 (-0.10-0.03)
Vacillating thoughts	74.6	1.16 (0.92-1.47)	0.02 (-0.03-0.07)	73.4	0.74 (0.58-0.95)*	-0.25 (-0.52-0.01)
Indecisiveness	79.7	0.92 (0.77-1.10)	-0.02 (-0.06-0.02)	79.8	0.80 (0.65-0.99)*	-0.12 (-0.36-0.12)
Death/Suicidal thoughts	46.6	0.88 (0.62-1.23)	-0.06 (-0.13-0.02)	41.5	0.79 (0.50-1.25)	-0.10 (-0.74-0.54)
Thoughts of death	49.2	1.14 (0.80-1.61)	0.002 (-0.07-0.07)	44.7	0.99 (0.66-1.50)	0.24 (-0.33-0.81)
Suicidal ideation	29.7	0.83 (0.50-1.37)	-0.07 (-0.17-0.03)	26.6	0.77 (0.40-1.49)	-0.17 (-0.97-0.64)
Suicide plan	16.1	0.43 (0.20-0.89)*	-0.21 (-0.37--0.04)*	13.8	0.09 (0.03-0.28)**	-1.67 (-2.67--0.66)**

^aPrevalence ratios (PR) and 95% confidence intervals (CI) calculated using those inactive as the reference group. β -values and 95% CI calculated using physical activity (log-transformed) as a continuous variable.

^bPR and β -values with corresponding 95% CI for each symptom calculated using Poisson regression models adjusted for age, physical health, educational attainment, depression severity and other symptoms.

* $p < .05$, ** $p < .01$