

REPORT CARD

Southern Ocean Acidification



ANTARCTIC CLIMATE
& ECOSYSTEMS CRC



PHOTO 2

ACE CRC Report Card

Southern Ocean Acidification

Introduction

There is now a clear scientific consensus that the increasing volume and rate of our carbon dioxide (CO₂) emissions are causing rapid and unprecedented changes in our oceans. These will have potentially serious impacts within the 21st century, for the sustainability and management of many marine and coastal ecosystems.

Ocean acidification science is rapidly evolving. Since the ACE CRC Position Analysis on ocean acidification was released in 2008¹ significant advances in knowledge have been achieved, including by ACE CRC researchers in the Southern Ocean.

This cold-water coral reef was recently discovered 800m below the ocean surface on the edge of the Antarctic continental slope. The corals are comprised of aragonite and will be amongst the first organisms to disappear if ocean acidification increases.

The aims of this document are:

1. To update the Australian Government on the latest developments in ocean acidification scientific research, highlighting the ACE CRC's efforts and niche in this rapidly expanding research arena since the first Position Analysis;
2. To outline the likely impacts of increased carbon dioxide absorption on the Southern Ocean in particular; and
3. To identify issues for Australia's consideration in future policy development.

Ocean acidification 101

Ocean acidification is the name given to the ongoing decrease in the pH of the world's oceans, caused by their uptake of anthropogenic carbon dioxide from the atmosphere.






The term pH is short for potentiometric hydrogen ion concentration, ie the concentration of hydrogen ions in a solution. The value assigned is measured on a logarithmic scale which means that a drop in pH of 1 unit eg from 8 to 7, represents ten times the increase in hydrogen ions.

When carbon dioxide dissolves in seawater it forms weak carbonic acid that rapidly releases a hydrogen ion and a bicarbonate ion.

The hydrogen ions generated make the ocean more acidic (ie lower the pH) and some react with carbonate ions to produce more bicarbonate ion.


Thus, the net effect of increasing carbon dioxide in seawater is to increase concentrations of hydrogen ions, carbonic acid and bicarbonate while decreasing concentrations of carbonate ions making it more difficult for marine organisms to produce calcium carbonate shells.


In addition to direct impacts on marine calcifiers, other equally concerning impacts of increasing carbon dioxide in our oceans are emerging, including: the influence of ocean chemistry on trace metal availability to organisms, combined effects of carbon dioxide with temperature and oxygen on zooplankton metabolism, egg and sperm motility and fertilisation success, settlement and recruitment success for fish larvae, pheromone-like communication and behaviour cues for fish, and reduced sound absorption and thus a noisier ocean that may affect mammals and other animals that communicate acoustically.

The Issue	What we know	What we don't know	What we're doing	What's at stake	Online Links
<ul style="list-style-type: none">Ocean acidification  PHOTO 3	<ul style="list-style-type: none">The global ocean currently absorbs one-quarter of the anthropogenic CO₂ added to the atmosphere each year^{2,3}.Of the anthropogenic CO₂ emissions absorbed by the world's oceans, the Southern Ocean absorbs over 40%.	<ul style="list-style-type: none">Whether the Southern Ocean will continue to absorb the same proportion of anthropogenic CO₂ in future.Where and when new additions of CO₂ will combine with naturally varying distributions of CO₂ in the ocean to regionally amplify acidification (e.g. the Ross Sea may experience delayed impacts)³⁵.	<ul style="list-style-type: none">Quantifying Southern Ocean processes that influence the uptake of CO₂.Measuring and monitoring Southern Ocean carbon chemistry and ocean-atmosphere interaction.	<ul style="list-style-type: none">The continued climate mitigation service of the global ocean:<ul style="list-style-type: none">- Atmospheric CO₂ levels are tempered by the oceanic uptake of CO₂ i.e. without the ocean current atmospheric CO₂ concentration would be ~450 ppm³⁶ instead of ~387 ppm³⁷.- This hidden ocean 'service' has been estimated to represent an annual subsidy to the global economy of US\$60 – US\$400 billion per year³⁸.	<ul style="list-style-type: none">Global Carbon Project 2008: www.globalcarbonproject.org
 PHOTO 4	<ul style="list-style-type: none">The uptake of CO₂ into the ocean is driving a change in ocean chemistry: lowering ocean pH⁵ and the concentration of carbonate ions⁶ available to organisms that build calcium carbonate shells or skeletons (known as calcifiers).	<ul style="list-style-type: none">How Southern Ocean non-calcifiers will respond to changes in CO₂, pH and carbonate ion availability.How acidification will interact with other ocean variables likely to change in the future (e.g. temperature and oxygen).	<ul style="list-style-type: none">Studying past, present and likely future impacts of ocean acidification on Southern Ocean organisms including:<ul style="list-style-type: none">- pteropods and deep-sea corals (aragonite calcifiers) and- foraminifera and coccolithophores (calcite calcifiers).	<ul style="list-style-type: none">The role of Southern Ocean calcifiers in the global carbon cycle.	<ul style="list-style-type: none">Royal Society Report 2005: royalsociety.org/Ocean-acidification-due-to-increasing-atmospheric-carbon-dioxide/
 PHOTO 5	<ul style="list-style-type: none">Some calcifiers are more at risk than others: shells can be made from aragonite or calcite (and aragonite is more easily dissolved than calcite)⁷.	<ul style="list-style-type: none">How calcifiers made from aragonite and calcite will respond to changes in pH and carbonate ion availability.		<ul style="list-style-type: none">Fossil fuel CO₂ emissions are currently tracking above all scenarios used by the IPCC³⁹. If this trend in emissions continues, organisms would be subject to the lowest ocean pH experienced in the past ~20 million years by 2300 and the rate of fall in pH may possibly be the most rapid ever experienced⁴⁰.	<ul style="list-style-type: none">Ocean Acidification 2007: www.abc.net.au/catalyst/stories/s2029333.htm
 PHOTO 6	<ul style="list-style-type: none">Continued acidification will push the ocean towards chemical compositions not seen for a long time⁸:<ul style="list-style-type: none">- pre-industrial pH was 8.2⁹- current average pH is 8.1¹⁰- we expect a drop to 7.8 by 2100 under 'business as usual' CO₂ emissions¹¹.	<ul style="list-style-type: none">What areas of the Southern Ocean are most at risk of continued ocean acidification.	<ul style="list-style-type: none">Developing comprehensive Southern Ocean ecosystem models to illuminate:<ul style="list-style-type: none">- populations at risk,- communities at risk, and- regions at risk.	<ul style="list-style-type: none">The health and function of ocean ecosystems in vastly more acidic waters.	<ul style="list-style-type: none">CO₂ emissions and trends 2008: www.globalcarbonproject.org/carbonbudget/index.htm
 PHOTO 7	<ul style="list-style-type: none">Polar pH is changing at twice the rate of tropical waters¹².	<ul style="list-style-type: none">What the rate of acidification will be in Southern Ocean waters.		<ul style="list-style-type: none">Polar ecosystems are forecast to be among the first to be affected by acidification⁴¹.	<ul style="list-style-type: none">High Latitude Acidification 2009: tos.org/oceanography/issues/issue_archive/issue_pdfs/22_4/22-4_fabry.pdf

Impacts on Southern Ocean organisms



Ocean acidification is expected to impact on micro-organisms such as phytoplankton and zooplankton first. The impacts felt at this level feed up through the food web to higher trophic levels such as fish, birds and mammals. Here we summarise the current state of knowledge on the impact of ocean acidification on all trophic levels in the Southern Ocean in turn for both calcifying and non-calcifying organisms. We currently know most about the likely effects of ocean acidification on calcifying organisms as the bulk of our efforts have been directed to these 'sentinel' organisms. We note, however, that there is increasing concern that ocean acidification could affect the non-calcifying species. Non-calcifying phytoplankton (diatoms and flagellates) and non-calcifying zooplankton (copepods and krill) dominate the Southern Ocean ecosystem and the mechanisms of impacts of acidification on these and other higher trophic levels, such as fish, remains a serious gap in our knowledge.

Southern Ocean calcifiers	What we know	What we don't know	What we're doing	What's at stake	Online Links
<ul style="list-style-type: none"> Calcifying Phytoplankton - coccolithophores  <p>PHOTO 8</p>	<ul style="list-style-type: none"> Coccolithophores show both negative and positive responses to ocean acidification in laboratory experiments^{13,14}. The Southern Ocean coccolithophore <i>Emiliana huxleyi</i> appears to be shifting its habitat south due to environmental pressures¹⁵. 	<ul style="list-style-type: none"> How Southern Ocean coccolithophores respond to changing atmospheric levels of CO₂. Whether effects measured in laboratory studies differ due to experimental design (e.g. pH can be changed experimentally using CO₂ gas or by introducing acid). 	<ul style="list-style-type: none"> Studying the natural variability of Southern Ocean coccolithophores since the last glacial maximum. Following best practice guides for ocean acidification methods that have been recently established. 	<ul style="list-style-type: none"> <i>Emiliana huxleyi</i> is one of the most prominent producers of calcium carbonate in the world's ocean¹³. Increasing CO₂ can stimulate their growth through enhanced photosynthesis or hamper growth through enhancing dissolution of their calcium carbonate shells. As they play a significant role in regulating marine carbon cycling and atmosphere - ocean CO₂ exchange⁴² their likely response to future ocean acidification is of the utmost importance. 	<ul style="list-style-type: none"> Coccolithophores 2010: earthobservatory.nasa.gov/Features/Coccolithophores/ Guide to Best Practices 2010: epoca-project.eu/
<ul style="list-style-type: none"> Calcifying Zooplankton - foraminifera - pteropods 	<ul style="list-style-type: none"> Some species of modern planktonic foraminifera in the Southern Ocean have 30-35% lighter shell weights than their counterparts from pre-industrial times¹⁶. 	<ul style="list-style-type: none"> How calcifiers such as pteropods will respond to ocean acidification in the Southern Ocean (as they are made of a more soluble form of calcium carbonate than foraminifera). 	<ul style="list-style-type: none"> Investigating the impacts of ocean acidification on pteropod shells. Monitoring plankton and calcifier biodiversity in the Southern Ocean. 	<ul style="list-style-type: none"> Shelled zooplankton play important roles in the Southern Ocean food web as well as transporting atmospheric carbon to the deep ocean on their death⁴³. Models predict the Southern Ocean will become unstable for pteropod shell production as early as 2030⁴¹. 	<ul style="list-style-type: none"> Southern Ocean Sentinel 2010: www.abc.net.au/catalyst/stories/2886137.htm
<ul style="list-style-type: none"> Southern Ocean Corals 	<ul style="list-style-type: none"> Cold-water corals are likely to experience difficult water chemistry conditions by 2100¹⁰ (some as early as 2020¹⁷). Some Mediterranean corals survive in a different form, i.e. adapt, in undersaturated waters¹⁸. 	<ul style="list-style-type: none"> Whether Southern Ocean corals respond to acidification in the same way and at the same rate as tropical corals. Whether Southern Ocean corals are able to adapt to undersaturated waters. 	<ul style="list-style-type: none"> Investigating the distribution of Southern Ocean corals above and below the current aragonite saturation horizon in the Southern Ocean. 	<ul style="list-style-type: none"> Warm water corals are important fish habitats and have been demonstrated to be very susceptible to ocean acidification⁴⁴. Little is known about the potential impacts of acidification on coldwater coral-associated fish⁴⁵. 	<ul style="list-style-type: none"> Acid Test – can we save our oceans from CO₂? 2009: na.oceana.org/sites/default/files/reports/Acidification_Report1.pdf

Southern Ocean non-calcifiers	What we know	What we don't know	What we're doing	What's at stake	Online Links
<ul style="list-style-type: none"> Bacteria Phytoplankton <ul style="list-style-type: none"> diatoms 	<ul style="list-style-type: none"> A substantial fraction of the organic carbon in the oceans depends upon breakdown by heterotrophic bacteria¹⁹. Growth of some diatoms are enhanced with increasing CO₂ in lab experiments²⁰ although other researchers have found CO₂ enhanced productivity to be nutrient dependent²¹. 	<ul style="list-style-type: none"> Whether ocean acidification will impact bacterial processes and/or role in the marine ecosystem. Whether non-calcifying phytoplankton will have a competitive advantage in a high CO₂ ocean. 	<ul style="list-style-type: none"> Investigating impacts of ocean acidification on key members of the Southern Ocean community such as: <ul style="list-style-type: none"> bacteria, and phytoplankton through mini-cosm experiments. 	<ul style="list-style-type: none"> Changes in bacterial activity and growth efficiency can profoundly affect the oceanic net carbon balance. Diatoms contribute about half of all marine primary production and are known to actively take up CO₂ for photosynthesis⁴⁶. 	<ul style="list-style-type: none"> Ocean Acidification Effects on Cyanobacteria 2010: www.co2science.org/subject/o/summaries/acidificationcyano.php Ocean Acidification Effects on Diatoms 2010: www.co2science.org/subject/o/summaries/acidificationdiatoms.php
<ul style="list-style-type: none"> Zooplankton <ul style="list-style-type: none"> krill copepods  <p>PHOTO 9</p>	<ul style="list-style-type: none"> Antarctic krill <i>Euphausia superba</i> show impaired embryonic development at high CO₂ levels²². It also appears that survival and recruitment of early larval stages would be significantly reduced under elevated CO₂²³. In contrast to krill studies, high CO₂ exposure did not significantly affect survival, body size or developmental speed of the copepod <i>Acartia tsuensis</i>²⁴. 	<ul style="list-style-type: none"> Whether all life stages of krill respond negatively to high CO₂ concentrations. Whether copepods will have a competitive advantage in a high CO₂ ocean. 	<ul style="list-style-type: none"> Investigating the effects of acidification on the survival and development of all life stages of Antarctic krill. Determining Antarctic krill growth, recruitment and survival. 	<ul style="list-style-type: none"> Changing krill distribution is expected to impact predators including seals, penguins and whales³⁴. Planktonic copepods compose the bulk of the biomass in most pelagic zooplankton communities and are an important food source for higher trophic organisms including krill and fishes²⁴. 	<ul style="list-style-type: none"> The world's most abundant crustacean 2010: www.antarctica.gov.au/science/southern-ocean-ecosystems/krill
<ul style="list-style-type: none"> Fish 	<ul style="list-style-type: none"> Tropical juvenile clownfish olfactory (smell)senses are detrimentally affected as the acidity of seawater increases²⁵ and larval fish survival to adulthood is dramatically decreased by ocean acidification²⁶. 	<ul style="list-style-type: none"> Whether ocean acidification will impact upon Southern Ocean fish behaviour or population structure. 	<ul style="list-style-type: none"> Determining effects of ocean acidification on lower trophic levels likely to impact fish populations. 	<ul style="list-style-type: none"> Replenishment of fish stocks requires successful recruitment from larval to adult stage. Additional CO₂ absorbed into the ocean reduces recruitment success and has far-reaching consequences for the sustainability of fish populations. 	<ul style="list-style-type: none"> Ocean Acidification Effects on Fish 2010: www.co2science.org/subject/o/summaries/acidfish.php
<ul style="list-style-type: none"> Birds & Mammals <ul style="list-style-type: none"> penguins seabirds seals whales 	<ul style="list-style-type: none"> We know that seabird and penguin breeding success is diet dependent²⁷. A more acidic ocean will be a noisier place for whales^{28,29}. 	<ul style="list-style-type: none"> How impacts on krill, phytoplankton and zooplankton will be felt by higher trophic levels. 	<ul style="list-style-type: none"> Documenting foraging, population and reproductive success for: <ul style="list-style-type: none"> penguins seabirds seals and whales. 	<ul style="list-style-type: none"> Projected future CO₂ emissions^{28,29} imply significant increases in noise within the auditory range critical for environmental, military, and economic interests. 	<ul style="list-style-type: none"> A noisier ocean at lower pH 2010: www.nature.com/ngeo/journal/v3/n1/full/ngeo719.html

Impacts on Southern Ocean ecosystems

There is increasing concern that ocean acidification could affect the functioning of whole marine ecosystems, and particularly polar ecosystems. However, the mechanisms by which population impacts will occur have not been identified. The majority of our research to date has been laboratory-centric, making it difficult to extrapolate single-species impacts into whole of natural ecosystem settings. The best analogues we have for high-CO₂ ocean environments are natural marine ecosystems near volcanic vents. These ecosystems point the way toward dramatically different ecosystems once pH drops below 7.8.

Southern Ocean ecosystems	What we know	What we don't know	What we're doing	What's at stake	Online Links
<ul style="list-style-type: none"> Ecosystem Structure  <p>PHOTO 10</p>	<ul style="list-style-type: none"> Naturally high CO₂ marine environments elsewhere in the world exhibit major shifts in marine ecosystem structure as pH drops³⁰. In one study, the tipping point for structural change appears to be at a pH of 7.8³⁰. Southern Ocean ecosystems may be more or less sensitive. 	<ul style="list-style-type: none"> How changes in populations and communities of both calcifying and non-calcifying organisms will impact the greater Southern Ocean ecosystem structure. What impacts ocean acidification will have on higher trophic levels (e.g. fish). Whether ocean acidification will act with or against multiple stressors (e.g. temperature). 	<ul style="list-style-type: none"> Investigating impacts of ocean acidification on key members of the Southern Ocean community (such as phytoplankton) through mini-cosm experiments. 	<ul style="list-style-type: none"> Decreases in carbonate ion availability are likely to compromise the fitness of calcifiers and could shift the ecological and evolutionary competitive advantage to non-calcifying organisms⁴⁵. 	<ul style="list-style-type: none"> Ocean Acidification Network: www.ocean-acidification.net
<ul style="list-style-type: none"> Ecosystem Function  <p>PHOTO 11</p>	<ul style="list-style-type: none"> Calcifiers are important components of the Southern Ocean carbon cycle as they transport carbon to the deep ocean^{31,32} and contribute to the regulation of marine carbon cycling and the exchange of CO₂ between the ocean and atmosphere³³. The organisms at risk of acidification in the Southern Ocean are important components of the food web, as they are prey for higher trophic level species including krill, Antarctic fur seals, Adelie penguins and whales³⁴. 	<ul style="list-style-type: none"> Whether ocean acidification will affect processes and functions other than calcification (e.g. reproduction rates). How impacts on non-calcifying organisms will change the Southern Ocean ecosystem. How impacts on keystone species such as krill will then impact upon higher trophic levels (e.g. seals, whales and seabirds) and ecosystem services (e.g. fisheries). 	<ul style="list-style-type: none"> Addressing the Southern Ocean's role in global ocean circulation and ecosystem processes. Quantifying trophic links in Antarctic marine predators. Documenting foraging behaviour, population trends and reproductive success for penguins, seabirds, seals and whales. 	<ul style="list-style-type: none"> The full impact of ocean acidification on organisms is still in its infancy. We are not yet able to predict how impacts may propagate through marine ecosystems to ultimately effect higher trophic levels (e.g. fish) and the climate services the ocean provides us (i.e. continued absorption of a significant proportion of atmospheric CO₂)⁴⁵. We don't know how impacts will be felt in the natural setting on non-calcifying organisms and as these organisms hold important places in the Southern Ocean food chain (e.g. sea ice diatoms and krill) this of serious concern. 	<ul style="list-style-type: none"> Guide to Ocean Acidification for Policy Advisers and Decision Makers 2009: epoca-project.eu/index.php/what-do-we-do/outreach/rug/oa-the-facts.html Ocean Acidification Questions Answered 2010: epoca-project.eu/index.php/what-do-we-do/outreach/rug/oa-questions-answered.html

REFERENCES

- 1 Antarctic Climate & Ecosystems Cooperative Research Centre (2008) Position Analysis: CO₂ emissions and climate change: ocean impacts and adaptation issues. ISSN: 1835-7911
- 2 Sabine et al (2004) The oceanic sink for anthropogenic CO₂. Science 305: 367-371
- 3 Sabine C and Feely R (2007) The oceanic sink for carbon dioxide. In: Greenhouse Gas Sinks (ed) Reay et al, CABI Publishing pp. 31-49
- 4 Khatiwala et al (2009) Reconstruction of the history of anthropogenic CO₂ concentrations in the ocean. Nature 462: 346-349
- 5 Caldeira K and Wickett ME (2003) Anthropogenic carbon and ocean pH. Nature 425: 365
- 6 Doney et al (2009) Ocean Acidification: The Other CO₂ Problem. ARMS 1: 169-192
- 7 Mucci A (1983) The solubility of calcite and aragonite in seawater at various salinities, temperatures, and one atmosphere total pressure. AJS 283: 780-799
- 8 Pelejero et al (2010) Paleo-perspectives on ocean acidification. TEE 25: 332-344
- 9 Key et al (2004) A global ocean carbon climatology: Results from GLODAP. GBC 18: GB4031, doi:10.1029/2004GB002247
- 10 Orr et al (2005) Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. Nature 437: 681-686
- 11 Feely et al (2009) Ocean acidification: Present conditions and future changes in a high-CO₂ world. Oceanography 22: 36-47
- 12 Bindoff N and Willebrand J (2007) IPCC AR4 Chapter 5, Observations: Oceanic Climate Change and Sea Level, Cambridge University Press, Cambridge, 387-432
- 13 Riebesell et al (2000) Reduced calcification of marine plankton in response to increased atmospheric CO₂. Nature 407: 364-366
- 14 Iglesias-Rodriguez D (2008) Phytoplankton Calcification in a High-CO₂ World. Science 320: 336-340
- 15 Cubillos et al (2007) Calcification morphotypes of the coccolithophorid *Emiliania huxleyi* in the Southern Ocean: changes in 2001 to 2006 compared to historical data. MEPS 348: 47-54
- 16 Moy et al (2009) Reduced calcification in modern Southern Ocean planktonic foraminifera. Nature Geoscience 2: 276-280
- 17 Guinotte et al (2006) Will human induced changes in seawater chemistry alter the distribution of deep-sea scleractinian corals? Frontiers in Ecology and the Environment 4: 141-146
- 18 Fine and Tchernov (2009) Scleractinian Coral Species Survive and Recover from Decalcification. Science 315: 1811
- 19 U.S. Department of Energy Workshop Report (2008) Carbon Cycling and Biosequestration Integrating Biology and Climate Through Systems Science. March 4-6 and 17-18, 2008
- 20 Wu et al (2010) CO₂-induced seawater acidification affects physiological performance of the marine diatom *Phaeodactylum tricornutum*. Biogeosciences 7: 2915-2923
- 21 Yoshimura et al (2009) Impacts of elevated CO₂ on phytoplankton community composition and organic carbon dynamics in nutrient-depleted Okhotsk Sea surface waters BGD 6: 4143-4163
- 22 Kawaguchi et al (2010) Will krill fare well under Southern Ocean acidification? Biol. Lett. doi:10.1098/rsbl.2010.0777
- 23 Robinson (pers comm.) ACE CRC PhD Student Project: The effects of CO₂-induced ocean acidification on the survival and development of early larval stage Antarctic krill (*Euphausia superba*)
- 24 Kurihara H and Ishimatsu A (2008) Effects of high CO₂ seawater on the copepod (*Acartia tsuensis*) through all life stages and subsequent generations. MPB 56: 1086-1090
- 25 Munday et al (2009) Ocean acidification impairs olfactory discrimination and homing ability of a marine fish. PNAS 106: 1848-1852
- 26 Munday et al (2010) Replenishment of fish populations is threatened by ocean acidification. PNAS 107: 12930-12934
- 27 Lynnes et al (2004) Diet and reproductive success of Adelie and chinstrap penguins: linking response of predators to prey population dynamics. Polar Biol. 27 : 544-554
- 28 Hester et al (2008) Unanticipated consequences of ocean acidification: A noisier ocean at lower pH. GRL 35: L19601, DOI:10.1029/2008GL034913
- 29 Ilyina et al (2010) Future ocean increasingly transparent to low-frequency sound owing to carbon dioxide emissions Nature Geoscience doi: 10.1038/NGEO719
- 30 Hall-Spencer et al (2008) Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. Nature 454 : 96-99
- 31 Trull et al (2001) Moored sediment trap measurements of carbon export in the Sub-Antarctic and Polar Frontal zones of the Southern Ocean, south of Australia AJGR 106: 31489-3151015
- 32 Francois et al (2002) Factors controlling the flux of organic carbon to the bathypelagic zone of the ocean. GBC 16: doi:10.1029/2001GB001722
- 33 Holligan P and Robertson J (1996) Significance of ocean carbonate budgets for the global carbon cycle. GCB 2: 85-95
- 34 Reid and Croxall (2001) Environmental response of upper trophic-level predators reveals a system change in an Antarctic marine ecosystem. Proc. R. Soc. Lond. B 268: 377-384
- 35 McNeil et al (2010) A multi-decadal delay in the onset of corrosive 'acidified' waters in the Ross Sea of Antarctica due to strong air-sea CO₂ disequilibrium. GRL 37: doi:10.1029/2010GL044597
- 36 Doney et al (2009a). Ocean Acidification: The Other CO₂ Problem. ARMS 1: 169-192
- 37 Atmospheric CO₂ for September 2010 = 386.80ppm (Mauna Loa Observatory: NOAA-ESRL): co2now.org
- 38 Ocean Acidification Reference User Group (2009) This assumes a theoretical replacement cost based on sequestration of 2Gt C/yr at an expected future carbon credit price of \$30 – \$200/t CO₂
- 39 Raupach et al (2007) Global and regional drivers of accelerating CO₂ emissions. PNAS 104: 10288-10293
- 40 Turley C (2008) Impacts of changing ocean chemistry in a high-CO₂ world. Mineralogical Magazine 72: 359-362
- 41 McNeil B and Matear R (2008) Southern Ocean acidification: A tipping point at 450-ppm atmospheric CO₂. PNAS 105: 18860-18864
- 42 Nybakken (2001) Plankton and Plankton Community. In: Marine Biology: An Ecological Approach (ed) Cummings, San Francisco, pp. 38-93
- 43 Kleypas et al (2006) Impacts of ocean acidification on coral reefs and other marine calcifiers: a guide for future research. St. Petersburg Workshop Report 18-20 April 2005
- 44 Turley et al (2007) Corals in Deep-Water: Will the unseen hand of ocean acidification destroy cold-water ecosystems? Coral Reefs 26: 445-448
- 45 Raven et al (2005) Ocean acidification due to increasing atmospheric carbon dioxide. Policy document 12/05 Royal Society, London. The Clyvedon Press Ltd, Cardiff

CONTACTS

Donna Roberts

Ocean Acidification Project – ACE CRC
D.Roberts@acecrc.org.au

Tom Trull

Carbon Program Leader – ACE CRC
Tom.Trull@acecrc.org.au

Tessa Jakszewicz

Deputy CEO – ACE CRC
Tessa.Jakszewicz@acecrc.org.au

Tony Press

CEO – ACE CRC
Tony.Press@acecrc.org.au



ANTARCTIC CLIMATE
& ECOSYSTEMS CRC

ACE CRC Report Card

Southern Ocean Acidification

RC01-101123

ISBN: 978-1-921197-12-3

© Copyright Antarctic Climate & Ecosystems Cooperative Research Centre 2010

This work is copyright. It may be reproduced in whole or in part for study or training purposes subject to the inclusion of an acknowledgement of the source, but not for commercial sale or use. Reproduction for purposes other than those listed above requires the written permission of the Antarctic Climate & Ecosystems Cooperative Research Centre.

Requests and enquiries concerning reproduction rights should be addressed to:

The Manager
Communications
ACE CRC
Private Bag 80
Hobart
Tasmania 7001
Tel: +61 3 6226 7888
Fax: +61 3 6226 2440
Email: enquiries@acecrc.org.au
www.acecrc.org.au

Established and supported under the Australian Government's Cooperative Research Centres Program



The ACE CRC is a unique collaboration between its core partners the Australian Antarctic Division; CSIRO; University of Tasmania; the Australian Government's Department of Climate Change and Energy Efficiency; the Alfred-Wegener Institute for Polar and Marine Research (Germany); and the National Institute of Water and Atmospheric Research Limited (New Zealand) and a consortium of supporting partners.

PHOTOS

FRONT COVER PHOTO Steve Rintoul (CSIRO): The Southern Ocean.

PHOTO 2. Martin Riddle (Australian Antarctic Division): Cold-water coral reef, recently discovered 800m below the ocean surface on the edge of the Antarctic continental slope.

PHOTO 3. ACE CRC: Deploying sediment trap in the Southern Ocean aboard the Aurora Australis.

PHOTO 4. Russ Hopcroft (UAF, USA): Southern Ocean pteropod: *Limacina helicina antarctica*.

PHOTO 5. Gustaaf Hallegraeff (UTAS): *Emiliana huxleyi*: a single-celled planktonic alga known as a coccolithophore.

PHOTO 6. ACE CRC: Retrieving the sediment trap from the Southern Ocean aboard the Aurora Australis.

PHOTO 7. Wendy Pyper (Australian Antarctic Division): ACE CRC's 'Team Acid' sort calcifiers aboard the Aurora Australis in the Southern Ocean.

PHOTO 8. ACE CRC: Viewing foraminifera samples.

PHOTO 9. So Kawaguchi (Australian Antarctic Division): Antarctic krill: *Euphausia superba*.

PHOTO 10. Steve Nicol (Australian Antarctic Division): Antarctic fur seals.

PHOTO 11. Frederique Oliver (Australian Antarctic Division): Emperor penguins.