SOME ALGAE IN LAKES HUME AND MULWALA (1974)

VICTORIA

By

KANJANA VIYAKORNVILAS B.Sc. (Tas.)

A Thesis submitted in partial fulfilment of the requirements for the Degree of Bachelor of Science with Honours at the University of Tasmania. I hereby declare that this thesis contains no material which has been accepted for the award of any other degree in any university and that, to the best of my knowledge, the thesis contains no copy or paraphase of material previously published or written by another person, except when due reference is made in the text.

ACKNOWLEDGEMENTS

I wish to express my sincere thanks to Dr. P.A.Tyler, my supervisor, for his helpful advice and criticism. Mr. R.L.Croome for collecting all the samples. Mrs. R.Wickham for her technical advice. All members of the Botany Dept. for their assistance at various times of the year.

> K.Viyakornvilas, Botany Dept., University of Tasmania, November 1974.

CONTENTS

*		Page
Summary		1
Introduction		2
Materials and Me	ethod	2
Results		4
Systemat	ic account	13
Division	Chlorophyta-Class Chlorophyceae	13
17	Chrysophyta-Class Chrysophyceae	65
	Class Bacteriophyceae	68
11	Euglenophyta-Class Euglenophyceae	79
₹\$	Pyrrhophyta-Class Dinophyceae	86
51	Cyanophyta -Class Cyanophyceae	87
Discussion		
The trop	nic status of Lakes Hume and Mulwala	93
Geograph	ical distribution of desmids seen	95
Plates 1-14 and	Explanation of plates	96
Literature cite	1	116

SUMMARY

Lakes Hume and Mulwala have very similar plankton communities in which <u>Melosira granulata</u> Ralfs is dominant. The species composition and the plankton quotients show the two lakes are mesotrophic.Most of the algae seen are well-known and widespread. <u>Micrasterias Hardyi</u> G.S.West has only been · recorded from Australia.

INTRODUCTION

Lakes Hume and Mulwala, parts of the Murray River (fig.1) Victoria, are near the boarder of Victoria and N.S.W. The aims of this study are three-folded:

1) to identify most of the phytoplankton species in both lakes,

2) to determine the trophic status of both lakes from the species composition and the phytoplankton quotients,

3) to survey the geographical distribution of the desmids seen.

The algae will be identify solely from their morphology. Attempts will be made to put them into existing taxa. In recent years advances have been made on the identification of algae by the use of cultures and scanning electron microscope. The latter is particularly useful in the case of <u>Cosmarium sp.</u> or the diatoms, etc. where the wall structures are the taxonomic criterion (Lyon 1969). Culture experiments are useful when one wishes to delimit species. MATERIALS AND METHOD

20 (1974) Hume and 7 (1974) Mulwala fixed samples as well as some live samples, all were collected by Mr.R.L.Croome with a 20 μ pore net at various stations shown in fig. 1, were examined on a Carl Zeiss RA microscope fitted with a prism drawing system with internally reflected image, allowing binocular viewing while drawing. 1 (1968) Hume and 6 (1968-9) Mulwala fixed samples were also examined.

The 1974 samples were denoted as date/month of collection. The numbers in brackets following the Hume samples indicated the station numbers. All the 1968, 1969 samples were denoted by year/month/date/number of the sample.



G.

RESULTS

TABLE 1: THE ALGAL COMPOSITION OF LAKES HUME AND MULWALA (1968,

1	969	AND	1974)
	+		seen in 1974 samples
	* *		seen in 1974, 1968, 1969 samples
	210	9.6 612	not seen in 1974 samples
	aje .	dince Visual	seen only in 1968/1969 samples

Species		Hume	Mulwala
Division Chlorophyta	£0~44.65572668746874687468748878878878878878678787878	an andremeiningkoonderspilanseoon vaar	yndynastanning sonnymine y synanning a synannymine y synannymine y synannymi
Class Chlorophyceae			
Order Volvocales			
Eudorina elegans Ehrenb.		+	+
Pandorina morum (Muller) Bory		+	++
Pleodorina californica Shaw		-	. +
Sphaerocystis Schroeteri Chodat		++	++
Volvox cf. aureus Ehrenb.		+	*
" " Carteri Stein		*	*
" globator Linnaeus		650	*
Order Ulotrichales			
Ulothrix subconstricta G.S.West		esus;	++
Order Oedogoniales			
Oedogonium sp.		++	ting
Order Chlorococcales			
Actinastrum Hantzschii Lag.		ф.	+
Ankistrodesmus falcatus Corda		-	+
Botryococcus Braunii Kuetz.		**	4 +

Genus and species names are not underlined for simplicity of table.

Species	lume	Mulwala
Coelastrum microporum Naeg.	anticuleturrest websonali met terrest	1999 - 1999 -
" reticulatum (Dang.) Senn.	4.	*+
Dictyospaerium pulchellum Wood	-4-	+
Dimorphococcus lunatus A.Braun.	+	
Kirchneriella lunaris(Kirchner) Moeb.	+	- afa
Micractinium pusillum Fres.	+, *	.
Nephrocytium Agardhianum Naegeli	+	+
Oocystis lacustris Chodat	+	
" parva West & West	+	an
Pediastrum duplex Meyen forma	+	++
" var. clathratum (A.Brun.) Lag.	+	ağı.
" var. gracillimum West & West	+	4
" var. rugulosum Raciborski	+	*
" tetras (Ehrenb.) Ralfs	+	+
Scenedesmus denticulatus Lagerh. var.		
denticulatus May	-	+
" obliquus (Turp.) Kuetz. var.		
dimorphus May		+
" opoliensis Richter	+	- -
" quadricauda (Turp.) Breb.	+	+
" " var. quadrispina (Chodat)	
G.M.Smith	60%)	+
Selenastrum westii G.M.Smith	+	+
Tetraedron gracile (Reinsch) Hansgirg	+	+
"hastatum(")"	+	4327
" limneticum Borge var. gracile Prescot	t +	

	Species	Hume	Mulwala
Ϸ άδυξητη τη του	៹៹៳៹៹៹៹៹៶៷៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹៹	14 m2.400824 %Ac.100.40.407864.19664	nicht chiv 1 Ebwardiskiniker värbarnärkingslan i värschaberd
Tetraedror	n regulare Kuetz.	-	+
11	victorieae Wolosz var. major Smith	4-	ésar
Treubaria	triappendiculatum Bernard	4	+
Order Zygr	nematales		
Family Zyg	nemataceae		
Cf. Mougeo	otia sp.	+	Sat.
Spirogyra	sp 1	50m	+
1,1	sp 2	-tr	*
Family Des	smidiaceae		
Arthrodesm	us constrictus G.M.Smith		
	var. longispinus (Gronbl.)	- † -	*
68	convergens Ehrenb.	****	*
Closterium	aciculare T.West var. subpronum Wests		+
11	ceratium Perty	-	+
11	gracile Breb.	esta:	+
Ŧŧ	Kutzingii Breb.	ವ್ಯೆ	-1-
78	moniliferum (Bory) Ehrenb.	4	4 ~
f 9	striolatum Corda	÷	4500
18	Venus Kutz. var. inflatum Claassen 19	61+	- -
Cosmarium	bipunctatum Borgesen	+	Bite
\$ F	contractum Kirchn.	+	6148
**	granatum Breb. var. pyramidal Schmidle	+	
ŦŦ	magnificum Nordst.		*+
78	cf. pseudonitidum Nordst.	632	+
ŦŦ	punctulatum Breb. var. subpunctulatum		
	Borge		903 1

" quadrum Lund

Spec	cies	Hume	Mulwala
Euastrum anasatum (Ehrei	nb.) Ralfs	54787847974848976497878689844978786898449787868978978 64669	
" cuspidatum var	• goyazense Forster	+	+
" divergens Josh	ua	-	+
" elegans Kutz.		+	1000
Gonatozygon kinahani (A:	rcher.) Rabenh.	++	+
" monotaenium	De Bory	• •	
Micrasterias Hardyi G.S	.West	++	+
" decemdenta	ta (Nag.) Archer.	6 49	+
" mahabuleswa	arensis Hobson	çan.	++
" thomassian	a var. notata (Nordst.)	×.	
	Gronbl.	4-	1
Pleurotaenium Ehrenberg:	ii (Breb.) De Bory		
var. undu	latum Schaarschmidt	+	
Spondylosium planum (Wo	lle) West & West	+	*
Staurastrum anatoides S	cott & Prescott	-	*
" arctiscon ()	Ehrenb,) Lund var, glabru	m	
Will Construct of the Williams	est & West		aler aler
" asterias Nv	gaard		+
" dickiej Ral	fs	+	++
" grande Bulbu	n, var, parvus West	•	2
" Freemanii W	est & West	-b-	
" furcatum (E)	hrenh,) Breh		*
" lentocladum	Nordst	+	*
	rescott		т.
II nachanhanahi	um Nordst		T
Il natone Turn.	an	Ŧ	*

€	Species	Hume	Mulwala
	๚๛๛๚๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛	esalja realizativljena utjekovaljenovalje	1998 M M Raha C 1995 M S 2004 RD 1220 PC 10 PC 1
Staurastrum	pingue Teiling	++	++
11	" " (small form)	+ .	+
**	cf. pinnatum Turner var. subpinnatum		
	Schmidle	Day.	+
11	Playfairi Scott & Prescott	+	++
11	pseudosebaldi Wille var. planctonicu	n	
	Teiling		++
11	sebaldi Reinsch. var. ornatum Nordst	0 ***	*
"	cf. subavicula West	+	-
11	subgemmulatum West & West	+	+
"	tetracerum Ralfs	+	*+
n ,	" var. evolutum West & West	+	-
11	tohopekaligense Wolle	+	*
11	victoriense West		*
Staurodesmu	s cuspidatus (Breb.) Ralfs var.		
	tricuspidatus (Breb.)	+	500 1
11	triangularis (Lagerh.) Teiling	+	+
Division Ch	rysophyta		
Class Chrys	ophyceae		
Dinobryon di	ivergens Imhof.	+	++
יי ה	ertularia Ehrenb.	+	
Mallomonas	akrokomos Ruttner	•••	+
19	splendens (G.S.West) Playfair	+	
11	sp.	` + -	4.
Synura spin	osa Korsh	++	++

Species	Hume	Mulwala
Class Bacillariophyceae	##34*##@v=#0;#0#0###############################	<mark>Genergyweddan Marten O</mark> mmellan ei gynweddwer Anwr
Centrales		
Attheya Zachariasi J.Brun.	+	++
Cyclostella stelligera Cleve & Grun.	+	803
Melosira granulata Ralfs	++	++
" varians Agardh.	++	++
Rhizosolinia erensis H.L.Sm.	-+-	+
" " var. morsa West & West	• •	+
Pennales		
Asterionella formosa Hass.	+	+
Cymbella cf. gastroides Kutz.	+	+
" lanceolata Ehrenb.	+	-
Epithemia sorex Kutz.		++
Fragilaria capucina Desm.	+	+
"sp.	+	
Gomphonema Augur Ehr. var. Guatieri H.V.H.	800	+
" constrictum Ehrenb.	+	++
" var. capitatum Ehrenb.	· .	+
Cf. Gyrosigma attenuatum	- t -	++
Cf. Navicula hennedyi	afr	-
Nitzschia pelagica O.Mull.	+	+
Pinnularia abaujinsis (Pant.) Ross var.		
abaujinsis	400	+
Stauroneis fulmen var. capitata	6004 ·	x¢:
Surirella Engleri Q.Mull.	+	÷
" spinifera Hust.	+	*+

	Species	Hume	Mulwala
Surirella tenera G	reg.	. +	
Synedra acus Kutz.		+	++
n n var.	radians (Kg.) Hust.	+	÷+
" ulna (Nitz	.) Ehr. var. danica (Kg.) V.H. +	-
Tabellaria floccul	.osa (Roth) Kutz. var. fl	occulosa	
	Knudson	+	*
Division Euglenoph	yta		
Class Euglenophyce	ae		
Order Euglenales			
Euglena cf. gracil	is Kleb.	ára .	+
" spirogyra	Ehrenb.	+	+
Phacus tortus (Lem	.) Skvortzow	+	
" pleuronecte	s (Muell.) Dujardin	• •	4 433
Trachelomonas arma	tus var. longispina (Pla	vfair)	
	Dêfl	andre +	
u u baci	llifera Plavf.	+	•
	ata Stein var. australic	а	•
	Plaufair	~	
ll ll biss	itayiaii	ronata	
ursł	Iua (Ferty) Stern var to	I Una ca	
· •• •• ••	neum*		600
	" var. rectangularis		
	Br. Schroder	+	6334
" " scab	ora Playfair	6554	· +
" " supe	rba Swirenko	+	+
" " volv	ocina Ehrenb. var. punct	ata	
	Playfair.	+	*

Species	Hume	Mulwala
Division Pyrrhophyta	ለዚያው የሪድርስ እና 25 28 28 20 20 20 20 20 20 20 20 20 20 20 20 20	Bearlow Obertald Quarter Hornby Auropention
Class Dinophyceae		
Order Peridiniales		
Ceratium hirundinella (O.Muell) Dujardin	+	+
Glenodinium sp.	+	6×4
Peridinium sp.	+	 .
Division Cyanophyta		
Class Cyanophyceae		
Order Chroococcales		
Coelospharium Naegelianum Unger.	++	+
Microcystis aeruginosa Kutz.	+	+
" flos-aquae (Wittr.) Kirchner.	ديچ	*
Order Nostocales		
Anabaena limnetica G.M.Smith	+	6 4
" spiroides Kleb.	+	+
Lyngbya palmarum Bruhl et Biswas		+
Oscillatoria curviceps Agardh ex Gomont	65	+
" lacustris (Kleb.) Geitler	-	+
" nigra Vaucher	6725	+
" cf. subbrevis Schmidle		*
waterbergensis Claassen 1961		*
Schizothrix rivulis (Wolle) Drouet	+	

TABLE	2:	THE	PHYTOPLANKTONOUOTIENTS	OF	LAKES	HUME	AND	MULWALA
					and the second sec	and the second state of the second data and the second state of th	to be a second as a second	the sector of th

	an ann a chaoladhachadh an allan ain chaolan a san an deanna ann an bhailtean dhailte fha rada			
	No of species			
	Hume	Mulwala		
Myxophyceae	5	8		
Chlorophyceae	29	24		
Centrales	6	5		
Euglenideae	10	6		
Total	50	43		
Dësmidiaceae	34	30		
Phytoplankton quotients	$\frac{50}{34} = 1.47$	$\frac{43}{30} = 1.43$		

TAXONOMIC COMMENTS.

All dimensions are given in microns.

Abbreviations.

W = width	c p r = with processes
L = length	spr = without processes
Ist. = isthmus width	csp = with spines
dia. = diameter	ssp = without spines

Division Chlorophyta

Class Chlorophyceae

Volvocales

Éudorina elegans Ehrenberg. Pl 2 fig. 8

Smith 1920 Pl 19 fig. 1 p. 96

Prescott 1962 Pl 1 figs. 24-26

cell dia 8-10

colony dia 77-86

Eudorina is distinguished from <u>Pandorina</u> by its colony of distanced cells with one to several pyrenoids. (Prescott 1962). Playfair (1923) believed <u>P.morum</u> become regularly spherical and form <u>E.elegans</u>. Dr. P.Tyler (pers. comm.) thinks it probably does,too. When <u>P.morum</u> divides the mucilage expands and the colony then resembles (or equals) <u>E.elegans</u>.

E.elegans is widespread and has been recorded from Australia. (Hardy 1906, West 1909, Playfair 1916,1923, Thomasson 1973). Bailey 1895 <u>P.morum</u> (pl. VIII fig. 8) is probably <u>E.elegans</u>. Samples: Lake Hume 9/7(3), 11/9(5)

Lake Mulwala 10/2, 4/4, 680921/1.

Pandorina morum Bory Pl 2 figs 10,11 Smith 1920 Pl 19 figs 16,17 p. 95 Prescott Pl 1 fig 23 p. 75 (1962)

cell dia 8

colony dia 24-50

Playfair(1915)stated that the Australia plant differed from the European plant by having coenobia ovate. This is not correct because Smith (1920) and Prescott (1962) figured ovate P.morum from Wisconsin.

This alga is widespread. Previous Australian records-West (1909), Playfair (1915,1923).

Samples: Lake Hume 1/5(3), 30/5(5), 9/7(3), 10/7(3), 8/8(4),9/9(3), 11/9(5).

Lake Mulwala 4/4, 12/8, 681121/2, 680921/1.

Pleodorina californica Shaw Pl 2 fig. 9

Prescott 1962 Pl 2 fig. 1 p. 77

The colony (Pl 2 fig. 9) was viewed from the anterior pole where larger reproductive cells resided. (Prescott 1962).

Vegetative cells dia 4-5

Reproductive cells dia 8-9

Previous Australian record: none

Samples: Lake Mulwala 10/2.

Sphaerocystis Schroeteri Chodat Pl 4 figs. 10-12

Smith 1920 Pl 19 figs. 3-4

Prescott 1962 Pl 3 fig. 6 p. 83

Clusters of recently divided cells were somtimes within envelopes inside the colonial envolope.

cells dia 9-14

colony dia 40-126

Sphaerocystis Schroeteri and Planktosphaeria gelatinosa are so similar the only difference between them is the presence of several parietal, polygonal chloroplasts each with one pyrenoids in the latter. (Smith 1920). Pl 4 fig. 10 has one cell with three pyrenoids. It could be <u>Planktosphaeria gelatinosa</u>. Although it is possible the two species are forms of the same plant, West (1909) did not observed any life stages of <u>Sphaerocystis Schroeteri</u>(fig. 8, p. 76) that resemble <u>Planktosphaeria gelatinosa</u>.

Previous Australian record: West (1909).

Samples: Lake Hume 14/2(Dam), 7/3(3), 8/3(Tallangatta), 2/4(Tallan.)

5/4(Dam), 1/5(3), 2/5(4), 3/5(Dam), 30/5(3), 3/6(5), 8/7(5), 9/7(3), 10/7(3), 7/8(5), 9/9(3), 10/9(4), 11/9(5), 681122/3.

Lake Mulwala 10/2, 4/4 7/5, 4/6, 680921/1, 681121/2,

690206/1.

Volvox sp.

Three morphological forms of <u>Volvox sp.</u> were observed. (Pl 2 figs. 12-16). They could not be identified by Smith 1944 key because of the lack of all stages of asexual and sexual reproduction.

Volvox cf. aureus Ehrenberg. Pl 2 figs. 12-13

Smith 1920 Pl 18 fig. 20 p. 98

Smith 1944 fig. 3

.

The specimens may be <u>Volvox aureus</u> Ehrenberg. because of the presence of cytoplasmic connections. <u>Volvox aureus</u> is not the only species which possesses cytoplasmic connections. (Smith 1944).

cell dia 5-6

Samples: Lake Hume 14/2(Dam), 7/3(3), 8/3(Tallan.), 31/5.

Volvox cf. Carteri G.M. Smith Pl 2 figs. 15,16 Smith 1920 Pl 18 fig 1 p. 99 Smith 1944 fig. 37 as Volvox Carteri Stein.

The specimens have no cytoplasmic connections. Smith (1944) depicted <u>Volvox Carteri</u> Stein with undivided gonidia of equal size. The plant seen (Pl 2 fig. 15) may have some gonidia divided therefore some gonidia are bigger than others.

cell dia 4-5

colony dia 780

Samples: Lake Hume 681122/3

Lake Mulwala 680921/1, 681121/2.

Volvox cf. globator Linnaeus Pl 2 fig. 14

The pentagonal to hexagonal cells suggest <u>Volvox globator</u> L. (Smith 1920 Pl 18 fig. 3 p. 98, Smith 1944 fig. 7)

cell dia 5-6

Volvox globator Playfair (Playfair 1915, Pl 44 figs. 5-6,

p. 341) and <u>Volvox globator var. australis</u> Playfair 1915, p. 342 are <u>Volvox Merrilli</u> Shaw according to Smith 1944.

Sample: Lake Mulwala 680921/1.

<u>Ulotricales</u>

Ulothrix subconstricta G.S.West Pl 5 fig. 5

The closest the Mulwala plants resemble is <u>Ulothrix</u> <u>subconstricta</u> G.S.West (Prescott 1962 Pl 6 fig. 11, p. 96, Ramanathan 1964 Pl 10 fig. 10G, p. 39). The plants differ from the above authers'in having chloroplasts extend throughout the entire length of cells but the length of chloroplasts compared with the cells'length is variable in <u>Ulothrix sp.</u> (Lokhorst & Uroman 1972).

The cell dimensions are comparable with those given by Ramanathan 1964. cell L 9, dia 6. There is no previous Australian record. Smith (1920) was of the opinion that <u>Ulothrix subconstricta</u> is the only one of the order which has become established as a true member of the plankton. Samples: Lake Mulwala 4/4, 680921/1.

Oedogoniales

Oedogonium sp. Pl 5 fig. 1

The Hume plants are <u>Oedogonium sp</u>.because cells possess scar rings. Also, they are unbranched filaments with cells showing a slight inflation toward one end. The chloroplasts are parietal, each containing several pyrenoids (Prescott 1962).

cell L 54-63

W 13-17

In differentiating the species of <u>Oedogonium</u>, size of vegetative cells, the shape and size of the oogonia, the form and decoration of the zygospore wall and the location of the antheridal cells are the most important differentiating and specific characters. (Prescott 1962). In the absence of these characters one cannot possibly do more than <u>Oedogonium</u> sp.

Samples: Lake Hume 3/5(Dam), 681122/3.

Chlorococcales.

Actinastrum Hantzschii Lag. Pl 3 figs. 1-5

There are two morphological forms of <u>Actinastrum</u> one of which is wider (Pl 3 figs. 1-3) than the other (Pl 3 figs. 4-5). Smith (1920) designated both forms as <u>Actinastrum Hantzschii</u> and Actinastrum gracillimum G.M.Smith respectively.

Playfair (1916), Haughey (1968 figs. 9,10) grouped <u>Actinastrum</u> with outer ends of cells acute and inner ends blunt of the same dimensions as both forms of the plants found in <u>Actinastrum</u> <u>Hantzschii</u>. Their grouping will be followed. Cell L 7-38, W 2-5.. Smith (1920) depicted a specimen (Pl 43 fig. 8) which resembled his <u>Actinastrum gracillimum</u> (Pl 43 fig. 4) and called it <u>Actinastrum</u> <u>Hantzschii var. elongatum</u> because of its larger size. It is possible that Smith s'<u>Actinastrum Hantzschii var. elongatum</u> are bigger cells of <u>Actinastrum gracillimum</u> and <u>Actinastrum gracillimum</u> and <u>Actinastrum Hantzschii</u> are forms of the same plant.

The multiple colony (Pl 3 fig. 5) is held together by lateral contact (Philipose 1967).

Previous Australian and New Zealand records: Playfair 1916, Scott & Prescott 1958, Haughey 1968.

Samples: Lake Hume 7/3(3), 1/5(3), 2/5(4), 3/5(Dam), 31/5(4),

3/6(5), 8/7(5), 9/8(5), 10/9(3), 11/9(5).

Lake Mulwala 4/4, 7/5, 4/6, 12/8.

Ankistrodesmus falcatus Corda Pl 3 fig. 6

Smith 1920 Pl 32 fig. 1

Prescott 1962 Pl 56 fig. 5

Philipose 1967 figs 121 a,c

Cell L 50

W 3

Ankistrodesmus falcatus Corda is widespread and has been recorded from Australia and New Zealand (Hardy 1906, West 1909, Playfair 1916, Scott & Prescott 1958, Thomasson 1973, and Maskell 1881, Haughey 1968).

Edward (1894) observed cells of <u>Ankistrodesmus falcatus</u> loosened in culture then curved like those of <u>Ankistrodesmus</u> <u>convolutus</u> (Philipose 1967 fig. 122 a-d) which are smaller. Haughey (1968) observed a gradation from curved, larger to straight, smaller cells in his samples.

Sample: Lake Mulwala 4/4.

Botryococcus Braunii Kuetz Pl 2 fig. 17

This alga is common and often abundant, especially in semi hard water lakes where it frequently is the dominant component of water bloom associations(Prescott 1962). It has been recorded from Australia. (Bailey 1893, Hardy 1906, West 1909, Scott & Prescott 1958, Thomasson 1973).

Samples: Lake Hume 2/5(4), 3/5(Dam), 681122/3.

Lake Mulwala 7/5, 681121/2. <u>Coelastrum microporum</u> Naeg. Pl 3 fig. 16 Haughey (1969) fig. 4:1,1a

cell dia including shealth 14 Previous Australian and New Zealand records: West (1909),Scott & Prescott 1958, Thomasson (1973), Haughey 1969. Samples: Lake Hume 3/5(Dam). <u>Coelastrum reticulatum</u> (Dang.) Senn Pl 3 fig. 15 Smith (1920) Pl 42 fig. 4 p. 161 Bourrelly (1966) Pl 34 fig. 6 Prescott (1962) Pl 53 fig. 6 p. 230

The plants have cell dia with sheath of 5-6 which is in the range recorded by Smith (1920). Philipose (1967) figured <u>Coelastrum reticulatum</u> (Dang.) Senn. (fig. 142) which looked neither like the specimens nor the illustrations given by the above authôrs.

Previous Australian record: Playfair (1918). Samples: Lake Hume 5/4(Dam), 3/5(Dam), 3/6(5).

Lake Mulwala 10/2, 690206/1. <u>Dictyosphaerium pulchellum</u> Wood Pl 3 figs. 10,11 Smith 1920 Pl 20 fig. 13 p. 105 Prescott 1962 Pl 51 fig. 6 Dictyosphaerium pulchellum Wood observed (P13 figs. 10,11) do not have cells as round as those depicted by the mentioned workers but, according to Smith (1920) the cells are sphaerical when mature, sphaerical to ovoid when young. The Hume & Mulwala cells are not as ellipsoid as those of <u>Dictyosphaerium ehrenbergianum</u> Nag.(Haughey 1968 fig. 72). <u>Dictyosphaerium pulchellum</u> was given as a synonym of Dictyosphaerium ehrenbergianum (Bailey 1898).

> Sometimes the colony envelope is not seen (Pl 3 fig. 11). cell L 5-7

cell dia. 4-5

Dictyosphaerium pulchellum is widespread (Philipose 1967). Previous Australian records: Schmidle 1896, Hardy 1906, West 1909, Croome & Tyler (1972), Thomasson (1973).

Samples: Lake Hume 8/3(Tallan.), 2/5(4), 3/5(Dam), 3/6(5), 8/7(5),

11/9(5).

Lake Mulwala 4/6.

Dimorphococcus lunatus A. Braun. Pl 3 fig. 14

Philipose 1967 fig. 115a p. 205

cell L 7

₩ 7-12

Previous Australian records: Scott & Prescott 1958, Thomasson 1973. Samples: Lake Hume 3/5(Dam), 3/6(5).

Kirchneriella lunaris (Kirchner.) Moeb. Pl 3 fig. 14

Smith 1920 Pl 34 fig. 4

Philipose 1967 fig. 131

The following dimensions are within the range given by the above authors.

.....

.

cell L 10-13

W 6-9

<u>Kirchneriella lunaris</u> is widespread (Smith 1920) and has been recorded from Australia (Bailey 1895, Schmidle 1896, Hardy 1906, West 1909, Playfair 1914, Thomasson 1973).

Bailey (1895) noted "-----neither a gelatinous envelope nor a regular arrangement of groups into four appears remarkable-----Âccording to Philipose (1967) the gelatinous envelope is visible only after stainking. No gelatinous envelope was observed but cells were in groups of four. P13 fig. 14.

Samples: Lake Hume 5/4(Dam), 1/5(3), 2/5(4), 3/5(Dam), 3/6(5).

Lake Mulwala 10/2, 7/5.

Micractinium pusillum Fres Pl 2 fig. 18

Smith (1920) Pl 28 fig. 1-3

Haughey (1968) fig. 9:3a

cells dia. 4-5

setae L 10-14

Haughey (1968) recorded this form with rather short setae from NZ.

Previous Australian record: none

Samples: Lake Hume 2/5(4), 3/5(Dam), 3/6(5).

Nephrocytium Agardhianum Naegeli Pl 3 figs. 24-26

Smith (1920) differentiated genera <u>Oocystis</u> and <u>Nephrocytium</u> by their cell shape.

Cells ovoid to elleptic, never curved-----Oocystis

Cells ovoid to reniform, frequently curved-----<u>Nephrocytium</u> Using the above key the plants (Pl 3 figs. 24-26) are <u>Nephrocytium</u>. They are <u>Nephrocytium Agardhianum</u> Naeg. (Philipose 1967, fig. 104b, p. 189). Cells L 12-21, W 5-11.

Previous Australian records: West 1909, Borge 1911, Playfair 1912, 1916, (varieties).

Samples: Lake Hume 2/5(4), 3/5(Dam), 31/5(4), 3/6(5), 10/7(3), 11/9(5). Lake Mulwala 4/6, 7/5. Occystis lacustris Chodat And Occystis parva West & West.

There are two morphological types of <u>Oocystis</u> in Lake Hume. One type has pointed ends which suggest polar thickenings (Pl 3 figs. 21-23). Their cells'shape and dimensions agree with those of <u>Oocystis lacustris</u> Chodat (Smith 1920 Pl 22 figs. 8-9 p. 112, Rehakova 1969 Tab. 1 fig. 6).

cell L 13-18

W 9-10

Another type (Pl 3 figs. 19-20) closely resembles <u>Oocystis parva</u> West & West (Smith 1920 Pl 22 fig. 6 p. 112). Although Smith (1920) stated this species had pointed ends, some of his cells (Pl 22 fig. 6) had rounded ends like the Hume plants. Rehakova 1969 figured <u>Oocystis parva</u> West & West of both round and pointed ends. The plants'dimensions agree with those of Oocystis parva given by Smith (1920).

cell L 14-15

W 7

West (1909) recorded both forms from Vic.

Playfair (1916) regarded both <u>Oocystis lacustris</u> & <u>Oocystis parva</u> as polymorphic forms of one organism. This is probably very correct. It is known what initiates polar thickenings. One may culture <u>Oocystis parva</u> and find out if they form polarthickenings in which case <u>Oocystis parva</u> and <u>Oocystis lacustris</u> are forms of one biological species.

Samples: <u>Oocystis lacustris</u>

Lake Hume 2/5, 31/5, 9/7.

Oocystis parva

Lake Hume 14/2(Dam), 5/4(Dam), 1/5(3), 2/5(4), 3/5(Dam),

30/5(3), 9/7(3), 9/9(3).

Pediastrum duplex Meyen

The type form (Pl 4 fig. 4) and three varieties of <u>Pediastrum duplex</u> Meyen were found. Pl 4 fig. 4 is the type form (Smith 1920 Pl 46 fig 16). Cell dia. 10-22.

Samples: Lake Hume 8/3(Tallan.), 1/5(3), 2/5(4), 3/5(Dam), 30/5(3),

3/6(5), 9/7(3).

Lake Mulwala 10/2, 4/4, 4/6,

680508/1, 680411/1, 690206/1.

<u>Pediastrum duplex var. clathratum</u> (Braun. A) Lagerheim Pl 4 figs. 1-2 (Smith 1920 Pl 47 fig. 3). Sulex (1969) in the recent Europeanmonograph recorded this variety as <u>var. duplex</u> and stated it is synonymous with <u>P.duplex var. clathratum</u>. Cell dia. 7-11. Samples: Lake Hume 5/7(2).

Lake Mulwala 681121/2.

Pediastrum duplux var. gracillimum W.& G.S West (Pl 4 fig. 3) (Smith 1920 Pl 47 figs. 8-11 p. 172, Haughey 1968 fig.6:3)

cell dia 8

Huber-Pestalozzi (1936) depicted <u>P.duplex var. gracillimum</u> with two cells in the middle as seen in one colony (Pl 4 fig. 6). Usually this <u>var. gracillimum</u> has no cells in the mid region as depicted in Pl 4 fig. 3.

Samples: Lake Hume 1/5(3), 3/5(Dam).

Lake Mulwala 12/8.

<u>Pediastrum duplex var. rugulosum</u> Raciborski Pl 4 fig. 7 Sulek 1969 Beil 4 fig. 4.

cell dia. 11-17

Bailey (1895) recorded this variaty as <u>P.duplex</u> from Queenland (Pl VII fig. 9).

Samples: Lake Hume 8/3(Tallan.)

Lake Mulwala 681121/2, 690206/1.

The arrangement of big cells with two cells in the midregion(Pl 4 fig. 5) is not recorded in the recent European monograph by Sulex 1969. Cell dia 15. Sample: Lake Hume 3/5(Dam).

Smith (1920) stated that in many cases the marginal cells of <u>P.duplex</u> and the variety <u>clathratum</u> have a tuft of long hyaline setae at the apex of the projection. Their function is to give buoyancy to the colonies when living under pelagic conditions. These hyaline setae are sometimes observed on <u>P.duplex</u> (type form) but not on the variety <u>clathratum</u>.

<u>P.duplex</u> and its varieties have been recorded from Australia and New Zealand (Bailey 1895, Hardy 1906, West 1909, Thomasson 1973, Haughey 1968).

Pediastrum tetras (Ehrenb) Ralfs Pl 4 figs. 8,9.

Prescott (1962) Pl 50 fig. 6

Sulex (1969) Taf 12 fig. 6

cell dia. 9-13

<u>P.tetras</u> has been recorded from Australia and New-Zealand (Bailey 1893, Hardy 1906, West 1909, Borge 1911, Scott & Prescott 1958, Spencer 1882, Nordstedt 1888). Samples: Lake Hume 3/5(Dam), 10/9(4).

ipres: bake nume y/ y(bam/, 10/)(1/.

Lake Mulwala 10/2, 4/4, 681121/2.

<u>Scenedesmus denticulatus</u> Lagerh. <u>var. denticulatus</u> May 1973 Pl 5 fig 16 May 1973 p. 444.

= Scenedesmus denticulatus Lagerh. (Haughey 1969 fig. 5:3; Philipose

1967 fig. 176 a-c).

cell L 13-14

₩ 6-8

and grow and

teeth L 2-4

Previous Australian and New Zealand records: May (1973, other varieties), Haughey 1969.

Trainor in 1966 regarded <u>Sc. denticulatus</u> as a stable species (May 1973).

Samples: Lake Mulwala 4/4, 7/5.

Scenedesmus obliquus (Turp.) Kuetz. var. dimorphus May 1973 p. 438. = Scenedesmus dimorphus (Turp) Kuetz. (Philipose 1967 fig. 160a,

Haughey 1968 fig, 10:4).

cell L 18-29

₩ 3-6

According to Philipose (1967) 8-celled colonies are always in subalternating series.A linear 8-celled colony was observed(P15 fig. 6).

May (1973) gave a list of previous Australian records of the variety. Haughey (1968) recorded it as <u>Sc. dimorphus</u> from NZ. Samples: Lake Mulwala 10/2, 4/4, 4/6.

Scenedesmus opoliensis Richter Pl 5 figs. 12-15.

Smith 1920 Pl 41 figs. 8-11

May 1973 p. 446

cell L 15-22

W 4-5

spine L 14-23

The navicular shape of the cells, their lateral contact in the median third only, together with the beak-like instead of rounded apex separate this species from <u>Sc. quadricauda</u> (Smith 1920). May (1973) distinguished <u>Sc. opoliensis</u> from <u>Sc. quadricauda</u> by having cells of coenobe join in median third of their length.

W.& G.S. West (1902 p. 197) stated <u>Sc. opoliensis</u> Richter is a synonym of <u>Sc. quadricauda var. opoliensis</u> (Richter) West & West. The record of a tendency of <u>Sc. opoliensis</u> cells to assume a more cylindric form like those of <u>Sc. quadricauda</u> (Haughey 1968) indicates the two species may be forms of the same organism. Only culture experiments would prove the forgoing statement. May (1973) regards <u>Sc. opoliensis</u> as one of the 14 recognised species in Australia. Previous Australia and New Zealand records: May (1973), Haughey (1968). Pl 5 figs. 12,15 are 2-celled coenobia which may be obtained under high B.O.D. loadings (Haughey 1968).

Samples: Lake Hume 5/7(2).

Lake Mulwala 4/4, 7/5.

Scenedesmus quadricauda (Turp) Breb. Pl 5 figs. 8-10

= Scenedesmus quadricaudus (Turp.) Breb. (May 1973 p. 445).

Pl 5 fig. 8 is the type form of <u>Sc. quadricauda</u>. The dimensions are in the range of the type form sland <u>var. maximus</u> s' as given by Smith (1920).

4-celled coenobia L 21-29

W 16-21

Spine L 15-23

Assmall form of <u>Sc. quadricauda</u> (Pl 5 fig. 10) could be either a daughter coenobium of <u>Sc. quadricauda</u> or <u>Sc. quadricauda</u> <u>var. parvus</u> West & West (Smith 1920 Pl 40 fig. 7). The dimensions are in the range given for var. parvus given by Smith (1920).

4-celled coenobia L 16

W 9

Spines_L 11

An 8-celled coenobium (Pl 5 fig. 9) may be <u>Sc. quadricauda</u> <u>var. Westii</u> (Smith 1920 Pl 41 fig. 3). The dimensions are smaller than those given by Smith (1920).

8-celled coenobium L 12-14

W 4

Despite the various variety names recorded in the literature, all <u>Sc. quadricauda</u>- like plants (Pl 5 figs. 8-10) will be grouped in <u>Sc. quadriquada</u>. There are experimentalresults and field observation that <u>Sc. quadricauda</u> change number of cells in a coenobium(2, 4, 8) and the size of cells(Komarek & Ruzicka 1969, May 1973). With these variation in mind it is doutful if <u>Sc. quadricauda</u> is a stable species in the sense that they do not vary greatly with conditions of culturing, an idea of Trainor s'in 1964, 1966 (from May 1973).

<u>Sc. quadricauda</u> have been recorded from Australia and New Zealand (May 1973, Haughey 1968, Maskell 1881). Samples: Lake Hume 2/5(4), 8/8(4).

Lake Mulwala 10/2, 4/4, 7/5.

Scenedesmus quadricauda var. quadrispina (Chodat) G.M.Smith.

The plant depicted in Pl 5 fig. 11 may be <u>Sc. quadricauda</u> <u>var. quadrispina</u> (chodat) G.M.Smith(Philipose 1967 fig.187d). The plant s'dimensions agree with those given by Philipose(1967) but its cells join at the median third of their length.

cells L 11

₩ 6

spines L 4

<u>Sc. quadricauda var. quadrispina</u> has been recorded from Australia as <u>Sc. quadricaudus var. quadrispina</u> (May 1973 p. 446). Sample: Lake Mulwala 4/4. <u>Selenastrum westii</u> G.M.Smith Pl 3 figs. 7-8

The plants are identified as <u>Selenastrum westii</u> because the strongly curved cells have their convex faces toward one another. (Smith 1920). Their dimensions are comparable with those given by Smith 1920 Pl 31 figs. 8-10 p. 132. Cells L 20, W 3. Smith (1920) stated <u>Selenastrum westii</u> had no pyrenoids. Sometimes one pyrenoid per cell was observed in Lakes Hume and Mulwala plants. Prescott (1962, p. 257) was not sure if pyrenoids of this species were lacking.

<u>Selenastrum species</u> are distinguished from <u>Ankistrodesmus</u> <u>species</u> by both vegetative cells (and autospores when liberated) have the characteristic convex faces toward one another in a colony (Smith 1920). Species of <u>Selenastrum</u> should be compared with those of <u>Kirchneriella</u> which have scattered lunate cells within a gelatinous envelope(Prescott 1962 p. 256).

There is no previous Australian record of the species. Samples: Lake Hume 31/5(4),

Lake Mulwala 10/2, 4/4.

<u>Tetraedron gracile</u> (Reinsch) Hansgirg Pl 4 figs. 13,14 Smith 1920 Pl 26 figs. 1-5, p. 122 Philipose 1967 figs. 69 a-c, p. 154

The plants' dimensions are in the range given by Philipose (1967). Wsp 34-36, Wsp 10-11.

There is no previous Australian record. Samples: Lake Hume 1/5(3), 3/5(dam),

Lake Mulwala 12/8.

Tetraedron hastatum (Reinsch) Hansgirg Pl 4 figs. 15,16 Scott & Prescott 1958 fig. 24:16

The plants differ from Smith s'(1920 Pl 25 fig. 18) in having two terminal spines.

Lcp 36

W of base of process 7

The general shape of the cell is similar to that of

Tetraedron limneticum Borge but there is no branching of the processes.

(Smith 1920). One specimen (Pl 4 fig. 16) had just branched suggesting <u>Tetraedron hastatum</u> and <u>Tetraedron limneticum</u> may be forms of the same biological species.

Previous Australian record: Scott & Prescott 1958. Playfair (1916) recorded <u>T. hastatum var. elegans</u> from NSW. Sample: Lake Hume 1/5(3).

<u>Tetraedron limneticum</u> Borge <u>var. gracile</u> Prescott Pl 4 figs. 17,18 Philipose 1967 fig. 73

Lcp 37-38

dia. of base of process 7-8

There is no previous Australian record. This variety differs from the type in being smaller and having bases of the processes adjoining so there is scarcely a cell body (Prescott 1962). Samples: Lake Hume 8/3(Tallan.), 3/5(Dam), 30/5(3).

The mentioned three <u>Tetraedron</u> species may be forms of the same biological species. As a rule in <u>T. limneticum</u> the processes are tetrahedral while in the majority of cases those in <u>T. gracile</u> are in the same plane (Thomasson 1965) but Thomasson (1965) found a plant which looks like <u>T. gracile</u> but has 6 processes which are directed from everywhere. (fig. 3:6) He called his plant <u>T. limneticum</u>. <u>Tetraedron regulare</u> Kuetz. Pl 4 fig. 20 Prescott 1962 Pl 60 fig. 25 p. 269 Philipose 1967 fig. 60a p. 145

cell dia. 21

Previous Australian records: Schmidle 1896, Scott & Prescott 1958.

Sample: Lake Mulwala 10/2.

<u>Tetraedron Victorieae</u> Wolosz. <u>var. major</u> G.M.Smith Pl 4 fig. 19 Smith 1920 Pl 24 figs. 21, 22 Philipose 1967 fig. 63d

Lcp 42

W 20

There is no previous Australian record.

Sample: Lake Hume 5/4(Dam), 2/5(4), 3/5(Dam).

Treubaria triappendiculatum Bernard Pl 3 fig. 17

Philipose 1967 fig. 32 p. 107

The spines were observed to vary from 3-4. The dimensions are comparable to those given by Philipose (1967).

cell csp 39-41 in dia.

spine 32-34 long and 2-3 broad at base.

Playfair (1918) recorded <u>Bernardia tetraedrica</u> n. sp. (Pl VII fig. 12) from NSW. His plant was slightly smaller than <u>Treubaria triappendiculatum</u>. Playfair s'plant may be the same as the Hume and Mulwala plants.

Sample: Lake Hume 2/5(4),

Lake Mulwala 10/2.

An identified plant PL 3 fig. 18

The filaments were arranged in a tetrahedron structure. Samples: Lake Hume 8/3(Tallan.), 30/5(3), 10/7(3), 7/8(3), 8/8(4).

Lake Mulwala 12/8.

Order Zygnematales

Family Zygnemataceae

Cf. Mougeotia sp. Pl 5 fig. 4

The plants may be <u>Mougeotia sp</u>. because of the band-like chloroplasts. No drawings of any plants were found to match the plants.

cell I 18-36

₩ 3-4

To identify <u>Mougeotia sp.</u> one must know its mode of reproduction (by conjugation or by aplanospores or both), structure and position of the zygospore formed, and finally the size of the vegetative cells(Prescott 1962).

Samples: Lake Hume 2/5(4), 9/9(3).

Spirogyra sp. 1 Pl 5 fig. 3

The curvature of the cell tempts one to identify it as <u>Spirotaenia condensata</u>. Since a lot of workers (West & West 1904 Pl 2 figs. 7-8, Krieger 1937 Taf 2 fig. 1, Bourrelly 1966 Pl 84 fig. 9, Prescott 1966 Pl 11 fig. 8) depicted <u>Spirotaenia condensata</u> as having much more rounded ends and a thicker band of chloroplasts than the Mulwala plant, the plant(Pl 5 fig. 3) is a cell of <u>Spirogyra species</u> s'filament. Reproductive structures (conjugation tubes, zygospores) are needed to identify the species.

- L 151
- W 16

Sample: Lake Mulwala 10/2.

Spirogyra sp. 2 Pl 5 fig. 2

- L 162
- W 41

Samples: Lake Hume 10/7(3).

Lake Mulwala 681121/2.

Family Desmidiaceae

Arthrodesmus convergens Ehrenb. Pl 5 fig. 17

= Staurodesmus convergens (Ehrenb.) Teiling 1948 var. Laportei Teil. West & West 1912 Pl CXVI figs. 7-12, p. 107 Smith 1924 Pl 84 fig. 6 p. 126 Frere-Irenee-Marie 1939 Pl 60 fig. 1 Teiling 1967 Pl 25 fig. 4 p. 588 Lsp 60, Wsp 55, spines L 10, Ist. 14

Previous Australia and New Zealand records: Borge 1896, Hardy 1906, Maskell 1883, 1889, Nordstedt 1888.

Sample: Lake Mulwala 680921/1.

Geographical distribution: widespread (West & West 1912).

Arthrodesmus constrictus G.M.Smith var. longispinus (Gronbl.) Pl 7 figs. 14,15

Apart from having more incurved spines the Hume and Mulwala plants resemble <u>Arthrodesmus constrictus var. longispinus</u> (Scott & Prescott 1961 Pl 36 fig. 2).Their dimensions are comparable.

Lssp 29-33, Wcsp 39-62, Ist. 5-6, Spines L 7-22 Previous Australian record: none

Samples: Lake Hume 14/2(Dam), 2/5(4), 3/5(Dam), 8/3(Tallan.).

Lake Mulwala 4/4, 7/5.

Geographical distribution: The Amazon Basin, Brazil (Scott, Gronblad, & Prescott 1965) Wisconsin, Canada(Smith

1921 forma).

<u>Closterium aciculare</u> T. West <u>var. subpronum</u> West & West Pl 6 fig. 1 West & West 1904 Pl XXIII figs. 4,5 p. 175

L 371-577-766

Apex 2

In this variety cells commenly more elongate than in the type, very slightly sigmoid (West & West 1904). Previous Australian records: West 1909, Scott & Prescott 1958 fig. 7:16, Thomasson 1973, Prescott & Scott 1952 forma). Samples: Lake Hume 8/3(Tallan.), 1/5(3), 2/5(4), 3/5(Dam.),31/5(4),

3/6(5), 5/7(2), 8/7(5).

Lake Mulwala 4/4, 7/5, 4/6.

Geographical distribution : Germany, New Zealand (West & West 1904).

W 5-7
Closterium ceratium Perty Pl 3 fig. 3

The closest plant for Pl 6 fig. 3 is <u>Closterium ceratium</u> (Pl XXIII figs. 6,8 West & West 1904). The dimensions are in the range given by West & West (1904) who also stated that this species' apices are the sharpest ones met with in the whole genus.

cell L 210, W 5

There is no previous Australian record. Sample: Lake Mulwala 4/4.

Geographical distribution: England, France, Germany, Switzerland, Austria & Galicia, Sweden (West & West 1904).

Closterium gracile Breb Pl 6 fig. 4

The morphology of Pl 6 fig. 4 resembles most to <u>Closterium gracile</u> Breb. (Pl XXI fig. 10 West & West 1904 p. 166, Krieger 1939 Taf. 30 figs 7-9). The dimensions are within the range given by West & West (1904).

cell L 188, W 5

Previous Australian records: Schmidle 1896, Bailey 1893, 1895 Pl X

fig. 18, Hardy 1906, Borge 1911,

Playfair 1914, 1915, Prescott & Scott 1952).

Sample : Lake Mulwala 4/4.

Geographical distribution: widespread (West & West 1904).

Playfair (1915) stated Brebisson s'plant had blunt ends. The shape of apices does vary. Frere-Irenee-Marie (1939) figured two <u>Closterium gracile</u> with very pointed apices. (Pl 3 figs. 15,16). Gronblad & Croasdale (1971) figured <u>Cl. gracile</u> with narrow apices than the Mulwala plant s'.

Closterium Kutzingii Breb Pl 6 fig. 2

West & West 1904 Pl XXV fig. 19 p. 186.

L 469, W 16, Apex 2

Previous Australia and New Zealand records: Mobius 1892, Schmidle 1896, Bailey 1898, Hardy 1906, West 1909, Prescott & Scott 1952, scort + prescott A1958, Thomasson 1973, Norstedt 1888, and Haughey 1969 who noted that a few cells from a polluted farm pond were noticeably more curved towards the extremities.(figs. 5, 5a).

Samples: Lake Hume 2/5(4), 3/5(Dam), 3/6(5), 9/9(3).

Lake Mulwala 12/8.

Geographical distribution: widespread (West & West 1904).

Closterium moniliferum (Bory) Ehrenb. Pl 6 figs. 5,6

West & West 1904 Pl XVI figs. 15,16

Krieger 1937 Taf. 18 figs 6,7

Smith 1924 Pl 52 fig. 10

Bicudo 1969 Tab. 117 fig. 38

L 201-260, W 38-46

Although West & West (1904) stated <u>Cl. moniliferum</u> had broad and rounded ends and one central row of pyrenoids, the species with acute ends and ones that had many rows of pyrenoids had been reported (Frere-Irenee-Marie 1939 Pl 5 fig. 2,1 and Prescott 1966 Pl 2 fig. 6 respectively).

Since cell dimensions are variable and the Hume & Mulwala plants sometimes were seen to exhibit one character each of <u>Cl.Leibleinii</u> (acute apices) and <u>Cl.Ehrenbergii</u> (many rows of pyrenoids). The three taxa may be forms of one species as suggested by Playfair 1914. Playfair (1914) did not provide drawings and description of the zygospores of <u>Cl.Ehrenbergii</u> and <u>Cl. moniliferum</u> which he claimed to be identical (p. 100).

Previous Australian and New Zealand records: Playfair 1914, Maskell 1883, Spencer 1882, Norstedt 1888. Samples: Lake Hume 30/5(3), 9/7(3), 10/7(3), 7/8(3), 8/8(4), 9/9(3).

Lake Mulwala 12/8.

Distribution: widespread (West & West 1904, Claassen 1961). Closterium striolatum Corda Pl 6 fig. 7

The truncate ends, the dimensions and the number of striation on the wall of the Hume plant agree well with <u>Cl. striolatum</u> (Krieger 1937 Taf. 28 fig. 9, Thomasson 1967 fig. 4:5). The Hume plant, Pl 6 fig. 7, has wider cell than Krieger s'drawing. However judging from the dimensions given by Krieger, his plant may be wider than the Hume s'.

L 230-330, W 29-30, Apices 9, 14 striation across wall.

Although West & West (1904) stated it exhibited little variation as far as the cell s'shape was concerned Bailey (1898) figured <u>Cl.striolatum</u> with obtuse round ends (Pl VII figs. 1a,b) and Borge (1903) figured <u>Cl. striolatum</u> which curved more than Wests'plants(Tab. 1 fig. 12).

<u>Cl.striolatum</u> Corda have been recorded from Australia & New Zealand (Raciborski 1892, Bailey 1898, Hardy 1906, Harrop 1869, Spencer 1882, Maskell 1881, 1883, Prescott & Scott 1952).

Based on the dimensions and the morphology given by Rich (1932) <u>Cl.Hutchinsonii</u> sp. nov.(figs. 5 d-g) are <u>Cl.striolatum</u>. Rich (1932) was not justified in creating this new species on the basis of minor morphological variations. It was stated <u>Cl. Hutchinsonii</u> had straighter inner margin and narrower apices.

Samples: Lake Hume 7/8(3), 10!9(4).

Distribution: widespread (West & West 1904).

Closterium Venus Kutz. var. inflatum Claassen 1961 Pl 6 fig. 8

Pl 6 fig. 8 was an empty cell with a distinct inflation in the mid-cell region. Apart from being more curved this specimen resembles Cl.Venus var inflatum (Tab. 6 fig. 17 Claassen 1961). L 54-153, W 9-13

They are longer than Claassen (1961) s'plants(L 68-70, W 9-10). Specimens which curved less than the drawing were also observed.

West (1909) and Maskell (1883) recorded <u>Cl.Venus</u> from Australia and New Zealand. The Hume plants also resemble <u>Cl.dianae var. compressum</u> (Kriger 1937 Taf. 19 fig. 14) but the latter are 2-3 times longer. Maskell (1883) noted <u>Cl.Venus</u> and Cl.dianae are similar and the former are smaller. It was proposed that sexual morphology be an essential feature in the identification of taxa within the <u>Cl.Venus-Cl.diana</u> complex(Cook 1963).

Sample: Lake Hume 8/8(4).

Geographical distribution: <u>Cl. Venus</u> are widespread(West & West 1904). Cosmarium bipunctatum Borgesen Pl 5 figs 22-24

West & West 1908 Pl LXXXV fig. 6

Forster 1969 Taf. 16 fig. 12

The presence of the two mid granular structures suggests <u>Cosmarium bipunctatum</u>. The Hume plants differ from the above authors' in having apecies devoid of granules and being slightly bigger.

L 25-27, W 21-23, Ist. 7-8

Pl 5 figs. 22, 24 are probably <u>Cos.bipunctatum</u> because they are similar to <u>Rig. 23</u> which has been identified as <u>Cos.bipunctatum</u>. From the end view the plant (fig. 22) seems to possess central granular structures but the front view does not review them.

There is no previous Australian record. Samples: Lake Hume 8/3(Tallan.), 2/5(Tallan.). Geographical distribution: Brazil, England, (West & West 1908)

e e ante transforme parties de

ພະຍຸການ

Sao Paulo (Borge 1918).

Cosmarium contractum Kirchn. Pl 7 fig. 6 West & West 1905 Pl LXI figs. 23-25 p. 170 Gronblad 1960 Pl XIV fig. 27 Scott & Prescott 1961 Pl 27 fig. 4

Another similar plant is <u>Spondylosium monoliforme</u> Lundell. Since the Hume plant consists of two cells and not a string of them, a character of <u>Spondylosium sp</u>.(West & Carter 1923), it may be <u>Cosmarium contractum</u>.

L 28-32, W 22-23, Ist. 10

Previous Australian records: Hardy 1906, West 1909. Sample: Lake Hume 7/3(3).

Geographical distribution: widespread (West & West 1905). <u>cosmarium granatum Breb.</u> <u>f. pyramidal</u> Schmidle P1 5 fig. 19 Gronblad, Scott & Croasdale 1964 Pl IV figs. 76, 77 Forster 1965 Taf. 5 fig. 4, Taf 11 fig. 29 Prescott 1966 Pl VIII fig. 46

The plant s'dimensions are in the range as those given by Forster (1965).

L 43, W 27, Ist. 7, Apex 7.

Bailey (1898) recorded <u>Cos.granatum var. gibbosum</u> Schmidle 1896(Pl 2 figs. 8a, b) which also looks like the Hume plant. It is possible that <u>Cos. granatum var. gibbosum</u> and <u>f. pyramidal</u> are the same plant.

Gronblad (1960) stated <u>Cos.granatum</u> Breb. is a most variable species. In 1942 he gave various forms of <u>Cos. granatum</u> (Taf. II figs. 22-25). They differed in size, the shape of sinus, the extent of cell wall granulation. Bailey (1898) gave no description for <u>Cos. supergranatum</u> n. sp. (Pl III fig. 5). His new species looks so much like the Hume plant they may be the same biological species. The Hume plant differs from the type in not having the sinus apical dilation as depicted in West & West 1904 Pl LXIII figs.1-4.

Previous Australian and New Zealand records: Borge 1896, Hardy 1906, West 1909, Borge 1911, Maskell 1881, Nordstedt 1888. Sample: Lake Hume 30/5(3).

Geographical distribution: widespread (West & West 1905, Claassen 1961). Cosmarium magnificum Norstedt Pl 5 fig. 20

The outline, the large and small pores in the mid region of each semicell and the size are similar to those of <u>Cos. magnificum</u> Norstedt (Norstedt 1888 Tab. 6 fig. 19 p. 62). The only difference is the shape and the width of the sinus. Nordstedt s'plant has narrow sinus with dilated ends.

L 118, W 91, Ist. 46

The spices of some Mulwala specimens are smooth like those of Nordstedt s'plant while others'are not. Skuja (1949) figured <u>Cos.magnificum</u> Nordst. <u>var. subcirculare</u> Skuja (Taf. XXXII fig. 8) with not smooth apecies.

Thomasson (1965) gave a photo of <u>Cos.decoratum</u> which is similar to <u>Cos. magnificum</u> in outline but is about 30 μ smaller in length and width. His photo was so dark any wall structure (if any) could not be seen(photo h fig. 9:8). <u>Cos. magnificum</u> seems closely related to <u>Cos. multiordinatum</u> West & West <u>var. africanum</u> Bourrelly (Bourrelly 1957 Pl 11 fig. 96) which, according to Lind (1967 Pl 6 fig. 4) have the big and small granules confined to the central area.

There is no previous record of <u>Cos. magnificum</u> in Australia. Nordstedt (1888) first reported it from New Zealand. Maskell (1889 p. 6) stated <u>Cos. gemmiferum</u> is <u>Cos. magnificum</u>. His 1883 drawing of <u>Cos. gemmiferum</u> (Pl XXIV fig. 4) showed neither the big and small central pores nor the characteristic outline of <u>Cos. magnificum</u> which is truncate. He stated Cos. gemmiferum was a big desmid but gave

38

no dimensions.

Samples: Lake Mulwala 4/4, 680508/1

Geographical distribution: Australia & New Zealand and perhaps

Africa (as Cos.multiordinatum West &

West var. africanum Bourrelly).

Cosmarium pseudonitidum (cf.) Nordst. Pl 7 figs. 19-22

The closest plant that looks nearly like the Mulwala plants in all views is <u>Cosmarium pseudonitidum</u> Nordstedt(Nordstedt 1872 fig. 4 p. 46). The Mulwala plants are smaller than Nordstedt s' plant whose dimensions are in brackets.

> L 26-32 (42) W 24 (33) Ist. 6 (10)

Mucilagenous strands were sometimes observed around the plants (Pl 7 fig. 22). Nordstedt (1872) did not mentioned any mucilagenous strands.

There is no previous record of the species. Sample: Lake Mulwala 4/4.

Cosmarium punctulatum Breb. var. subpunctulatum (Nordst.) Borgesen Pl 5 fig. 21

West & West 1908 Pl LXXXIV figs. 15-20

Smith 1924 Pl 57 fig. 29

Forster 1965 Taf. 7 fig. 10

Smith (1924) described this variety as "semicells with granules of central area conspicuously larger than those near the margin. The arrangement of granules at centre generally not following a definite pattern".

L 49, W 37

Gronblad (1926 Taf. II figs. 75,76) Cos.multiundulatum n. sp.

resemble the Hume plants. His new species is probably <u>Cos. punctulatum</u> <u>var. subpunctulatum</u>. Norstedt (1888) raised this variety to a specific rank (Tab. 5 fig. 8).

Previous Australian and New Zealand records: Borge (1896, 1911, Scott & Prescott 1958 fig. 14:7, and Mobius 1892, Hardy 1906, West 1909, Bailey 1893 as <u>Cos. subpunctulatum</u> (Breb.) Nordstedt. Samples: Lake Hume 8/3(Tallan.), 2/5(Tallan.).

Geographical distribution: widespread (West & West 1908).

Cosmarium quadrum Lund Pl 5 fig. 18

West & West 1912 Pl C figs. 3-6, p. 21

Thomasson 1965 fig. 9:5

L 65, W 64

The dimensions are in the range given by West & West 1912. From the end view the intermingle of large and small granules was observed only in the mid region.

Previous Australian and New Zealand records: Bailey 1898, West 1909, Flint 1966.

Samples: Lake Mulwala 4/4, 680411/1.

Geographical distribution: widespread (West & West 1912).

Euastrum anasatum (Ehrenb.) Ralfs Pl 7 fig. 8

Krieger 1937 Taf 58 fig. 1 p. 484

The Mulwala Plant dimensions are within the range given by Krieger (1937).

L 85, W 40, Its. 12

The facial swellings on each semicell were not observed. Their number and disposition along with the proportion of semicell base and apex do vary (Prescott & Scott 19(6).

Previous Australian records: Harrop 1869, Mobius 1892, Bailey 1893, Borge 1896, 1911, Maskell 1883, Norstedt 1888). Hardy (1906) doubted if <u>Euastrum affine</u> Ralfs could be <u>Euastrum anasatum</u> Ralfs. From Pl 35 figs. 11,12 West & West 1905 of <u>E.affine</u> the morphology of these two plants are quite different.

West & West (1905, Pl XXXVI fig. 10) and Frere-Irenee-Marie (1939 Taf. 40 fig.8) recorded with one lateral swelling on each side of a semicell and called <u>E.anasatum</u>. Sometimes such a plant is called <u>E.obesum var. crassum</u> (Pl 1 fig. 23 Prescott & Scott 1942). <u>E.cuneatum</u> (Bourrelly 1957 Pl 2 fig. 14) also resembles the Mulwala plant. Perhaps all these taxa refer to the same biological species.

Sample: Lake Mulwala 4/4.

Geographical distribution: widespread (West & Carter 1923,

Claassen 1961).

Euastrum cuspidatum var. goyazense Forster 1969 Pl 7 figs. 9-11

The closest plant that looks nearly like the Hume and Mulwala s'is <u>Euastrum cuspidatum var. goyazense</u> Forster (Forster 1969 Taf. 9 fig. 20). Their sinus shape is the same. The Hume and Mulwala plants differ from Forster s'plants in having three pyrenoids per semicell instead of one, and they have less number of spines on polar lobes. Their dimensions are comparable.

Lcsp 22-26, Wcsp 23-29, Ist. 5

Forster (1969) stated its basinonym was <u>Euastrum subtile</u> Borge <u>var. goyazense</u> (Forster 1964 Taf. 40 fig. 19), its synonym was <u>Euastrum cuspidatum Wolle var. subtile</u> (Borge).

There is no previous Australian record but this plant has been seen in Lake Sorell, Tas. (Dr. P.A.Tyler pers. comm.). Samples: Lake Hume 5/4(Dam), 1/5(3), 3/6(5), 8/7(5).

Lake Mulwala 4/4, 7/5, 4/6.

41

Geographical distribution: Brazil (Scott, Gronblad & Croasdale 1965) as <u>Euastrum cuspidatum Wolle var. subtile</u>, (Borge 1918) as <u>E. subtile</u>,

North Carolina (Schumacher et al 1966)

as Euastrum cuspidatum Wolle.

Euastrum divergens Josh. Pl 7 fig. 12

The Mulwala plant resembles closely <u>Euastrum divergens</u> Josh. (fig. 5:2 Scott & Prescott 1958). The original plant of Joshua s'(1886 Pl 23 fig. 9) had more pointed spines on each lobe and its sinus is wider open.

L 51-60, Wcsp 48-56, Its. 7-12

There are many varieties of <u>Euastrum divergens</u> because of the variation in the pattern, length and the number of terminal spines on each lobe, the shape of the central protuberance.

Previous Australian records: Scott & Prescott (1958), Borge (1896) recorded <u>Euastrum divergens var. australianum</u> Borge (Tab II fig. 15) from Queenland. His plant had fewer spines which were more pointed and a more open sinus when compared with the Mulwala plant.

The Mulwala <u>Euastrum divergens</u> were seen to have spine ornamentation on each semicell lobe.

Sample: Lake Mulwala 10/2.

Geographical distribution: Burma (Joshua 1885), Africa (Lind 1967),

Australia.

<u>Euastrum elegans</u> (Breb.) Kutz. <u>var. elegans</u> forma Pl 7 fig. 7 West & Carter 1923 Pl 38 fig. 16 Gronblad, Scott & Croasdale 1964 Pl IX fig. 210 p. 13

L 31, W 19, Ist. 5

These dimensions are comparable with those given by the above authors.

Previous Australia and New Zealand records: Scott & Prescott 1958, Maskell 1881, Nordstedt 1888. Geographical distribution: widespread (West & Carter 1923). Sample: Lake Hume 3/5(Dam). <u>Gonatozygon kinahani</u> (Archer) Rabenh Pl 7 figs. 2,3 Frere-Irenee-Marie 1939 Pl 66 fig. 25

Forster 1969 Taf. 1 fig. 9

West & West (1904) showed the chloroplast bands of <u>Gonatozygon kinahani</u> were not continuous as seen in those of the Hume and Mulwala and Forster s'plants.

L 185-198, W 9-13, pyrenoids 10-12 per chloroplasts band.

The dimensions are in the range given by the above authors.

West & Carter (1923) and Krieger (1937) figured <u>Roya</u> <u>cambrica</u> West <u>var. limnetica</u> West (Pl CLXVI fig. 14 p. 260 and Taf. 5 fig. 10 respectively) of the same morphology but of smaller size when compared with the Hume and Mulwala plants. The two taxa seem closely related.

<u>Gonatozygon kinahani</u> have been recorded from Australia and New Zealand(West 1909, Playfair 1912, 1914, Flint 1966). Samples: Lake Hume 14/2(Dam), 2/5(4), 9/9(3), 681122/3.

Lake Mulwala 10/2.

Geographical distribution: Canada (Frere-Irenee-Marie 1939),

Central Africa (Forster 1969),

Italy, Siam(var.), U.S.A.(Wests 1904).

Gonatozygon monotaenium De Bory Pl 7 fig. 4

West & West 1904 Pl 1 figs. 1-7 p. 30

Forster 1965 Taf. 1 fig. 4

The dimensions fall into the range given by West & West 1904. L 117, W 8, apex W 9. Previous Australian and New Zealand records: Hardy 1906, West 1909, Flint 1966.

Sample: Lake Hume 14/2(Dam).

Geographical distribution: widespread (West & West 1904, Claassen 1961). <u>Micrasterias Hardyi</u> G.S. West Pl 6 fig. 9

West 1909 Pl 5 fig. 1, p. 57

Lcp 202-242 Lsp 102-108 Wcp 213-248 Ist. 20-27

Polar lobe process length 0-7

The variations within the population were observed to be 1) The sinus shape. This may be wider open than that of the drawing (Pl 6 fig. 9).

2) The furcation along the length of each process may be longer or shorter than that of the drawing (Pl 6 fig. 9).

3) The polar lobe processes may not be present.

An empty cell was found and all the wall ornamentation was as shown in the drawing.

Previous Australian records: West 1909, Tyler 1970.

Dr. Tyler (pers. comm.) thinks this species may be an endemic form of Australia. I think may be it is an extreme form of the cosmopolitan <u>Micrastrias mahabuleshwarensis</u> Hobson because some reported drawings of <u>M.mahabuleshwarensis</u> could become those of <u>M. Hardyi</u> if the processes are longer or shorter, etc. Gronblad (1945) depicted <u>M. mahabuleshwarensis</u> with shorter processes... otherwise the plant becomes <u>M.Hardyi</u>. (Tab IV fig. 30). Kreiger s;(1939) <u>M.mahabuleshwarensis</u> (Taf. 109 fig. 10) looks like <u>M.Hardyi</u> with shorter processes and long polar lobe processes. <u>M.mahabuleshwarensis</u> var.europaea Nordst.(Krieger 1939 Taf.110 fig. 5) may become <u>M.Hardyi</u> if the polar lobe processes are reduced and their processes become longer with deeper incision of lateral lobules.

A genetic link of <u>M.Hardyi</u> with <u>M.tropica var. indivisa</u> (Nordst.) Eichl et Racib via <u>M.berganii</u> Hauge had been suggested (Tyler 1970).

Samples: Lake Hume 14/2(Dam), 7/3(3), 8/3(Tallan.), 5/4(Dam), 1/5(3), 2/5(4), 3/5(Dam), 31/5(4), 3/6(5), 8/7(5), 9/7(3), 7/8(3), 681122/3.

Lake Mulwala 4/4.

Geographical distribution: Tas., Vic., NSW, (Australia-Tyler 1970).

Micrasterias decemdentata (Nag.) Arch. Pl 6 fig. 12

Bailey 1898 Pl 2 fig. 18

Krieger 1939 Taf. 104 fig. 8 p. 34

Bourrelly 1957 Pl 4 fig. 38, Pl 5 figs. 39-40

Claassen 1961 Tab. 16 fig. 5 p. 585

Gronblad & Croasdale 1971 Pl IV fig. 47 p. 11

Tyler (1970) included the Mulwala plant in <u>M.laticeps</u> <u>var. minor</u> Nordst. I think the plant may be <u>M.decemdentata</u> because it differs from Nordstedt s'original plant (1869 Taf. 2 fig. 14 p.220). and Bicudo et al (1972) who studied polynorphism in <u>M.laticeps</u> did not record any form with lateral branched lobules like the Mulwala plant. The Mulwala plant agrees in shape and size with M.decemdentata (Bailey 1898 Pl 2 fig. 18). The other mentioned workers reported <u>M.decemdentata</u> of 30_M smaller to about half the size of the Mulwala plant.

L 95, W 100

Playfair (1912) recorded <u>M.truncata</u> mixed form(Pl lvi fig. 3) with one semicell of <u>M.decemdentata</u> and another semicell Previous Australian records: Mobius 1894, Bailey 1895, 1898, Schmidle 1896, Borge 1911, Tyler 1970 as <u>M.laticeps var. minor</u> Sample: Lake Mulwala 10/2.

Geographical distribution: seem widespread.

<u>Micrasterias mahabuleshwarensis</u> Hobson typica-wallichii transition Pl 6 fig. 11

Tylor (1970) fig. 17 e

Lep 167, Lsp 127, Wep 146.

The wall ornamentation was not shown in the drawing because the ornamentation was hard to observe& against the cell contents.

Previous Australian records: Mobius 1894, Tyler 1970.

West & West (1905) described the variation within <u>M.mahabuleshwarensis</u> which had resulted in the naming of numerous varieties.

Samples: Lake Mulwala 4/4, 690206/1.

Geographical distribution: This species is principally tropical and subtropical in its distribution (West & West 1905).

<u>Micrasterias thomassiana var. notata</u> (Nordst.) Gronblad Pl 6 fig. 10 West & West 1905 Pl 50 fig. 7 as <u>M.denticulata var. notata</u> Nordst. Krieger 1939 Taf. 140 fig. 1 p. 111

Scott & Prescott 1961 Pl 17 fig. 6

Flensburg 1967 Pl IIIc

Tyler 1970 fig. 29a

L 251, W 210, Ist. 38.

The dimensions are comparable with those given by the above authors. The plant also closely resembles the South African plant <u>M.denticulata</u> Breb.var. subnotata West(Claassen 1961 Tab.12 fig. 2, Tab.13 fig. 1) which is synonymous with <u>M.thomassiana var</u>. notata (Krieger 1939). Krieger (1939) gave various synonyms of the taxon. The intergradation of <u>M.denticulata</u> and <u>M.thomassiana</u> has been mentioned (Tyler 1970).

Previous Australian record: Tyler (1970), Prescott & Scott-Sample: Lake Hume 7/8(3). 1952).

Geographical distribution: England, Germany, Sweden, Finland, NZ,

(West & West 1905), Tasmania & Victoria (Tyler 1970), Indonesia (Scott & Prescott 1961), Canada (Stein & Garrath 1968).

<u>Pleurotaenium Ehrenbergei</u> Breb. De Bory var. undulatum Schaarschmidt. Smith 1924 Pl 54 fig. 9 p. 15

> L 444-523 W 31-42 Apex 24 Ist. 33

<u>Pleutaenium Ehrenbergei</u> have been recorded from Australia and New Zealand (Mobius 1894, Schmidle 1896, West 1909, Nordstedt 1888). This variety <u>undulatum</u> differs from the type in having the undulation more or less throughout the whole semicell s' length (Smith 1924). Prescott & Scott (1952) recorded the variety. Sample: Lake Hume 30/5(3).

Geographical distribution: widespread (Claassen 1961).

Spondylosium planum West & West Pl 7 fig. 5

West & Carter 1923 Pl CLX figs. 23-24

Gronblad 1926 Taf. 3 fig. 123

Frere-IreneeMarie 1939 Pl 61 fig. 17

Prescott & Scott 1960 Pl 60 figs. 6-8

L 14, W 14, Ist. 7.

There is no previous Australian record. Flint (1966) recorded it from New Zealand. Samples: Lake Hume 14/2(Dam),

Lake Mulwala 680411/1, 681121/2, 690202/1 Geographical distribution: New Zealand (Nordstedt 1888). Australia, West & Carter (1923) : Norway, U.S.A.,

Canada.

Staurastrum anatoides Scott & Prescott Pl 10 fig. 1

The Mulwala plant is a form of <u>Staurastrum anatoides</u> This taxon was named by Scott and Prescott (1958). Scott & Prescott (1961) described <u>St.anatoides var. javanicum</u> (Pl 56 fig. 1) whose end view is almost identical to that of the Mulwala plant. The front view differs slightly. Scott & Prescott s'semicell has a row of granules above the isthmus while the Mulwala plant has two long isthmal spines per semicell. Their dimensions are comparable. I think <u>St. quebecense</u> Frere-Irenee=Marie (Frere-Irenee-Marie 1939, Pl 47 fig. 6 and Pl 54 fig. 5) may be the same biological species as <u>St. anatoides</u> Scott & Prescott. Their dimensions are comparable and on focusing up the curved lines in the middle and at the base of each process of the semicell from the end view as depicted by Frere-Irenee-Marie (1939 Pl 47 fig. 6) become bifurcate verrucae

Previous australian record: Scott & Prescott 1958. Sample: Lake Mulwala 690206/1. (L 44, Wcp 73, Ist. 11). Geographical distribution: Australia

> Indonesia (Scott & Prescott 1961) Perhaps Canada as <u>St.quebecense</u> (Frere-Irenee-Marie 1939).

<u>St.arctiscon</u> (Ehrenb.) Lund <u>var. glabrum</u> West & West Pl 10 figs. 7,8 Smith 1924 Pl 84 figs. 3-5, Pl 83 fig. 15, p. 125 Frere-IreneeMarie 1939 Pl 57 fig. 3

St.arctiscon var. glabrum have smooth processes. There are

two sizes (Pl 10 figs. 7,8). As a rule, there are six processes in the upper whorl and nine in the lower whorl (Thomasson 1973). The processes may be either bifurcate or trifurcate (Frere-Irenee-Marie 1939, Thomasson 1973).

> Lsp 34-57 Lcp 62-102 Wsp 26-37 Wcp 57-95 Ist. 14-25

<u>St.arctiscon var. glabrum</u> and <u>St.leptacanthum</u> Nordstedt are similar plants (Frere-Irenee-Marie 1939 Pl 48 fig. 2, Pl 58 figs. 10,11). The latter have fewer processes, 4 in the upper whorl, 6 in the lower whorl (Frere-Irenee-Marie 1939). Forster (1969) <u>St.leptacanthum var. Borgėi</u> morpha minor is <u>St.arctiscon var</u>. <u>glabrum</u>. Gronblad (1947 fig. 37) <u>St.tohopekaligense</u> is also <u>St.arctiscon var. glabrum</u> judging from the end view.

<u>St.arctiscon var. glabrum</u>, <u>St.leptacanthum</u>, and <u>St.tohopekaligense</u> seem to be three well-defined morphological species to me.

Thomasson (1973) stated the number of arms of <u>St.arctiscon</u> <u>var. glabrum</u> varies. He reported the species with only three processes in the lower whorl. This plant morphology of <u>St.arctiscon var. glabrum</u> seems odd.

Previous Australian and New Zealand records: Thomasson (1973, smalle form), Flint 1966. Samples: Lake Mulwala 4/4, 680411/1, 680508/1, 690206/1. Geographical distribution: seem widespread. St.asterias Nygaard Pl 9 fig. 12

Nygaard (1926) Pl VI fig. 62 p. 232

The end view of the Mulwala plant agrees well with that of Nygaard s'original plant. As for the front view the Mulwala plant has more convex apecies. Their dimensions agree.

Lsp 31, Wcp 39, Ist. 11.

Previous Australian records: none Sample: Lake Mulwala 4/4.

Geographical distribution: Australia, Japan (Hinode 1966 Pl 9 fig. 5), Malaya (Nygaard 1926).

Starastrum dickiei Ralfs Pl 9 figs. 1-5

= <u>Staurodesmus dickiei</u> (Ralfs) Lillier (Teiling 1967 fig. 29:3, p 598). West & Carter 1923 Pl CXXIX fig. 14 p. 3-4

Frere-Irenee-Marie 1939 Pl 44 fig. 10 p. 275

L 33-38

Wap 33-43

Ist. 8-11

According to Gronblad, Scott & Croasdale 1964 the plants (Pl 9 figs. 4,5) are <u>St.dickiei var. maximus</u> (Pl XII figs. 277,278). The spines of <u>St.dickiei</u> may be straight or slightly incurved (Claassen 1961).

Previous Australian records: Raciborske 1892, Hardy 1906, West 1909.

Samples: Lake Hume 8/3(Tallan.), 2/5(4), 3/5(Dam), 5/4(Dam), 7/8(3).

a server a server a server server

er negerige

Lake Mulwala 12/8, 680508/1, 680921/1.

Geographial distribution: widespread (West & Carter 1923).

<u>Staurastrum grande</u> Bulnh. <u>var. parvum</u> West Pl 9 fig. 6 = <u>Staurodesmus grande</u> (Bulnh.) Teiling (Teiling 1967 Pl 23 fig. 5) West & West 1912 Pl CXX figs. 2,3

Frere-Irenee-Marie 1939 Pl 45 fig. 4, 1951 Pl 2 fig. 5

L 64, W 61, Ist. 14

The dimensions agree with those given by West & West (1912). The chloroplast appears stellate from the end view.

Previous Australian record: West (1909)-<u>St.grande</u> Bulnh. Sample: Lake Mulwala 680508/1.

Geographical distribution: Australia, Canada (Frere-Irenee-Marie

1939, 1951), Norway & Scotland (Wests 1912). Staurastrum Freemanii West & West 1902 Pl 9 fig. 9

The Hume plant is a form of <u>St.Freemanii</u> West & West 1902 (West & West 1902 Pl 21 fig. 21 p. 177).which has two processes in the end view. The Hume plant s'end view has three processes the base of which has no accessory spines otherwise it is very similar to <u>St.Freemanii var. triquetrum</u> (West & West 1902 Pl 21 fig. 22). The Hume plant dimensions are slightly bigger than <u>St.Freemanii</u> var. triquetrum s' (West & West 1902).

> Lssp 30 Wcsp 63-72

<u>St.Freemanii</u> and <u>St.patens</u> Turner (Pl 9 fig. 10) seem closely related. West & West 1902 stated <u>St.Freemanii</u> differs from <u>St.patens</u> in its smooth lateral margins, its much more produced angles, and in the characteristic pairs of spines at the apices.

Previous Australian records: Scott & Prescott (1958). Sample: Lake Hume 7/3(3).

Geographical distribution: Australia, Ceylon (West & West 1902),

Indonesia (Scott & Prescott 1961, variety). The central highland of New Guinea as <u>St.freemanii facies triquetrum</u> (Brook & Hine 1966). Sturastrum furcatum (Ehrenb.) Breb. Pl 9 figs. 13,14 West & Carter 1923 Pl CLV figs. 1-4 p. 173 Smith 1924 Pl 83 figs. 1-3 p. 118 Frere-Irenee-Marie 1939 Pl 48 fig. 7

The end view of the Hume and Nulwala plants'suggests <u>St.furcatum</u>. From the published figures of <u>St.furcatum</u> and <u>St.tohopekaligense var. brevispinum</u> G.M.Smith (Smith 1924 Pl 82 figs 8-11, Thomasson & Tyler 1971 figs. 10 g,h) the main difference of these two taxa is: from the end view the shorter processes are closer to the longer ones in the former. Since the Hume and Mulwala plants'dimensions also are in the range between those of <u>St.furcatum</u> and <u>St.tohopekaligense var. brevispinum</u>, these two taxa may be synonymous.

> Lcp 38-42 Lsp 28 Wcp 33 Wsp 19 Ist. 9-11

Judging from the almost identical end view morphology and the dimensions of <u>St.furcatum</u> and <u>St.tohopekaligense</u> Wolle given by West & Carter (1923) these two taxa may be varieties of the same biological species. One variety has longer processes than the other. That <u>St.furcatum</u> and <u>St.tohopekaligense</u> may be synonymous had been suggested by Chuah (1969).

Previous Australian records: West (1909), Chuah (1969). Samples: Lake Hume 14/2(Dam), 7/3(3), 8/3(Tallan.), 9/7(3).

Lake Mulwala 680411/1.

Geographical distribution: widespread (West & Carter 1923, Claassen 1961).

Staurastrum leptocladum Nordst Pl 8 figs. 6-11

The plants resemble <u>St.leptocladum</u> (Smith 1924 Pl 78 figs. 6,7). They are smaller and have no subapical spines. Smith (1924) and Scott, Gronblad & Croasdale (1965) stated Nordstedt in his orginal plant did not mention the subapical spines because they may have been lost. Rows of granules at the subapical region were observed (Pl 8 fig. 11).

Lsp	23-32
Lcp	44-61
Wcp	67-92
Ist.	5

One plant with dichotypecal semicells (Pl 8 fig. 7) suggests a relationship with <u>St.nodulosum</u> Prescott (Pl 8 fig. 5). The smaller size and the presence of dichotypical specimens of <u>St.leptocladum</u> and <u>St.nodulosum</u> features suggest <u>St.nodulosum</u> is a small form of <u>St.leptocladum</u>. Perhaps Prescott (1936) should not have created the new species, <u>St.nodulosum</u>. The processes of <u>St.leptocladum</u> are not always parallel or nearly so. Thomasson (1960) recorded <u>St.leptocladum var. elegans</u> with upwardly curved processes (fig. 22) from New Zealand.

Scott & Gronblad (1957) recorded a variety of <u>St.leptocladum</u> which resembles the Hume and Mulwala plants. They did not give any dimensions.

Previous Australian and New Zealand records: West 1909, Thomasson (1960, variety).

Samples: Lake Hume 1/5(3), 2/5(4), 3/5(Dam), 31/5(4), 3/6(5), 8/7(5), 9/8(5), 11/9(5).

Lake Mulwala 4/6.

Geographical distribution: widespread.

Staurastrum nodulosum Prescott Pl 8 figs. 1-5, 12, 13, 15-17 Prescott 1936 Pl LXIV figs. 2,3

The plants fit the description of <u>St.nodulosum</u> (Prescott 1936 p. 506-507). Dichotypical specimens of <u>St.nodulosum</u> semicell and another semicell with upwardly curved processes were often encountered (Pl 8 fig. 3) therefore there are two morphological forms of <u>St.nodulosum</u> in both lakes, namely

 Forms:
 Processes upwardly curved
 Processes divergent

 (Pl 8 figs. 1, 15, 16)
 (Pl 8 figs. 2, 4, 5, 17)

 Lsp
 16-19-24
 15-30

 Lcp
 36-53-68
 39-60

 Wcp
 38-53-67
 44-79

 Ist.
 3-5
 4-6

The variations within the populations are the size, the degree of divergence of the processes and the shape of the sinus (Pl 8 figs. 1-4). There are more specimens of the divergent <u>St.nodulosum</u> in both lakes.

Thomasson (1965) recorded <u>St.caledonense</u> f. major Thomm. (fig. 11:8) with upwardly curved processes and better-developed terminal spines. He suggested his plant was very similar to St.nodulosum. Since some <u>St.nodulosum</u> in Lakes Hume and Mulwala had upwardly curved processes Thomasson s'plant may be grouped in <u>St.nodulosum</u>.

In 1974 Thomasson noted <u>St.chaetopus</u> Hinode from NZ (fig. 5:7,8). His plants especially fig.5:7 looks like <u>St.nodulosum</u> (Pl 8 fig. 15). The descriptions of <u>St.chaetopus</u> given by Smith (1924 Pl 76 figs. 21-24) and Brook (1960 Tab. 89 fig. 1) did not fit Thomasson s'<u>St.chaetopus</u> therefore his plants may be <u>St.nodulosum</u>. The plant (Pl 8 fig. 12) is probably a form of <u>St.nodulosum</u>. Only one specimen was seen. The only clostest plant is <u>St.volans</u> West & West as depicted by Rich (1932 figs. 14 a-d) but <u>St.volans</u> is much smaller and the plant s'dimensions are within the range of <u>St.nodulosum</u> found. Since dimensions vary <u>St.volans</u> may be a form St.nodulosum and hence St.leptocladum.

St.volans-like plant			St.volans in Rich (1932)
	Lsp	23	14-17	
	Lcp	46	not given	
	Wcp	54	32-44	
	Ist.	6	4-6	

Judging from the shape and dimensions, the plant (Pl 8 fig. 13) with well-develope apical verrucae could be grouped in <u>St.nodulosum</u>.

Previous Australian records: Thomasson & Tyler 1970.

Samples: Lake Hume 14/2(Dam), 8/3(Tallan.), 2/4(Tallan.), 5/4(Dam),

1/5(3), 2/5(4), 3/5(Dam), 3/6(5), 5/7(2), 8/7(5), 10/7(3).

Lake Mulwala 10/2, 4/4, 7/5, 4/6.

Geographical distribution: Australia (Thomasson & Tyler 1970),

Panama Canal, Africa (Prescott 1936). Staurastrum Playfairi Scott & Prescott Pl 8 fig. 14

The plants which are of the same size as the divergent form of <u>St.nodulosum</u> but have granules on their semicell wall and an isthmal row of granules (Pl 8 fig. 14) resemble <u>St.Playfairi</u> Scott & Prescott (Scott & Prescott 1961 Pl 52 fig. 4 p. 101).

I agree with Chuah (1969) that Scott & Prescott (1961) should not introduce a new taxon when they stated that <u>St.Playfairi</u> was synonymous with St.volans var. elegans (Playfair 1913).

Chuah (1969) had observed a janus form of <u>St.Playfairi</u> from Wood & lake, Tasmania. Previous Australian record: Chuah (1969).

Samples: Lake Hume 2/5(4), 3/5(Dam), 5/4(Dam).

Lake Mulwala 4/4, 680508/1.

Geographical distribution: Australia, Indonesia (Scott & Prescott 1961).

Chuah (1969) suggested a relation between <u>St.Playfairi</u>-<u>leptocladum-volans</u>. The presence of dichotypical plants of <u>St.nodulosum-leptocladum</u> indicates culture experiments could provide an evidence for grouping <u>St.Playfairi</u>, <u>St.nodulosum</u>, <u>St.volans</u> and <u>St.leptocladum</u> into one biological species.where <u>St.leptocladum</u> may have many varieties. <u>Staurastrum pachyrhynchum</u> Nordstedt Pl 9 figs. 7,8 = <u>Staurodesmus pachyrhynchus</u> (Nordstedt) Teiling (Teiling 1967) West & West 1912 Pl CXXI fig. 8 p. 151

Teiling 1967 p. 499

The plants' dimensions are in the range given by the above authors.

L 30, W 24, Ist. 14.

Sometimes the plants had a spine on each corner.

There is no previous Australian record. Thomasson (1973) recorded <u>Staurodesmus pachyrhynchus</u> that does not look like the Hume and Mulwala plants. He recorded it from New Zealand (figs.4:14-16). Samples: Lake Hume 14/2(Dam), 8/3(Tallan.), 5/4(Dam), 2/5(4),8/7(5).

Lake Mulwala 4/4, 7/5. Geographical distribution: widespread (West & West 1912). Staurastrum patens Turner Pl 9 fig. 10

The Mulwala plant is <u>St.patens</u> Turner (Turner 1892 Tab.14 fig.11.p.108). In the end view the accessory spine at the base of each process as depicted by Turner is not seen. The plant differs from West s'<u>St.patens var.planctonicum</u> in not having two small denticulations at the base of each process (West 1909 Pl 6 fig.14). The Mulwala plant s'dimensions are smaller than Turner s' original <u>St.patens</u> (Turner 1892 Tab.14 fig.11) but are comparable to those of St.patens var.planctonicum given by West (1909).

> Lssp 28 Lcsp 41 Wssp 28-30 Wcsp 47-53 Ist. 14

Previous Australian records: West (1909) recorded <u>St.patens var.planctonicum</u> from Vic. He stated Playfair in 1907 recorded the same variety under the name of "<u>forma australica</u>". Sample: Lake Mulwala 680508/1 Geographical distribution: Australia, East India (Turner 1892). <u>Staurastrum pingue</u> Teiling Pl 11 figs. 1-6 <u>Staurastrum pingue</u> Teiling (small form) Pl 11 figs. 7-13 Teiling 1942 Abb. 3 Brook 1959 Pls XVII & XVIII figs. 2,6,12 Forster 1967 Taf. 10 fig. 11 Chuah 1969 fig. 1 b Thomasson & Tyler 1971 figs. 3 a-d, 7 a

From the literature survey most of the Australian plants (Chuah 1969, Thomasson & Tyler 1971) including those of the Hume s' and Mulwala s' have no apical ornamentation as depicted by Brook (1959) and Forster (1967). Teiling (1942) figured one out of three plants with had no apical ornamentation (Abb.3).

The variations within the two lakes populations are the size, the sinus shape, the degree of process furcation. <u>St.pingue</u> small form occur with the usual ones. Brook (1959) and Forster (1967) gave the detail of various sinus shape.

Pl 11 fig. 3 is a form of <u>St.pingue</u> which approaches <u>St.planctonicum</u> Teiling (Brook 1959 Pl XVIII fig. 12). It is <u>St.pingue</u> stout form according to Thomasson & Tyler (1971). The plants (Pl 11 figs. 1,2,4-6) are the slender form which have bulbous sinus (Chuah 1969).Pl 11 figs. 1,2 was described as <u>St.gracile var. bulbosum</u> by Thomasson & Tyler (1971 fig. 3f). Pl 11 figs. 7-13 are small form of <u>St.pingue</u> formerly known as <u>St.paradoxum var. evolutum</u> West & Carter 1923 (Brook 1959 Pl XVIII figs. 2,6,12). The angles of one semicell alternate with those of the other as seen from the end view (Pl 11 fig. 11). This is one characteristic of <u>St.pingue</u> (Brook 1959).

	small	form	St.pingue
qal	22-23		30-34-40
Wcp	34-56		52-68- 102
Ist.	5		6-12

Previous Australian and New Zealand records: Thomasson & Tyler 1971, Thomasson 1973, Thomasson 1960.

Samples: Lake Hume 14/2(dam), 7/3(3), 8/3(Tallan.), 1/5(3), 2/5(4), 3/5(dam), 30/5(3), 31/5(4), 10/7(3), 8/7(5), 9/8(5), 11/9(5), 681122/3.

Lake Mulwala 4/4, 4/6, 7/5, 680921/1, 681121/2, 690206/1. Geographical distribution: widespread.

Samples: St. pingue (small form) Lake Hume 7/3(3), 8/8(4),

Lake Mulwala 4/4.

Staurastrum cf.pinnatum var. subpinnatum Pl 9 fig. 17

West & West 1902 Pl 21 fig. 33 Skuja 1949 Taf.XXXIV fig. 7 Bourrelly 1957 Pl 17:149

Only one specimen was seen. The end view (pl 9 fig. 17) resembles closely that of <u>St.pinnatum</u> (Turner) <u>var. subpinnatum</u> (Schmidle) given by the above workers. Due to the presence of the cell contents it was not seen if the Mulwala plant had the apical spines as depicted by the mentioned workers. To positively identify the plant the front view is needed. Scott & Gronblad (1957) figured <u>St.comptum</u> Wolle <u>var. pinnatiforme</u> (Pl XXV fig.II) with bifurcate small processes otherwise the end view is similar to the Mulwala plant. Its front view and those of <u>St.pinnatum var.subpinnatum</u> as given by West (1902) and Skuja (1949) are quite different.

Previous Australian record: none. Schmidle (1896) recorded <u>St.subpinnatum</u> which resembles the Mulwala except the small subprocesses do not terminate in three spines. (Taf.IX fig. 20). Samples: Lake Mulwala 4/4.

Geographical distribution: widespread.

<u>Staurastrum pseudosebaldi</u> Wille <u>var. planctonicum</u> Teiling Pl 10 fig.3 Forster 1969 Taf.41 fig.4

> The plants' dimensions agree well with Forster s' (1969). Lcp 51-77 Lsp 40 Wcp 79-96 Ist. 10-14

Thomasson (1974) provided photos of <u>St.manfeldtii var</u>. <u>fluminense</u> Schumacher some of which (figs. 5:1,3) are similar to the Mulwala plants, others are not (figs. 1:9-15). His plants have 59

granules on the semicell wall while, due to the presence of chloroplasts, it was hard to see if the Mulwala plants had such granules. Forster (1969) did not depict any granules. Thomasson (1974) stated his plants may be related to <u>St.pseudosebaldi var</u>. <u>planctonicum</u> as depicted in Forster (1969 Taf.41 figs.4-6).

Playfair (1908) recorded this variety from Australia. Samples: Lake Mulwala 4/4, 680921/1. Geographical distribution: Australia,

Brazil (forster 1969),

Canada (Bourrelly 1966, forma).

Staurastrum sebaldi Reins.var. ornatum Nordst. Pl 10 figs. 2,4

The presence of apical verrucae in a trigular shape as seen from the end view indicates the plants are <u>St.sebaldi</u> Reinsch. <u>var. ornatum</u> Nordst.(Teiling 1947 fig.13). The plants (Pl 10 figs. 2,4) resembles <u>St.sebaldi var. ornatum</u> (Lind & Croasdales 1966 Pl 2 fig. 15, Pl 1 fig. 1 respectively). Lind & Croasdale (1966) who gave an account of the variation in <u>St.sebaldi var. ornatum</u> stated it was a variable desmid. The front and end view of <u>St.sebaldi var.</u> <u>ornatum</u> (Frere-Irenee-Marie 1939 Pl 54 fig. 7) did not look like the Mulwala plants. Rows of granules at the base of each process as seen in Pl 10 fig. 4 were also in Lind & Croasdale s'plant (1966 Pl 1 fig. 9). <u>St.sebaldi var. ornatum</u> = <u>St.manfeldtii</u> in West & Carter (1923) according to Teiling (1947).

> Lsp 49 Wcp 66-94 Ist. 13

Previous Australian records: Hardy 1906, Scott & Prescott-1958.

Sample: Lake Mulwala 680921/1.

Geographical distribution: widespread (Lind & Croasdale 1966).

60

Staurastrum subavicula West Pl 9 fig. 11

The closest plant that looks like the Hume plant Pl 9 fig. 11 is <u>St.subavicula</u> as depicted by Lind (1967 Pl 8 fig. 5 p. 380).

Lssp 28, Wcp 47, Ist. 14.

These dimensions are comparable with those given by Lind (1967). The Hume plant, seen only once, may be this species. Sample: Lake Hume 7/3(3).

Geographical distribution: Africa (Lind 1967)

Central Norway (Bergan 1946). Staurastrum subgemmulatum West & West Pl 9 fig. 18 Claassen 1961 Taf. 31 figs. 1,2 p. 602

The plant is not <u>St.sagittarium</u> Nordst. because <u>St.sagittarium</u> have their processes curve downwards and have much shorter cell body.(Nordstedt 1888 Pl 4 fig. 6 p. 37, Thomasson & Tyler 1971 figs. 11 a,b p. 307). The plant s' dimensions agree with those given by Claassen (1961).

L 33, Wcp 50, Ist. 10.

There is no previous Australian record of the species. Samples: Lake Hume 2/4(Dam), 2/5(4), 3/5(Dam), 8/3(Tallan.).

Lake Mulwala 10/2.

Geographical distribution: Transvaal Province, Africa(?) (Claassen 1961). Staurastrum tetracerum Ralfs Pl 9 figs. 15,16

West & Carter 1923 Pl 149 fig. 2

Frere-Irenee-Marie 1939 Pl 49 fig. 19

Gronblad & Scott 1955 figs. 7,8,24 as <u>St.bibrachiatum</u> Reinsch Type II Gronblad 1960 Pl VIII fig. 174

Lsp 8, Lcp 23, Wcp 31, Ist. 4.

The dimensions are comparable with the above authors'.

Gronblad & Scott (1955) stated <u>St.tetracerum</u> may be a form of the variable <u>St.bibrachiatum</u> Reinsch.

Previous Australian records: Borge 1896, Scott & Prescott 1958, Thomasson 1973. Maskell 1881 recorded <u>St.tetracerum</u> Kutzing (not Ralfs) from New Zealand.

Samples: Lake Hume 8/3(Tallan.)

Lake Mulwala 4/4, 680921/1.

Geographical distribution: widespread (West & Carter 1923). <u>Staurastrum tetracerum var. evolutum</u> West & West Pl 9 figs. 19-21 Thomasson & Tyler 1970 fig. 8g p. 309-310

This species has been discussed and recorded from Australia (Thomasson & Tyler 1971).

Lsp 22, Wcp 32, Ist. 4.

Samples: Lake Hume 14/2(Dam), 1/5(3), 2/5(Tallan.), 2/5(4), 30/5(3),

8/7(5).

Staurastrum tohopekaligense Wolle Pl 10 fig. 9

West & Carter 1923 Pl CLV fig. 12 p. 178

The dimensions agree well with those given by West & Carter (1923).

Lcp 66-74 Lsp 31-33 Wsp 20-24 Wcp 65-72 Ist. 14-17

Processes length 24

Previous Australian and New Zealand records: Scott & Prescott (1958), Thomasson (1972, 1973), Flint (1966). Samples: Lake Hume 14/2(Dam), 7/3(3).

Lake Mulwala 680411/1, 680508/1, 690206/1.

Geographical distribution: widespread (West & Carter 1923).

Staurastrum victoriense West Pl 10 figs. 5,6 West (1909) Pl 5 figs. 13,14 p.67 Thomasson & Tyler (1971) fig. 10d

> Lsp 53 Wsp 43 Wcp 89-103 Ist. 36 Width of processes 5-7

The cell bodies are slighty bigger than West s'plant (1909) but are of the same size as Borge s'<u>St.nudibrachiatum</u> (Borge 1903 Taf.4 fig.20) therefore the Australian plants only differ from Borge s'Brazilian <u>St.nudibrachiatum</u> only in the narrower processeswith entire apices, and in the greater number of processes on each semicell (West 1909).

I agree with Thomasson & Tyler (1971) that <u>St.nudibrachiatum</u> as depicted by Gronblad (1945 Pl 11 fig. 239) s'and West s' (1909) plants are too similar to be described as different species. I doubt if West s'plants (1909) did differ significantly from Borge s'(1903). The number of terminal spines and that of the processes have, in natural populations, been shown to vary. For example: the variation in the number of processes of <u>St.sagittarium</u> Nordst.in Arthur s' lake, Tasmania (Thomasson & Tyler 1971).

Previous Australian records: West (1909), Thomasson & Tyler 1971.

Samples: Lake Mulwala 680508/1, 690206/1. Geographical distribution: Australia.

> <u>St.victoriense</u> may be closely related to <u>St.nudibrachiatum</u> Borge which had been recorded from Brazil (Borge 1903, Gronblad 1945).

<u>Staurodesmus cuspidatus</u> (Breb.) Ralfs <u>subsp. tricuspidatus</u> (Breb.) Pl 7 fig. 13

= Staurastrum cuspidatum(Frere-Irenee-Marie 1939 Pl 55 fig. 1 p. 280)

The Hume plants have shorter spines than Teiling s'plants. (Teiling 1948 fig. 61).

Lcp 27, Wcp 29, Ist. 5, spines length 5

Teiling (1948) stated <u>Arthrodesmus constrictus</u> G.M.Smith belong to <u>Staurodesmus cuspidatus</u>.

Previous Australian and New Zealand records: Hardy (1906), Borge 1896, West 1909, Scott & Prescott (1958), Maskell (1883), Nordstedt (1888), (all the authors recorded <u>Staurastrum cuspidatum</u>). Thomasson (1973) recorded <u>Staurodesmus cuspidatus</u> from Sydney. Samples: Lake Hume 14/2(Dam), 5/4(Dam), 2/5(4), 3/6(5), 10/9. Geographical distribution: widespread (West & Carter 1923 as

Staurastrum cuspidatum).

Staurodesmus triangularis (Lagerh.) Teiling 1948 Pl 7 figs. 16-18 Teiling 1948 fig. 64, 1967 p. 517

> Lssp 16-20 Wssp 12-16 Ith. 3-4 spines length 19-24

Teiling (1948) stated this species is synonymous with <u>Arthrodesmus quiriferus</u> West & West and <u>Arthrodesmus triangularis</u> Lagerheim. Two features of <u>Staurodesmus triangularis</u> described by Teiling (1967) are observed in the Hume and Mulwala plants, these features are: the apex somewhat elevated above the bases of the spines, often retuse in the median part, isthmus mostly elongate.

The degree of divergency of the spines and their length were found to vary.

Previous Australian record: Thomasson (1973).

64

Samples: Lake Hume 3/5(Dam), 3/6(5), 11/9(5).

Lake Mulwala 4/4, 7/5, 4/6. Geographical distribution: seem widespread. Division Chrysophyta

Class Chrysophyceae

Dinobryon divergens Imholf Pl 12 fig. 8 Huber-Pestalozzi 1941 Abb. 289

The basal part of the species'loria is wavy in a characteristic manner. It occurs in oligotrophic waters(Huber-Pestalozzi 1941).

Loria L 35-47

W 7-9

The dimensions agree well with those given by the above author. The variation in the loria shape is observed in both lakes and is given in detail by Ahlstrom (1937 Pl 1 figs. 1-36).

Previous Australian records: Thomasson (1973).

Flint (1966) recorded this plant from New Zealand. Samples: Lake Hume 2/5(4), 3/5(Dam), 5/7(2), 8/7(5), 10/7(3), 7/8(3).

Lake Mulwala 7/5, 680508/1, 680921/1, 681121/2.

Dinobryon sertularia Ehrenb. Pl 12 fig. 9

Huber-Pestalozzi 1941 Abb. 290

The loria dimensions agree well with Huber-Pestalozzi s'

L 29-41

W 9-11

There is loria shape variation in Lakes Hume and Mulwala. Huber-Pestalozzi (1941) and Ahlstrom (1937 Pl 2 figs. 1-16) provided detail of loria shape variations. The outline of <u>Dinobryon sertularia</u> may be slightly undulated.

> Previous Australian records: Playfair (1912, 1915, 1921), Thomasson (1973).

According to Playfair (1912) Wesenberg-Lund & Kofoid regarded species of <u>Dinobryon</u> as forms of one biological species. No polytypical colonies of many species'loria were observed in the two lakes. I agree with Playfair (1912) that wherever colonies of different species lorea are observed the many species belong to the same biological species.

Samples: Lake Hume 3/5(Dam), 5/7(2), 8/7(5), 10/7(3), 8/8(4), 10/9(4). <u>Mallomonas akrokomos</u> Ruttn. Pl 12 fig. 5 Huber-Pestalozzi 1941 Abb. 109 a p. 93

L 30, W 5

The spiral flagellum shown in Huber-Pestalozzi s'drawing may have been lost. Harris (1958) who gave the description of <u>Mallomonas akrokomos</u> as seen with the optical and the electron microscopes stated that the tail may be straight or bent as seen in the Mulwala plant. Only one plant was seen.

Previous Australian record: Croome & Tyler (1973). Sample: Lake Mulwala 4/4.

Mallomonas splendens (G.S.West) Playfair Pl 12 fig. 6

Synonym: Lagerheimia splendens G.S.West

West 1909 Pl 6 figs. 4-8 p. 74

Playfair 1921 Pl 11 fig. 3 p. 108

The mentioned drawings show three setae while the Hume & Mulwala plants have four. Playfair (1921) stated there may be 2-4 setae before and behind.

L 24-41

W 9-10

Setae length 22-31

The spiral striation on the cell wall was observed clearly on one empty cell (Pl 12 fig. 6).

66

Previous Australian records: West (1909 as <u>Lagerheimia</u> splendens), Playfair (1912,1921), Thomasson (1973). Samples: Lake Hume 1/5(3), 2/5(4), 3/5(Dam), 10/5(3), 10/9(4),11/9(5).

Lake Mulwala 4/4, 7/5, 4/6. Mallomonas sp. Pl 12 figs. 10-12

Without the knowledge of the scale shape and morphology which are important taxonomic criteria (Harris 1960, 1970) I cannot positively identify the <u>Mallomonas sp</u>.(Pl 12 figs. 10-12). The lack of rear bristles suggests they may be a form of <u>Mallomonas tonsurata</u> Teiling (Huber-Pestalozzi 1941 Abb. 108, Harris 1970 fig. 1d) but the plants have only one instead of two chromatophores as shown in Huber-Pestalozzi s'drawing. The spiral flagellum as depicted in Huber-Pestalozzi s'drawing may have been lost.

Harris (1970) pointed out that the imperfect form of <u>Mallomonas intermedia</u> Kisselew (fig. 1e) resembles the perfect form of <u>Mallomonas tonsurata</u> Teiling (fig. 1d). There is a possibility that the Hume and Mulwala plants seen may be the imperfect form (i.e. the form that lack the features that have been included in the diagnosis-Harris 1970) of Mallomonas intermedia.

L 23-29

W 11-12

Samples: Lake Hume 5/4(Dam), 2/5(4), 3/5(Dam), 3/6(5), 11/9(5).

Lake Mulwala 4/6.

Synura spinosa Korch Pl 12 fig. 7

The plants were identified as <u>Synura spinosa</u> Korch because their scales were observed under the oil immersion to have thickened apical ends as shown in Huber-Pestalozzi (1941 Abb. 195 p. 139). Their scales'dimensions are comparable with those given by Huber-Pestalozzi (1941). L 4, W 3, Tail length 4. No flagella were observed. To confirm the plants are <u>Synura spinosa</u> electron micrographs of the scales and the whole organism as given in Petersen & Hansen (1956) are needed.

Petersen & Hansen (1956) stated the shape of the cell is of no taxonomic valve. Within the same species, there are few-celled colonies with nearly sphaerical cells and large, many-celled colonies with long-stalked cells. These two cell shape and colony were observed in lakes Hume and Mulwala.

There is no previous Australian record of the species. Samples: Lake Hume 30/5(3), 5/7(2), 8/7(5), 9/7(3, abundant), 10/7(3), 7/8(3), 8/8(4), 681122/3.

Lake Mulwala 4/4, 681122/3 (mostly long-stalked cells). Class Bacillariophyceae

Centrales

Attheya Zachariasi J. Brun. Pl 13 figs. 1-6

Huber-Pestalozzi 1942 Abb. 518

There are two forms of Attheya Zachariasi in the two lakes. Form 1 (Pl 13 fig. 6) is relatively wider and shorter than Form 2 (Pl 13 fig. 1). More of the slender form 2 are present.

	Form 1	Form 2
Lssp	2128	25-75
Lcsp	52-95	80-137
W	18-33	14-18

Intermediates of the two forms were present. Pl 13 figs. 2-5 show the vegetative cell division of <u>Attheya Zachariasi</u>. The frustules and chromatophores elongate, followed by the formation of two sets of daughter cells'spines. The aaughter cells soon break apart.

Previous Australian record: Playfair (1912).

68
Samples: Lake Hume 2/4(Tallan.), 5/4(Dam), 8/3(Tallan.), 1/5(3),

2/5(4), 3/5(Dam), 31/5(4), 3/6(5), 8/7(5), 9/7(3), 10/7(3), 11/9(5).

Lake Mulwala 4/4, 4/6, 690206/1.

Cyclostella stelligera Cleve & Grun. Pl 13 fig. 7

Huber-Pestalozzi 1942 Abb, 484c p. 397

The diameter 16 µis within the range given by Huber-Pestalozzi (1942). Only one plant was seen.

Previous Australian records: Playfair 1912 (Pl LVI fig.26) Croome & Tyler 1972, Thomasson 1973.

Sample: 3/5(Dam).

<u>Melosira granulata</u> Ralfs Pl 13 figs. 8 a,b Huber-Pestalozzi 1942 Abb. 451,452

There are two major forms of <u>Melosira granulata</u> in the two lakes. Intermediate forms are present. The first form has wider but shorter frustules. (Pl 13 fig. 8a).

L 18-25

W 8-17

The spines of end frustules were observed to vary in length and in number from none to four.

The second form is much slender and each end frustule has one to two spines which also vary in length.

- L 22-31
- W 5-8

Previous Australian and New Zealand records: West (1909), Playfair (1912), Thomasson (1973), and Flint (1966).

Samples: All samples from both lakes.

Melosira varians Agardh, Pl 13 fig. 9

Huber-Pestalozzi 1942 Abb. 447

The plants may be wider than Huber-Pestalozzi s'plants whose dimensions are in brackets.

L 22-32 (8-35) W 11-23 (9-13)

Previous Australian record: Playfair (1914 p. 123). Samples: Lake Hume 1/5(3), 3/5(Dam), 30/5(3), 31/5(4), 5/7(2),

8/7(5), 10/7(3), 7/8(3), 8/8(4).

Lake Mulwala 4/4, 7/5, 4/6, 12/8, 680921/1, 681121/2.

Rhizosolenia eriensis H.L.Smith Pl 13 fig. 13

Huber-Pestalozzi 1942 Abb. 514

Lssp 50-54 Lcsp 97-103 W 3-5

Previous Australian and New Zealand records: Thomasson (1973), and Flint (1966).

Samples: Lake Hume 2/5(4), 11/9(5).

Lake Mulwala 4/4, 4/6.

<u>Rhizosolenia eriensis</u> H.L.Smith <u>var. morsa</u> G.S.West Pl 13 figs. 10-12 Huber-Pestalozzi 1942 Abb 515

> Lssp 24-49 Lcsp 69-107 W 4-15

Previous Australian record: Playfair (1912).

Samples: Lake Hume 2/4(Tallan.), 3/5(Dam), 11/9(5),

Lake Mulwala 4/4, 4/6

Pennales

Asterionella formosa Hass. Pl 14 fig. 12

Huber-Pestalozzi 1942 Abb. 531a.

Patrick & Reimer 1966 Pl 9 figs. 1,3 p. 159.

l 28-57

Previous Australian and New Zealand records: Thomasson (1973), and Flint (1966), Wood (1960).

Samples: Lake Hume 2/5(4), 9/7(3), 10/7(3), 9/8(5), 11/9(5).

Lake Mulwala 12/8.

Cymbella cf. gastroides Kutz., Cymbella cf. gastroides var minor Pl 14 fig. 16

Heurck (1896) Pl 1 figs. 35,36 p. 146

No drawings were found to match the plants better than <u>Cymbella stroides</u> Kutz. given by Heurck (1896). Their dimensions are within the range of the type form and <u>var. minor</u> given by Heurck (1896).

L 69-160

Striation in $10 \mu = 7-10$

Cymbella gastroides is common in fresh waters (Heurck 1896). There is no previous Australian record. Samples: Lake Hume 5/4(Dam), 5/7(2), 8/7(5), 7/8(3), 8/8(4).

Lake Mulwala 4/4.

Cymbella lanceolata Ehr. Pl 14 fig. 17

Heurck 1896 Pl 1 fig. 37

The plants' dimensions are comparable with those given by the above author.

l 116

Striation in 10 μ = 8

Previous Australian record: Wood (1960). This species is very common in fresh waters (Heurck 1896).

Sample: Lake Hume 7/8(3).

<u>Cymbella gastroides</u> Kutz. and <u>Cymbella lanceolata</u> Ehr. may be forms of the same plant. Both have beaded striation, their dimensions and the number of striation in 10 pare comparable. Their valves'view differs slightly (pl 14 figs. 16,17).

Epithemia sorex Kutz. Pl 14 fig. 5

Judging from the morphology and dimensions given by Heurck (1896 Pl 9 fig. 351 p. 295) the Hume and Mulwala plants (Pl 14 fig. 5) may be Epithemia sorex Kutz.

L 44

Striation in 10 μ = 15

Bourrelly (1968) gave photos of <u>Epithemia sorex</u> which showed the beaded striation very clearly (Pl 100 figs. 10-12). <u>~ NZ</u> Previous Australian records:West (1909), Wood (1960). Samples: Lake Hume 5/4(Dam), 5/7(2), 10/7(3).

Lake Mulwala 4/4, 7/5, 4/6, 12/8, 680508/1, 680921/1, 681121/2.

Fragilaria capucina Desmazieres Pl 14 fig. 6

The closest plant that looks nearly like the Hume and Mulwala is <u>Fragilaria capucina</u> Desm. (Huber-Pestalozzi 1942 Abb. 527a p. 441). Bourrelly (1968) figured <u>Fragilaria capucina</u> Desm. (Pl 63 fig. 9) which seemed to have wider central area than the plants. He gave no dimensions. The plants'dimensions are comparable with those given by Huber-Pestalozzi (1942).

L 50

Striation in 10 $\mu = 13$

Previous Australian record: Wood (1960). Samples: Lake Hume 10/7(3)

Lake Mulwala 12/8.

Fragilaria sp. Pl 14 fig. 7

I could find any drawings to match the plants. They seem to be <u>Fragilaria sp</u>.which differs from <u>Fragilaria capucina</u> (Pl 14 fig. 6) in being longer and having coarser striation.

l 169

Striation in 10 $\mu = 8$

Samples: Lake Hume 5/7(2), 8/7(5).

Gomphonema constrictum Ehrenb. Pl 14 fig. 14

Heurck 1896 Pl 7 fig. 296 p. 270

Bourrelly 1968 Pl 96 figs. 1,2

The plants' dimensions are within the range given by Heurck (1896). The costae on each striae were not observed.

L 36-46

Striation in 10 μ = 11-12

Common in fresh waters (Heurck 1896). West (1909) recorded this species from Australia. Wood (1960) recorded it from NZ. Samples: Lake Hume 9/8(5).

Lake Mulwala 4/4, 4/6, 680411/1, 680921/1, 690206/1. <u>Gomphonema constrictum var. capitatum</u> Ehr. Pl 14 fig. 15 Heurck 1896 Pl 7 fig. 297 p. 270. Wood (1960) Pl 54 figs. 161 a.b.

The dimensions are within the range given by Heurck (1896).

L 43-55

W 13-14

Striation in 10 $\mu = 11-13$

Hirano (1964) recorded <u>Gomphonema constrictum</u> Cleve (not Ehrenb.) <u>var. capitata</u> (Ehrenb.) Cleve (Pl 10 fig. 14) with stipulated striation. The Hume and Mulwala plants seem to have smooth striation.

This variety is common in fresh waters (Heurck 1896). Previous Australian and New Zealand records: Wood (1960); Flint (1966). Samples: Lake Hume 3/5(Dam), 8/7(5), 9/7(3), 10/7(3), 7/8(3), 10/9(4).

Lake Mulwala 7,65, 4/6, 12/8.

Cf. Gyrosigma attenuatum Ku tz. Pl 14 fig.9

<u>Gyrosigma sp.</u> have longitudinal and transverse striation on valve surface perpendicular to each other while in <u>Pleurosigma sp</u>. they are not at 90° angle to each other (Patrick & Reimer 1966). Only longitudinal striation was seen in the Hume and Mulwala plants therefore they may be <u>Gyrosigma sp</u>. The side view (Pl 14 fig. 9) resembles <u>Gyrosigma attenuatum</u> Kutz. (Bourrelly 1968 Pl 77 fig. 9).

L 98-110

₩ 11-15

Previous Australian records of the species: West (1909). Spencer (1882) recorded <u>Gyrosigma attenuatum</u> from New Zealand. Samples: Lake Hume 7/3(2), 5/4(Dam), 1/5(3), 2/5(4), 3/5(Dam), 5/7(2),

8/7(3), 9/7(3), 7/8(3), 8/8(4), 10/9(4).

Lake Mulwala 10/2, 4/4, 12/8, 4/6, 680508/1, 680921/1,

681121/2, 690206/1.

Nitzschia pelagica O Mull. Pl 13 fig. 15 Huber-Pestalozzi 1942 Abb. 571 p. 477

The dimensions slightly differ from those given by Huber-Pestalozzi (1942) which are in brackets.

L 31-41 (35-54)

W 3-4 (2)

Striation in 10 μ = 11 (18-20)

Colonies of 4-5 frustules were observed. There is no previous Australian record.

Samples: Lake Hume 3/5(Dam).

Lake Mulwala 10/2, 4/4.

Pinnularia abaujensis (Pant.) Ross var. abaujensis Pl 14 fig. 4 Patrick & Reimer 1966 Pl 58 fig. 2 p. 612

The plants' dimensions are in the range given by the above authors.

L 88-130

W 12

Striation in 10 M = 9-11

The only difference from Patrick & Reimer s'plants is the Mulwala plants have shorter striae in the mid valve area.

There is no previous Australian record of the species. Samples: Lake Mulwala 4/4, 12/8.

Surirella Engleri O Mull. Pl 14 fig. 2

The closest plant seems to be <u>Surirella Engleri fa</u>. <u>angustior</u> (Huber-Pestalozzi 1942 Abb. 594b p. 493). This fa. is narrower than the Hume and Mulwala plants. The other forms of the species have an inward curvature in the mid region of the valves. From the literature survey the Hume & Mulwala plants may be a form of <u>Surirella Engleri</u> possibly <u>fa. angustior</u> with wider valve which is possible since the other forma of the species are wider than fa. angustior.

		Mulwala	Su.Engleri fa. angustior (Hube	er-
			Pestalozzi 1942 s')	
	\mathbf{L}	242	173-290	
	W	46	3036	
	Stri	2 (بر 10 ation	2	
	Ther	e is no previous Aus	tralian record.	
Samples:	Lake	Hume 8/7(5).		

Lake Mulwala 12/8.

Surirella spinifera Hust. Pl 14 fig. 1

Huber-Pestalozzi 1942 Abb. 627

The plants are <u>Surirella spinifera</u> Hust. because of the presence of one thorn at one end of the frustules. Their dimensions are comparable with those given by Huber-Pestalozzi (1942).

L 116-197

₩ 46-67

Striation in 10 μ = 20

The drawing (Pl 14 fig. 1) has one end less pointed than Huber-Pestalozzi s'drawing (1942 Abb. 627).

There is no previous Australian record of the species. Samples: Lake Hume 1/5(3), 30/5(3), 10/7(3),

Lake Mulwala 4/4, 680411/1, 681121/2, 690206/1. Surirella tenera Greg. Pl 14 fig. 3

The plants are probably a form of <u>Surirella tenera</u> Greg. They differ from the type form (Huber-Pestalozzi 1942 Abb. 623a) in having two thorns in the mid region (raphe). One thorn seems bigger than the other. Huber-Pestalozzi (1942) figured <u>Su.tenera</u> <u>var.nervasa</u> (Abb. 623b,c) with one big mid region thorn and eleven small ones.

The plants' dimensions agree with those of the type form given by Huber-Pestalozzi (1942).

L 132

W 40

Striation in 100 μ = 19

Wood (1960) recorded <u>Su. tenera var. splendidula</u> (Pl 56 fig. 198) from Lake Dobson, Australia. No dimensions were given. The Hume and Mulwala plants differ from his plant in having one end more pointed and the straight in place of the undulate median line. A plant from Lake Sorell, Tas. (negative 710424/1:7) seems wider across the middle of the valve otherwise it resembles the Hume and Mulwala plants.

Samples: Lake Hume 2/5(4),

Lake Mulwala 4/6. <u>Synedra acus</u> Kutz. Pl 13 figs. 16, 20 Huber-Pestalozzi 1942 Abb. 536a p. 458

There is a slight variation from the dimensions given by Huber-Pestalozzi (1942, in brackets).

> L 122-246 (100-300) W 6-7 (5-6)

Striation in 10 μ =9-10 (12-14)

Pl 13 fig. 16 is probably a form of <u>Synedra acus</u> because of the lack of striation in the central area. The valve is slightly inflated in the mid region. Its dimensions are in the range of Synedra acus s'given by Huber-Pestalozzi (1942).

- L 195
- W 5

Striation in 10 M = 11

Synedra acus Kutz. var. radians (Kutz.) Pl 13 figs. 18,19

The plant may be <u>Synedra acus</u> Kutz. <u>var. radians</u> (Kutz.) Hust. (Huber-Pestalozzi 1942 Abb. 536b) because of a pair of short-striae in the central region (Pl 13 fig. 19). Huber-Pestalozzi (1942) stated this variety had the same dimensions as the type form.

- L 235
- W 5

Striation in 10 $\mu = 10$

Wood(1960,West (1909) recorded Synedra acus Kutz. from Australia.

Samples: Synedra acus and its variety were found in

Lake Hume 5/7(2), 8/7(5), 7/8(3), 10/9(4).

Lake Mulwala 4/4, 680921/1, 681121/2.

Synedra ulna (Nitz.) Ehr. var. danica (Kutz.)V.H. Pl 13 fig. 17 Patrick & Reimer 1966 Pl 7 fig. 10 p. 151

The dimensions are comparable with those given by the above authors.

L 169-220

W 7

Striation in 10 μ = 8-10

West (1909) recorded <u>Synedra ulna</u> from Yan Yean Reservoir Vic. Flint (1966) recorded this variety from New Zealand. Samples: Lake Hume 31/5(4), 5/7(2), 8/8(4), 10/9(4).

Tabellaria flocculosa (Roth) Kutz. var. flocculosa Knudson 1952 Pl 14 fig. 18

Knudson 1952 figs. 5 a-i

Patrick & Reimer 1966 Pl 1 fig. 5

L 19-23

Number of septa 4-5

Knudson (1952) should not have created a new species, <u>Tabellaria quadriseptata</u> whech is distinguished from <u>T.flocculosa</u> by having 4 septa per frustule and the valve view has the same extent of inflation in the mid and both end regions when she knew these two features of T.flocculosa were greatly variable (p.434-435). The new species, <u>T.quadriseptata</u>, were created on the basis of very variable features and no biological concept of species was taken into account. Patrick & Reimer (1966) accepted <u>T.flocculosa</u> and <u>T.quadrisepta</u> as two distinct species. Huber-Pestalozzi (1942) figured a zig-zac colony of <u>Tabellaria fenestrata</u> (Lyngb.) Kg. (Abb. 519) which is very similar to the Hume and Mulwala plants (Pl 14 fig. 18) but according to Knudson (1952) and Patrick & Reimer (1966) the colony morphology of <u>T.fenestrata</u> is a straight line.

Previous Australian and New Zealand records: West (1909), Wood (1960), Flint (1966), thomasson (1973).

Samples: Lake Hume 30/5(3), 9/7(3), 8/7(5), 7/8(3).

Lake Mulwala 680921/1.

Cf. Navicula hennedyi Pl 14 fig. 13

A photo of an Australian diatom given by Wood (1960,fig.65b) resembles the Hume plant. Wood (1960) gave no dimensions. His references were not available. The Hume plant may belong to this taxon.

L 19, W 14, Striation 10 in 10 μ .

Sample: Lake Hume 10/9(4).

Stauroneis fulmen var. capitata Pl 13 fig. 14

The Mulwala plant resembles markedly Wood s'(1960) photo of <u>Stauroneis fulmen var. capitata</u> (Pl 53 fig. 105). He gave no dimensions.

L 132

Sample: Lake Mulwala 690206/1

Division Euglenophyta

Class Euglenophyceae

Order Euglenales

Euglena cf. gracilis Kleb. Pl 2 figs. 3,4

The closest taxon for the Mulwala plants is <u>Euglena</u> <u>gracilis</u> Kleb. (Prescott 1962 Pl 85 fig. 17 p. 393, Leedale 1967 fig. 11 p. 21). The plants resemble <u>Euglena gracilis</u> in having many chloroplasts and a somewhat pointed posterior end. The plants are slightly shorter than the above authors'plants and have a more pointed anterior end.

- L 31-32
- W 11-13

There is no previous Australian record. Haughey (1968) recorded it from New Zealand. Samples: Lake Mulwala 10/2, 7/5. <u>Euglena spirogyra</u> Ehrenb. Pl 2 figs. 1,2 Prescott 1962 Pl 86 fig. 15 p. 394 Plaufair 1912 Pl IV fig. 2 p. 119-120

L 136-149

W 11-21

The dimensions are in the range given by Prescott (1962). Claassen (1961) recorded <u>Euglena spirogyra var. major</u> from South Africa (152-196 **x** 22-26).

The plant (Pl 2 fig. 1) appeared twisted. Twisting occurs when it is irretated (Leedale, Meeuse, Pringheim 1965). Straight plants were also observed. It is said (Leedale et al 1965) a system of papillae or protuberances which are believed to be exudations emitted through pores arranged along the straie of the periplast are observed only in cultures with soil though not in all of them and in habitats rich in iron. The system of protuberances was sometimes observed (Pl 1 fig. 1). They may be lacking as seen in the plant (Pl 1 fig. 2).

The colour and ornamentation of the wall vary (Leedale et al 1965). Deformities of <u>Euglena spirogyra</u> as stated by Pringsheim (Leedale et al 1965) espectially in the tail end were not seen. Playfair (1912) stated that as a rule the granulate striae run obliquely from left to right but Pringsheim (Leedale et al 1965) observed the occurrence of right and left helices in the same clone and could afford no explation. Physiologically the paramylon of <u>Euglena spirogyra</u> was established to have a β 1:3 glucan structure which is very similar to the laminarin of Phaeophyceae rather than the starch of Chlorophyceae (Leedale et al 1965).

Euglena spirogyra is widespread and has been recorded from Australia and New Zealand (Playfair 1912 and Haughey 1969). Samples : Lake Hume 2/4 (Tallan.), 2/5(4), 8/8(4), 30/5(3).

Lake Mulwala 10/2, 4/4.

Phacus tortus (Lem.) Skvortzow Pl 2 figs. 5,6

Haughey (1968) stated that most specimens of <u>Phacus</u> <u>tortus</u> when swimming actively appear curved throughout most of their length while cells at rest appear flat. The curved or twisted plant (Pl 2 fig. 6) resembles the swimming form of <u>Phacus tortus</u> given by Haughey (1968 fig. 4:1e p. 734). The flat plant (Pl 2 fig. 5) resembles the curved one in having the longitudinal striation on the periplast and numerous chloroplasts. However the flat plant does not look exactly like Haughey s'figure of <u>Phacus tortus</u> resting form (Haughey 1968 fig. 4:1a,b). The former is narrower and has a longer tail. From the dimensions given by Haughey (1968) his <u>Phacus tortus</u> may be even narrower tha the Hume plant.

Hu	me	Haughey (1968)
L with tail	112-134	80 • 108
W	4357	30-44
tail length	37-52	32-50

Jahn (1949) recorded <u>Phacus longicauda</u> (fig. A) with body flattened but not twisted and <u>Phacus tortus</u> (fig. 81 B,C) as similar to <u>Phacus longicauda</u> except that the body is twisted. Haughey (1968) suggested specimens thought to be <u>Phacus longicauda</u> (Ehr.) may belong to this species. He suggested further on the taxonomy of Phacus. There is no previous Australian record of <u>Phacus tortus</u>, Playfair (1921) recorded <u>Phacus longicauda</u>, which may be the same plant as <u>Phacus tortus</u> (resting form) according to Haughey in 1968). Haughey (1968) recorded <u>Phacus tortus</u> from New Zealand. Samples: Lake Hume 2/5(4), 5/7(2). <u>Phacus pleuronectes</u> (Muell.) Dujardin Pl 2 fig. 7 Playfair 1921 Pl V fig. 1 p. 123 Prescott 1962 Pl 88 fig. 16 p. 402

L 41

W 29-32

Previous Australian records: Mobius 1894, Playfair 1921, Bailey 1895, Scott & Prescott 1958.

Sample: Lake Hume 8/8(4).

<u>Trachelomonas armatus var. longispina</u> (Playfair) Deflandre Pl 12 fig. 24 Playfair 1915 Pl IV fig. 20

Prescott 1962 Pl 83 fig. 27 p. 411

The dimensions agree with those given by Prescott (1962). Test dia. 30

L 40

Previous Australian record: Playfair 1915.

Sample: Lake Hume 2/5(4).

Trachelomonas bacillifera Playfair Pl 12 figs. 16-19

Playfair 1915 Pl 3 figs. 15,16

The plants' dimensions are in the range between those of the type form and those of the var. minima given by Playfair (1915).

Lssp 25-35

Wssp 22-30

Upon focusing up the spines on the wall (Pl 12 fig. 19) were seen as granular structures (Pl 12 figs. 16,17). The flagella pore could be seen when the plant was tilted. (Pl 12 fig. 17). In 1921 Playfair figured <u>Trachelomonas bacillifera</u> with longer and blunt spines (Pl VII figs. 2,3) and stated they were reddish yellow. The Hume and Mulwala plants were brick red.

Previous Australian records: Playfair (1915, 1921). Samples: Lake Hume 8/3(Tallan.), 5/4(Dam), 1/5(3), 2/5(4), 3/5(Dam),

3 0/5(3), 31/5(4), 3/6(5), 8/7(5), 9/7(3), 10/7(3).

Lake Mulwala 4/4, 7/5, 4/6.

Trachelomonas caudata Stein var, australica Playfair Pl 12 fig. 25

The closest plant that may be the plant, seen only once, is <u>Trachelomonas caudata</u> Stein <u>var. australica</u> Playfair (Playfair 1915 Pl V fig. 2). The dimensions are within the range given by Playfair (1915).

> L 87 W 18 Collar L 19 W 5 Cauda L 28 W 5

Playfair (1915) recorded this species from NSW. Sample: Lake Hume 7/8(3).

Trachelomonas hispida var. coronata Lemmermann Pl 12 fig. 22 Prescott 1955 Pl 1 fig. 23

1962 Pl 83 fig. 30

The Hume plants are slightly smaller than Prescott s'.

Test dia 19

L 24

Sample: Lake Hume 2/5(4).

Trachelomonas hispida (Perty) Stein var. rectangularis Br. Schroder Pl 12 figs. 20,21

The closest plant that resembles the Hume plants is <u>Trachelomonas hispida var. rectangularis</u> (Playfair 1915 Pl III fig. 10 p. 21). Their dimensions are comparable.

Lssp 30-41

Wssp 20-21

The colour was dark orangy red.

Samples: Lake Hume 2/5(4), 5/7(2), 7/8(3).

Trachelomonas scabra Playfair Pl 12 figs. 13-15

The curved neck of Pl 12 fig. 15 suggests <u>Trachelomonas</u> <u>euchlora</u> (Ehr.) Lemm. which may or may not be smooth-walled (Playfair 1915 Text figure 4 p. 16 and Pl 3 fig. 1). The Mulwala plant is slightly shorter than Playfair s'<u>Tr.euchlora</u>.

Ls collar	22-25
W	20-21
Collar L	4
W	5

Since Prescott (1962) named a plant with a curved neck <u>Tr.scabra var. longicallis</u> Playf. (Pl 1 fig. 8) all the similar plants observed with with square but rounded corners cells and a collar will be grouped in <u>Tr.scabra</u> Playf. no matter if their necks are curved. It is not possible to observed the wall ornamentation (if any) because the cell contents were present. Playfair s'(1915) Tr.euchlora and Tr.scabra seem closely related.

> ∾ بر مونور ۲۰۰۰ ور مراجع این مورده و

Previous Australian record: Playfair (1915). Sample: Lake Mulwala 7/5. Trachelomonas superba Swirenko Pl 12 fig. 23 Messikommer 1927 Taf. VI fig. 7

> Lcsp 36-42 Wcsp 27-30

Morphologically this species was observed to differ from <u>Tr. bacillifera</u> (Pl 12 figs. 16-19) in not having the "granules" or pore-like structures upon focusing up. Prescott (1962) showed the species had spines only at the anterior and posterior ends (pl84 fig. 10). It was found that the plants seemed to possess spines at the anterior and the posterior ends when they were viewed in one plane only. Upon focusing they were seen to have spines all over the tests.

Samples: Lake Hume 5/4(Dam), 2/5(4), 3/5(Dam), 3/6(5), 5/7(2), 8/7(5). Lake Mulwala 4/6.

Trachelomonas volvocina Ehr. var. punctata Pl 12 fig. 26

The Hume and Mulwala plants were identified as <u>Trachelomonas volvocina var. punctata</u> because its shape, punctate wall and colour agree with Playfair s'description although they were bigger. The colour was dark red.

Test dia. 19 (Playfair s'plants test dia. 13-14)

Prescott (1962) recorded <u>Tr. volvocina var. punctata</u> Playf. of even smaller tests (Pl 83 fig. 12 p. 419, test dia. 10, L 20). therefore the dimensions vary.

Previous Australian record of the variety: Playfair (1915). Schmidle (1896) and Haughey (1968) recorded the tpye form with no wall punctation from Australia and New Zealand. Samples: Lake Hume 8/3(Tallan.), 5/4(Dam), 1/5(3), 2/5(4), 3/5(Dam),

30/5(3), 31/5(4), 3/6(5), 5/7(2), 8/7(5), 9/7(3), 10/7(3), 11/9(5).

Lake Mulwala 10/2, 4/4, 7/5, 12/8.

Division Pyrrhophyta

Class Dinophyceae

Order Peridiniales

<u>Ceratium hirundinella</u> (O.Muell.) Dujardin Pl 12 fig. 2 Prescott 1962 Pl 92 fig. 5 p. 437

L 253-266

The length of the only one posterior horn was observed to vary. The thecal structure of <u>Ceratium hirundinella</u> as viewed under the scanning electron microscope was given by Dodge et al (1970).

Previous Australian and New Zealand records: Mobius (1892), Playfair (1912), Thomasson (1973) and Flint (1966).

Samples: Lake Hume 14/2(Dam), 7/3(3), 8/3(Tallan.), 5/4(Dam), 1/5(3),

2/5(4), 3/5(Dam), 31/5(4), 3/6(5).

Lake Mulwala 4/6.

Peridinium sp. Pl 12 fig. 4

The pattern of the wall plates is needed in identifying Peridinium sp.

Samples: Lake Hume 8/3(Tallan.), 2/5(4), 30/5(3), 7/3(3).

Glenodinium sp. Pl 12 fig. 3

The characteristics of the genus given in Prescott (1962) were observed in the Hume plants : cell wall thin, plates obscure with a transverse furrow that extends completely around the cell. Sample: Lake Hume 8/3(Tallan.).

Division Cyanophyta

Class Myxophyceae

Order Chroococcales

Coelosphaerium Naegelianum Unger. Pl 3 figs. 12,13

The morphology and size of the plants suggest <u>Coelosphaerium</u> <u>Naegelianum</u> Unger. (Smith 1920 Pl 4 fig. 1 p. 35, Desikachary 1959 Pl 28 figs. 9, 16 p. 147) but no mucilagenous envelope was observed. Their dimensions agree well with those given by Desikachary (1959).

cell L 2-3

W 4-6

There is no previous Australian record.

Samples: Lake Hume 5/4(Dam), 1/5(3), 2/5(4), 3/6(5), 8/7(5), 10/7(3),

9/8(5), 11/9(5), 681122/3.

Lake Mulwala 4/4, 7/5, 4/6, 12/8.

Microcystis aeruginosa Kutzing Pl 1 figs. 6, 11

Smith 1920 Pl 5 figs. 1,2 p. 39

Prescott 1962 Pl 102 fig. 1 p. 456

cell dia. 3-4

There is no previous Australian record. Haughey (1968) recorded it from New Zealand.

This species is a frequent component of water blooms, especially in lakes with eutrophic characteristics (Prescott 1962). Samples: Lake Hume 14/2(Dam), 1/5(3), 7/3(3), 2/5(4), 3/5(Dam), 30/5(3),

3/6(5), 10/7(3), 7/7(3), 9/8(5), 11/9(5).

Lake Mulwala 10/2, 7/5, 4/6.

Microcystis flos-aquae (Wittr.) Kirchner. Pl 1 fig. 7

Smith 1920 Pl 5 fig. 3

Desikachary 1959 Pl 17 fig. 11 p. 94

The Mulwala plants'cell dia. is comparable to that given by Desikachary (1959). Cell dia. 7-8. Some workers put <u>Microcystis flos-aquae</u> (Wittr) Kirchner. as synonymous with <u>Microcystis aeruginosa</u> Kutzing (See Prescott 1962 for further discussion).

Previous Australian records: Scott & Prescott 1958, Thomasson 1973.

Sample: Lake Mulwala 4/6.

Anabaena limnetica G.M.Smith Pl 1 fig. 4

Using the key provided by Smith (1920) this slightly flexed <u>Anabaena</u>, seen only once, may be <u>Anabaena limnetica</u> G.M.Smith (Smith 1920 Pl 8 fig. 8 p. 57). The Hume dimensions agree with those given by Smith (1920).

cells	W	9-13
hetercyst	W	9
spore	W	18
	T.	31

This species differs from <u>Anabaena spiroides</u> Kleb. in that the filament did not coil and the vegetative cells and the akinetes are bigger. Desikachary did not describe this species from India (1959).

There is no Australian record of the species. Sample: Lake Hume 3/5(Dam).

Anabaena spiroides Klebahn Pl 1 figs. 1-3

Desikachary 1959 Pl 71 fig. 9 p. 395

cells	W	68
heterocysts	W	8–10
akinetes	W	10

L 15-23

The dimensions are comparable with those given by Desikachary (1959). Sometimes the heterocysts are in the mucilagenous envelope (Pl 1 fig. 3). There is no previous Australian record. Flint (1966) recorded <u>Anabaena spiroides</u> Kleb. from New Zealand. Samples: Lake Hume 14/2(Dam), 7/3(3), 8/3(Tallan.), 1/5(3), 2/5(4),

3/5(Dam), 5/7(2), 11/9(5).

Lake Mulwala 10/2, 4/4.

Lyngbya palmarum Bruhl et Biswas Pl 1 fig. 8

Desikachary 1959 Pl 55 fig. 6 p. 301

The Mulwala plant is <u>Lyngbya palmarum</u> Bruhl et Biswas (using Desikachary 1959 s'key p. 198-199). The sheath is yellow and the plant is slightly shorter than the Indian plant.

cell L 2

W 7

Prescott (1962) recorded a similar plant Lyngbya aerugineo-caerulea (Kutz.) Gomont (Pl 111 fig. 11 p. 498) but his plant had colourless sheath.

There is no previous Australian record of the species. Samples: Lake Mulwala 10/2, 7/5.

Oscillatoria curviceps Agardh. ex Gomont Pl 1 fig. 14

Using Desikachary s'key (1959 p. 198-199) the plant was identified as <u>Os.curviceps</u> Agardh. ex Gomont.(Pl 39 figs. 9,10) The Mulwala plant differs from Desikachary s'plant in having slightly bent apex. The dimensions agree well.

cell L 3

W 14

The cross walls of <u>Os.curviceps</u> may or may not be granulated (Prescott 1962 Pl 108 fig. 17 p. 487).

There is no previous Australian record. Sample: Lake Mulwala 10/2. Oscillatoria lacustris (Kleb.) Geitler Pl 1 fig. 10 Prescott 1962 Pl 109 fig. 15 p. 488 Bourrelly 1966 Pl 1 fig. 8

cell dia 5-6

ь 3-5

Prescott (1962) noted it was possible that subsequent study will establish <u>Os.lacustris</u> (Kleb) Geitler as a new fresh water species of <u>Trichodesmium</u>. According to Desikachary (1959)<u>Os.</u> <u>lacustris</u> (Kleb) geitler is synonymous with the fresh water Trichodesmium lacustre Kleb. (Pl 42 fig. 22 p. 246).

Sample: Lake Mulwala: 7/5.

Oscillatoria nigra Vaucher Pl 1 fig. 5

Using Desikachary s'key (1959 p. 198-199) the plant is identified as <u>Os.nigra</u> Vaucher.

cell L 2-4

. .

W 7

The plant has no granulated cross wall. Prescott (1962) stated the cross wall of <u>Os.nigra</u> may or may not be granulated while Desikachary (1959) stated it is. The drawings given by Prescott (1962, Pl 109 fig. 18) showed the end cells had rounded ends. The plant end cells have truncate ends like those of Desikachary (1959 p. 223).

This species has no previous Australian record. Spencer (1882) and Norstedst (1888) recorded it from New Zealand. Sample: Lake Mulwala 10/2.

Oscillatoria subbrevis (cf.) Schmidle Pl 1 fig. 13

The plant resembles closest <u>Os.subbrevis</u> Schmidle (Desikachary 1959 Pl 40 fig. 1 p. 207). The dimensions are in the same order as Desikachary s'. Cell L 3, W 7. Away from the apical cell of <u>Oscillatoria subbrevis</u> seen the cells seem to be divided into two by a cavity filled with granules.

Sample: Lake Mulwala 680921/1.

Oscillatoria waterbergensis Claassen 1961 Pl 1 fig. 12

The morphology and dimensions agree with those of the new species created by Claassen in 1961 (Pl 2 fig. 9 p. 565).

cell L 4-6

W 5

There is no previous Australian record. Sample: Lake Mulwala 680921/1.

Schizothrix rivularis (Wolle) Drouet Pl 1 fig. 9

The closest plant that looks like the Hume plant (pl 1 fig. 9) is Schizothrix rivularis (Wolle) Droret (Prescott 1962 Pl 114 fig. 3). The filament observed had fluorescent property. Prescott (1962) recorded the cells are quadrate 5-11 μ in diameter.

Cell dia. 5

L 7-12

There is no previous Australian record. Prescott (1962) stated this species, left standing, would dissociate from the bundle sheath and give out pigments to fluoresce the water. Samples: Lake Hume 3/5(Dam), 8/3(Tallan.).

Protozoa

Vorticella sp. Pl 12 fig. 1

The contracted stalk resembles that of <u>Vorticella sp</u>. (Jahn 1949 fig. 358 b). Jahn (1949) stated <u>Vorticella</u> s'stalk possessed internal myoneme which was capable of contracting. The sheath merely coiled like a spring.

Samples: Lake Hume 3/5(Dam), 11/9(5), 8/7(5).

More diatoms

Gomphonema Augur Ehr. var. Guatieri H.V.H. Pl 14 fig. 11

The Mulwala plants look identical with <u>Gomphonema</u> <u>Augur Ehr, var. Guatieri</u> H.V.H. as depicted by Heurck (1896 Pl VII fig. 302 p. 271). Their dimensions agree with those given by Heurck. (1896).

L 50

W 16

Striation 11 in 10 M

Samples: Lake Mulwala 10/2, 4/6.

There is no previous Australian record.

Unidentified diatoms

1) Pl 14 fig. 10 (L 58, W 12, Striation 11 in 10 μ). Samples: Lake Hume 5/7(2), 7/8(3), 10/9(4).

Lake Mulwala 680921/1.

2) Pl 14 fig. 18 (L 58, W 14, Striation 25 in 10 μ). Samples: Lake Hume 1/5(3), 7/8(3), 10/9(4).

Lake Mulwala 4/6, 12/8.

DISCUSSION

Assuming that desmids are inhabitants of oligotrophic (nutrient-poor)waters and Euglenineae, Myxophyceae, and a great majority of Chloroccoccales and Centrales have an eutrophic tendency, Nygaard (1949) proposed the phytoplankton hypothesis whereby the numbers of oligotrophic and eutrophic species found in plankton samples are used to calculate the compound phytoplanktonquotient which is defined as:

<u>Myxophyceae + Chlorococcales + Centrales + Euglenineae</u> Desmidieae

The quotient of less than 1.0 indicates oligotrophic lakes. Values of greater than 1.0 probably indicate eutrophic lakes (nutrient-rich) and those between 5-20 indicate a higher degree of eutrophy.

Brook (1959b, 1965) pointed out two major limitations to the use of the compound phytoplankton quotients to determine the trophic status of lakes:

1) Most of the desmids were found to be tychoplanktic, i.e. adventitous forms from other habitats, therefore the inclusion of all desmids collected into the denominator of the quotient gave a misleadingly-low value which indicated olightrophy.

2) Taxonomists have different opinions on the delimitation of species hence one investigator s'determination may not be comparable to the other s'.

Nygaard (1949)& Brook (1959b, 1965) were of the same opinion that the aim of this biological means of determining the trophic status of lakes must be the formulation of a quotient based on a limited number of planktonic species, whose status in the plankton and whose nutritional requirement have been adequately investigated. Brook (1965) stated quotients can be applied with a fair degree of reliability if the limitations mentioned above are observed. The Hume and Mulwala compound phytoplankton quotients (Table 2) indicate both lakes are in the same lower range of eutrophy. The desmids and other algae, seen in 1968-1969 samples only, were not taken into account since the corresponding 1974 samples lack them. The 1968-1969 Mulwala samples had high numbers of some desmid species, in particular, <u>Staurastrum arctiscon</u> (Ehrenb.) Lund <u>var. glabrum</u> West & West, <u>St.dickiei</u> Ralfs, <u>St.grande</u> Bulhn <u>var. parvus</u> West, <u>St.furcatum</u> (Ehrenb.) Breb., <u>St.tohopekaligense</u> Wolle, and <u>St.victoriense</u> West. This observation shows that Mulwala 1968-1969 conditions differ from those at present. Perhaps the lake was more oligotrophic. Brook (1965) stated that an increase in pH may be responsible for the disappearance of desmids.

The other approach is to characterise waters in terms of their dominant plankton species and to classify them according to the community or associations they support. The plankton of an oligotrophic lake may be characterised by a large numbers of desmids species although it may be dominated numerically by one or two species of diatoms or Chrysophyceae, and a eutrophic may be characterised not by a large number of species of Myxophyceae but by very large numbers of filaments or colonies of only one or two species as during a water bloom (Brook 1965). Using this approach the two lakes are oligotrophic (Table 2).

To average the results of the determination of the determination of the trophic status of Lakes Hume and Mulwala by the two approaches, the two lakes may be said to be mesotrophic. 94

Geographical distribution of desmids seen.

The world distribution of algae in order to characterise them according to climatic regions has only a tentative value since algae from all parts of the world have not been critically examined. Also, the delimitation of species especially in the desmids is very subjective. It depends on the judgement of the desmologists.

From the literature survey all desmids seen are cosmopolitan except:

1) <u>Micrasterias Hardyi</u> G.S.West has only been recorded from Australia.

2) <u>Cosmarium magnificum</u> Nordstedt, <u>Staurastrum asterias</u> Nygaard, and St.victoriense G.S.West have no European records.

3) The southern hemisphere <u>St.nodulosum</u> Prescott may be a form of the widespread <u>St. leptocladum</u> Nordstedt.

4) <u>St.anatoides</u> Scott & Prescott may be synonymous with <u>St.quebecense</u> Frere-Irenee-Marie in which case this alga has been recorded from both hemispheres.

EXPLANATION OF PLATES

Where there is no along-side scale, the bottom scale applies.

		PLATE 1
Fig.	1-3	Anabaena spiroides Kleb.
	4	" limnetica G.M.Smith
	5	Oscillatoria nigra Vaucher
	6	Microcystis aeruginosa Kutz.
	7	" flos-aquae (Wittr.) Kirchner.
	8	Lyngbya palmarum Bruhl et Biswas
	9	Schizothrix rivulis (Wolle) Drouet
	10	Oscillatoria lacustris (Kleb.) Geitler
	11	Microcystis aureginosa Kutz.
	12	Oscillatoria waterbergensis Claassen
	13	" Cf. subbrevis
	14	" curviceps Agard ex Gomont
		PLATE 2
	1,2	Euglena spirogyra Ehrenb.
	3,4	" cf. gracilis Kleb.
	5,6	Phacus tortus (Lem.) Skvortzow
	7	" pleuronectes (Muell.) Dujardin
	8	Eudorina elegans Ehrenb.
	9	Pledorina californica Shaw
	10,11	Pandorina morum (Muller) Bory
	12,13	Volvox cf. aureus Ehrenb.
	14	" cf. globator Linnaeus
	15,16	" cf. Carteri G.M.Smith
	17	Botryococcus Braunii Kuetz.
	18.	Micractinium pusillum Fres.

All genus and species names have not been underlined.

Fig.	1-5	Actinastrum	Hantzs	chii	Lag.			
	6	Ankistrodes	mus fal	catu	s Corda	1		
	7,8	Selenastrum	westii	G.M	.Smith			
	9	Kirchneriel	la luna.	ris	(Kirch	ner.) Moeb.		
	10,11	Dictyosphae	rium pu	lche	llum Wo	bod		
	12,13	Coelosphari	um Naeg	elia	num Ung	ger.		
	14	Dimorphocod	cus lun	atus	A.Bra	in.		
	15	Coelastrum	reticul	atum	(Dang	.) Senn.		
	16	11	micropo	rum	Naeg.			
	17	Treubaria t	riappen	dicu	latum	Bernard		
	18	Unidentifie	d plant					
	19,20	Oocystis pa	rva Wes	t &	West			·
	21-23	" la	custris	Cho	dat			
	24-26	Nephrocytiu	ım Agard	hian	um Nae	geli		
		PLA	TE 4					
	1,2	Pediastrum	duplex	Meye	n var.	clathratum (A.Brun.)	Lag.
	3		11	11	var.	gracillimum	West & W	est
	4	11	FT .	**	type	form		
	5,6	11	17	11				
	7	11	14	11	var.	rugulosum Ra	ciborski	
	8,9	Pediastrum	tetras	(Ehr	enb.)	Ralfs		
	10-12	Spherocysti	.s Schro	eter	i Choda	at		
	13,14	Tetraedron	gracile	(Re	insch)	Hansgirg		
	15,16	11	hastatu	m	11	f f		
	17,18	**	limneti	cum 1	Borge	var. gracile	Prescott	
	19	**	victori	eae	Wolosz	var. major S	mith	
	20	. 19	regular	e Ku	etz.			

		PLATE 5
Fig.	1	Oedogonium sp.
	2	Spirogyra sp. 2
	3	" sp. 1
	4	Cf. Mougeotia sp.
	5	Ulothrix subconstrict G.S.West
	6,7	Scenedesmus obliquus (Turp.) Kuetz. var. dimorphus May
	8-10	" quadricauda (Turp.) Breb.
	11	" " var. quadrispina Chodat
	12-15	" opoliensis Richter
	16	" denticulatus Lag. var. denticulatus May
	17	Arthrodesmus convergens Ehrenb.
	18	Cosmarium quadrum Lund
	19	" granatum Breb. var. pyramidal Schmidle
	20	" magnificum Nordst.
	21	" punctulatum Breb. var. subpunctulatum Borge
	22-24	" bipunctatum Borgesen
		PLATE 6
	1	Closterium aciculare T.West var. subpronum West & West
	2	" Kutzingii Breb.
	3	" ceratium Perty
	4	" gracile Breb.
	5,6	" monoliferum (Bory) Ehrenb.
	7	" striolatum Corda
	8	" Venus Kutz. var. inflatum Claassen
	9	Micrasterias Hardyi G.S.West
	10	" thomassiana var. notata (Nordst.) Gronbl.
	11	" mahabuleswarensis Hobson
	12	decemdentata (Nag.) Archer.

Fig.	1	Pleurotaenii	um Ehrenbergii (Breb.) De Bory	
	2,3	Gonatozygon	kinahani (Arch.) Rabenh.	
	4		monotaenium De Bory	
	5	Spondylosium	n planum (Wolle) West & West	
	6	Cosmarium co	ontractum Kirchn.	
	7 .	Euastrum ele	gans Kutz.	
	8	" ana	asatum (Ehrenb.) Ralfs	
	9-11	" cuspie	latum var. goyazense Forster	
	12	" diver	gens Joshua	
	13	Staurodesmu	s cuspidatus (Breb) Ralfs var. tricuspidatus	
	14,15	Arthrodesmu	s constrictus G.M.Smith var. longispinus Gro	nb.
	16-18	Staurodesmu	s triangularis (Lag.) Teiling	
	19-21	Cosmarium c:	f. pseudonitidum Nordst.	
		PLA	re 8	
	11	Staurastrum	Playfairi Scott & Prescott	
All	the oth	er figures a:	re of Staurastrum nodulosum Prescott	5
		PLA	LE 9	
Fig.	1-5	Staurastrum	dickiei Ralfs	
	6	11	grande Bulhn. var. Parvus West	
	7,8	11	pachyrhynchum Nordst.	
	9	11	Freemanii West & West	
	10	18	patens Turner	
	11	11	sp.	
	12	11	asterias Nygaard	
	13,14	19	furcatum (Ehrenb.) Breb.	
	15,16	ŧŢ	tetracerum Ralfs	
	17	11	cf. subpinnatum Turner var. subpinnatum Sch	mid.
	18	19	subgemmulatum West & West	
	19-21	. 19	tetracerum Ralfs var. evolutum West & West	

Fig.	1	Staurastrum	anatoides Scott & Prescott
	2,4	!!	sebaldi Reinsch. var. ornatum Nordst.
	3	11	pseudosebaldi Wille var. planctonicum Teil.
	5,6	tt ·	victorience West
	7,8	tt.	arctiscon (Ehrenb.) Lund var. glabrum Wests
	9	· • • • •	tohopekaligense Wolle
		PLAT	<u>E 11</u>
	1-6	Staurastrum	pingue Teiling
	7-13	11	" " (small forms)
		PLAT	DE 12
	1	Vorticella s	•de
	2	Ceratium hir	undinella (O.Muell) Dujardin
	3	Glenodinium	sp.
	4	Peridinium s	sp.
	5	Mallomonas a	krokomos Ruttner
	6	17 E	plendens (G.S.West) Playfair
	7	Synura spinc	osa Korsh.
	8	Dinobryon di	vergens Imhof.
	9	n se	ertularia Ehrenb.
	10-12	Mallomonas s	sp.
	13-15	Trachelomona	s scraba Playfair
	16-19	F1	bacillifera Playfair
• .	20,21	11	hispida (Perty) Stein var. rectangularis
	22	11 A	" var. coronata Lem m.
	23	11	superba Swirenko
	24	**	armatus var. longispina (Playfair) Deflandre
	25	. H.	caudata Stein var. australica Playf.
	26	18	volvocina Ehrenb. var. punctata Playf.

Fig.	16	Attheya Zachariasi J.Brun.
	7	Cyclostella stelligera Cleve & Grun.
	8	Melosira granulata Ralfs
	9	" varians Agardh.
	1 0-12	Rhizosolinia erensis H.L.Sm var. morsa West & West
	13	Rhizosolinia erensis H.L.Sm
	14	Stauroneis fulmen var. capitata
	15	Nitzschia pelagica Ø.Mull.
	16	Synedra acus Kutz.
	17	" ulna (Nitz.) Ehr. var. danica (Kg.) V.H.
	18,19	" acus Kutz. var. radians (Kg.) Hust.
	20	" acus Kutz.
		PLATE 14
	1	Surirella spinifera Hust.
	2	" Engleri Ø.Mull.
	3	" tenera Greg.
	4	Pinnularia abaujinsis (Pant.) Ross var. abaujinsis
	5	Epithemia sorex Kutz.
	6	Fragilaria capucina Desm.
	7	Fragilaria sp.
	8	Tabellaria flocculosa (Roth) Kutz. var. flocculosa Knudson
	9	Cf. Gyrosigma attenuatum
	10	Unidentified diatom
	11	Gomphonema Augur Ehr. var. Guatieri H.V.H.
	12	Asterionella formosa Hass.
	13	Cf. Navicula hennedyi
	14	Gomphonema constrictum Ehrenb.
	15	" " var, capitatum Ehrenb.
	16	Cymbella gastroides (cf.) Kutz. 17 Cymbella lanceolata Ehrenb
		18 Unidentified diatom



PLATE I



500 M




















100 /

PLATE 12





LITERATURE CITED

AHLSTROM E.H. (1937) Studies on variability in the genus <u>Dinobryon</u> Trans. Amer. Microscop. Soc. Vol. LVI (2).

BAILEY (1893) Contributions to the Queensland Flora. Botany Bullet. No VI

(1895) " " " " " " " " <u>No XI</u> (1898) " " " " <u>No XI</u> BERGEN G. (1946) Desnidiaceae from Tisleia, Central Norway. Nytt. Maj. f. Naturv. B. 88

BICUDO (1969) Contribution to the knowledge of the desmids of the state of Sao Paulo (Brazil). Nova Hedwigia Band XVII BICUDO & SORMUS L. (1972) Polymorphism in the desmid <u>Micrasterias</u> laticeps and its taxonomic implications. J.Phycol <u>8</u> p. 237-242. BORGE O. (1896) Australische Susswasserchlorophyceans. Bihang Till Svenska Vet.-Akad. Handolingar Band 22 Afd. III No 9.

(1903) Die Algen der esstan Regnellschen Expdition II Desmidiacean. Arkiv. for Botnik Band I.

(1911) Susswasseralgen ars Queenland. Algologische Notizen. 6-7. Bot. Not.

(1918) Die von Dr. Lofgren in Sao Paulo gessammelten Susswasseralgen. Arkiv. for Botanik Band 15 No 13 p. 1-108 BOURRELLY P. (1957) Algues d'eau douce du Soudan Francais, region du Macina (A.O.F.) Bulletin de 1'I F.A.N. T. XIX

(1968) Les algues d'eau douce algues Jaunes et brines (1966) Quelques Algues d'eau douce du Canada Int. Revue ges. Hydrobiol <u>51</u> (1) p. 45-126

BROOK A.J. (1959) <u>St.paradoxum</u> Meyen and <u>St.gracile</u> Ralfs in thefresh water plankton (British) and a Revision of <u>St.anatinum</u> group of Radiate Desmids. Trans. Roy. Soc. Edin. vol. LXIII Part III No 26. BROOK A.J. (1959b) The status of desmids in the plankton and the determination of phytoplankton quotients. J.Ecol. 47

(1965) Planktonic algae as indicators of lake types, with special reference to the Desmidiaceae. Limnol. and Oceanogr. 10 (3) p. 403

BROOK A.J. & HINE A.E. (1966) <u>St.freemanii</u> from the Central Highland of New Guinea. J. Phycol. <u>3</u> p. 66-73

CHUAH T.I. (1969) Taxonomic studies of the genus <u>Staurastrum</u> B.Sc (Hons.) Thesis, Uni. of Tas.

CLAASSEN M.I. (1961) A contribution to our knowledge of the fresh water algae of the Transvaal Province. Bothalia VII part 3. CONRAD W. (1933) Revision du Mallomonas Perty (1851) including Pseudo-mallomonas Chodat (1920). Mem. du Roy. D' histore Naturelle de Belgique Mem. No 56

COOK P.W. (1963) Variation in vegetative and sexual morphology among the small curved species of <u>Closterium</u>.Phycologia <u>3</u> (1) p. 1-18

CROOME R.L. & TYLER P.A. (1972) Physical and chemical limnology of Lake Leake and Tooms Lake, Tas. Arch. Hydrobiol. <u>70</u> (3) p.341-4 DODGE J.D. & CRAWFORD R.M. (1970) The morphology and fine structure of <u>Ceratium hirundinella</u>. J. Phycol <u>6</u> p. 137-149

EDWARDS A.M. (1894) On species in the Desmidiae. Microscopic Jour. October 1894

FLINT E.A. (1966) Additions to the check list of fresh water algae of New Zealand. Trans. of the Roy. Soc. of NZ <u>3</u> (7). FORSTER K. (1964) Desmidiaceen aus Brasilien 2 Teil Bahia Goyaz, Pianhy und Nora-Brasilien. Hydrobiologia Acta Hydrobiologica Hydrographica et Protistologica XXIII Fasc. 3-4. FORSTER K. (1965) Beitrag zur Desmidieen-Flora der Torne-Lappmark in Schwedisch-Lappland. Arkiv for Botanik Serie 2 Band 6 Nr 3.

(1966) Beitrag zur Desmidiaceen des ost-allgaius Sonderdruck aus den Berichten Der Bayerischen Botanischen Gesellschaft. Band XXXIX

(1967) <u>Staurastrum pingue</u> Teiling und einige andere Staurastren aus dem Titisee (Schwarzwald). Arch. Hydrobiol/suppl. XXXIII (1) p. 121-126

(1968) Beitrag zur Desmidiaceenflora des Ost-Allgaus Sonderdruck aus den Berichten der Bayerischen Botanischen Gesellschaft. Band XL

(1969) Amazonische Desmidiean 1 Teil: A real Santarem Amazona Limnologia et occologia Regionalis Systemae fluminis Amazonas Vol. 2

FRERE-IRENEE-MARIE (1939) Flora Desmidiale De la Region de Montreal. Leprarie P.Q.

(1951) Desmidius de la Region quebeg. Le Canadian Vol.LXXVIII No 10 GRONBLAD R. (1920) Finnlandische Desmidiaceen aus keuru. Acta Societatis pro fauna et flora Fennica <u>47</u> (4).

(1926) Beitrag zur Kenntnis der Desmidiacean Schlesiens. Societas Scientiarum Fennica Commentationes Bioloicae II (5).

(1945) De Algis Brasiliensibus, Praccipue Desmidiaceis, in regione inferiore fluminis Amazones. A professore August Ginzberger (Wien) Anno MCMXXVII Collectis. Acta societatis Scientiarum Fennicae Nova series B. Tom II No 6.

(1960) Contributions to the knowledge of fresh water algae of Italy. Societas Scientiarum Fennica Commentations Biologicae XXII (4).

GRONBLAD R. & CROASDALE H. (1971) Desmids from Namibia (S.W.Africa)

Acta Botanica Fennica 93

GRONBLAD R., PROWSEG.A. & SCOTT A.M. (1958) Sudanese desmids. Acta Botanica Fennica 58

GRONBLAD R. & SCOTT A.M. (1955) On the variation of <u>Staurastrum</u> <u>bibrachiatum</u> as an example of variability in a desmid species. Acta Soc. pro Fauna et Flora 72 (6) p. 1-11

GRONBLAD R., SCOTT A.M. & CROASDALE H. (1964) Desmids from Uganda and Lake Victoria collected by Dr.E.Lind. Acta Botanica Finnica <u>66</u> HARDY A.D. (1906) The fresh water of Victoria part III. Victorian Naturalist XXIII p. 18-22.

HARROP E.D. (1869) Conversations on Desmidiaceae, with a list of species found in Tasmania. Roy. Soc. Tas. Pap. Proc. p.19-22. HARRIS K. (1958) A study of <u>Mallomonas insignis</u> and <u>Mallomonas</u> <u>akrokomos</u>. The Jour. of general Microbiol. 19 (1).

(1970) Imperfect forms and the taxonomy of <u>Mallomonas</u> Jour. of Gen. Microbiol. <u>61</u> p. 73-76

HARRIS K. & BRADLEY D.E. (1960) Ataxonomic study of <u>Mallomonas</u> Jour. of Gen. Microbiol. <u>22</u> (3).

HAUGHEY A. (1968) The planktonic algae of Auckland sewrage treatment ponds. NZ Jour. of marine and fresh water research <u>2</u> (4).

(1969) Further planktonic algae of Auckland sewrage treatment ponds and other waters. NZ Jour. of marine and fresh water_research 3 (2).

HEURCK (1896) A treatise on the diatomaceae. Willium Wesley & Son Publisher London.

HINODE T. (1966) Desmids from the northern district of Tokushima Prefecture (3). Jour. Jap. Bot. 41 (10).

HIRANO M.(1961-2) Fresh water algae collected by the joint Thai-Japanese Biological Expedition to South-East Asia 1961-2.

Nature and life in South-East Asia vol. V.

119

HIRANO M. (1964) Fresh water algae of Afghanistan: Results of the Kyoto University Scientific Expedition to the Karakoram & Hudukush 1955 vol. III.

(1972) Desmids from Cambodia, with special reference to Phytoplankton of Lakes Grands Lacs: Contributions from the biol. lab. Kyoto Uni. vol. 23 (3/4) p. 123-157.

HUBER-PESTALLOZZI (1936) Phytoplankton aus und Sumpfen Javas gesammelt von Prof. C. Schroter-Zurich. Schweizerische Botanis Gesellsch Berichte <u>46</u> p. 131-168.

(1941) Die Binnengewasser von Prof. Dr. August Thienemann. Band XVI Teil 2 Halfte 1 Das phytoplankton des Swsswasswassers.

(1942) same as above Halfte 2.

ISLAM N. (1970) Contributions to the knowledge of desmids of East Pakistan part 1. Nova Hedwigia Band XX.

JAHN T.L. (1949) How to know the Protozoa. W.M.C.Brown Company Publishers Dubuque Iowa.

JOSHUA W. (1886) Burmese Desmideae, with descriptions of new species occurring in the neighbourhood of Rangoon. Linn. Soc. Jour. <u>21</u> p. 634. KNUDSON B,M. (1952) The diatom genus Tabellaria 1) Taxonomy and Morphology. Ann. Bot. NS XVI p. 421

KRIEGER W. (1939) Die Desmidiaceen vol. XIII

Section I part 2 No I Conjugatae

(1937) same as above

Section I part 1 No 1-4.

LEEDALE F.G. (1967) Euglenoid Flagellates. Prentice-Hall Inc. LEEDALE G.F., MEEUSE J.D. & PRINSHEIM E.G. (1965) Structure and Physiology of <u>Euglena spirogyra</u>. Arch. Mikrobiol. <u>50</u> p. 133-155 LIND E.M. (1967) Some East African desmids. Nova Hedwigia Band XIII p. 361-387. LIND E.M. (1971) Some desmids from Uganda. Nova Hedwigia XXII. LIND E.M. & CROASDALE H. (1966) Variation in the desmid <u>Staurastrum</u> <u>sebaldi var. ornatum</u> Jour. of Phycol. <u>2</u> (3) p. 111-116 LOKHORST G.M. & VROMAN M. (1972) Taxonomic study on three fresh water <u>Ulothrix</u> species. Acta Bot. Neerl. <u>21</u> (5) p. 449-480. LYON T.L. (1969) Scanning electron microscopy : a new approach to the desmidiaceae. J. Phycol. <u>5</u> p. 380-382. MASKELL N.M. (1881) Contribution towards a list of NZ desmiaceae.

Tran. Proc. NZ 13 p. 297-317.

(1883) On the NZ desmidiceae: additions to Catalogue and notes on various species. Trans. Proc. NZ Inst. <u>15</u> p. 237-259. MAY V. (1973) Algal genus <u>Scenedesmus</u> in Australia. Contributions from the N.S.W. National Herbarium vol. <u>4 no</u> 7. MESSIKOMMER E. (1927) Biologische Studien in Tomfmoor von

Rabenhausen. Mitt. Bot. Mas. Uni. Zurich 122.

MOBIUS M. (1892) Australische Sufswasseralgen III Flora 3

(1894) same as above . Senckenbergische Naturforschande Gesellschaft <u>18</u> p. 310-350.

NORDSTEDT O. (1872) Bidrag till kamedomen one sydligare morges desmidiaceae. Roy. Soc. Catalog. of Scientifig papers vol. VIII p. 1-50.

(1888) Fresh water algae collected by Dr. S. Berggren in NZ & Australia. Kogl. Svenska. Vetenskaps. Akademiens Handlingar Bandet 22 No 8.

NYGAARD G. (1926) Plankton from two lakes of the Malayan region Dansk. Natur. For. Vidensk. Meddel. <u>82</u> p. 197-240.

NYGAARD G. (1949) Hydrobiological studies on some Danish ponds and lakes. Part II The quotient hypothesis and some new or little phytoplankton organism. Det Kongelige Danske Videnskabernes Selskab Biologiske Skrifter Bind VII nr I.

PATRICK R. & REIMER C.W. (1966) The diatom of the United Stated

exclusive of Alaska ana Hawaii Vol. 1. Monographs of the Academy of Natural Sciences of Philadelphia No 13.

PETERSEN J.B. & HANSEN J.B. (1956) On the scales of some <u>Synura</u> species. Biol. Medd. Dan. Vid. Selsk. <u>23</u> (2).

PHILIPOSE M.T. (1967) Chlorococcocales. Indian council on Agriculture Research. New Delhi.

PLAYFAIR G.I. (1912) Plankton of the Sydney water supply. Proc. Linn. Soc. N.S.W. XXXVII p. 512.

(1914) Contributions to a knowledge of the biology of the Richmond River. Proc. Linn. Soc. N.S.W. XXXIX p. 93.

(1915) The genus <u>Trachelomonas</u>. Proc. Linn. Soc. N.S.W. XL p. 1-41

(1916) <u>Oocystis</u> and <u>Eremosphaera</u>. PROC. LINN. SOC. N.S.W. vol. XLI p. 107-147

(1918) New and rare fresh water algae. Proc. Linn. Soc. N.S.W. vol. XLIII p. 497-543.

(1921) Australian fresh water flagellates. Proc. Linn. Soc. N.S.W. vol. 46. p. 99-146.

(1923) Notes on fresh water algae. Proc. Linn. Soc. N.S.W. vol. VLVIII (3) p. 206-228.

PRESCOTT G.W. (1936) Notes on the algae of Gatun Lake, Panama Canal. Trans. Amer. Microscop. Soc. Vol. IV p. 501-509.

(1955) Algae of the Panama Canal and its tributaries. Part 1 Flagellated organisms. Ohio J. Sci. Vol. LV (2) p. 99-121.

(1962) Algae of the Western Great Lakes Area . WM. C. Brown Company Publishers.

(1966) Algae of the Panama Canal and its tributaries Part 2 Conjugales. Phykos. <u>5</u> p.1-49.

PRESCOTT G. W. & CROASDALE H. (1937) New or noteworthy fresh water algae of Massachusetts. T ans. of Amer. Microscop. Soc. Vol LVI (3). PRESCOTTG.W. & SCOTT A.M. (1942) The fresh water algae of southern United States. Part 1 Desmids from Mississippi, with description of new species and varieties. Trans. Amer. Microscop. Soc. 61 (1) p. 1-29.

(1945) The fresh water algae of the Southern United Stated. Part 3 The desmid genus <u>Euastrum</u>, with descriptions of some new varieties. The Amercan Middland Naturalist <u>34</u>. (1952) Some south Australian desmids.

Trans. Roy. Soc. S. Aust. 75 p. 55-69.

PROWSE G.A.(1957) Introduction to the desmids of Malaya. Malayan Nat. J. <u>11</u> p. 42-58.

REHAKOVA H. (1969) Die Variabilitat der Arten der Gattung <u>Oocystis</u> A. Braun. IN Studies in Phycology edited by Fott B. Academia Publishing House of the Czechoslovak Academy of Sciences (1969). RICH F. (1932) Contributions to our knowledge of fresh water algae of Africa. S. African Trans. Vol. XX part 2.

RAMANATHAN K.R. (1964) Ulotrichales. Indian Council of Agricultural Research New Delhi.

SCHMIDLE W.(1896) Susswasseralgen aus Australien. Flora <u>82</u> p. 297 SCHUMACHER G.J. KIM Y.L. et al (1966) Additions to the fresh water in North Carolina. Part VII j. Elisha Mitchell Scientific soc. <u>82</u> (2). SCOTT .M. & GRONBLAD (1957) New and intersting desmids from the Southern United States. Acta Societatis Scientiarum Fennica Nova Series B II no 8

SCOTT A.M., GRONBLAD R. & CROASDALE H. (1965) Desmids from the Amazon basin, Brazil. Acta Botanica Fennica <u>69</u> p. 1-93. SCOTT A.M. & PRESCOTT G.W. (1958a) Notes on Indonesian fresh water algae III. New varieties of some little known <u>Staurastra</u>. Reinwardlia vol. 4 part 3.

123

SCOTT & PRESCOTT (1958) Some fresh water algae from Arnham land in the Northern territory of Australia. Reprinted from the American Australia Scientific expedition to Arnham land. Vol. 3 SCOTT A.M. & PRESCOTT G.W. (1961) Indonesian desmids. Hydrobiologia Acta Hydrobiologica Hydrographica et Protistologica <u>17</u>. SKUJA H. (1949) Zur Susswasseralgen-Flora Burmas. Nova Acta Regiae Societatis Scientiarum Upsaliensis ser IV vol. 14 (5). Smith G.M. (1920) Phytoplankton of the inland lakes of Wisconsin. Part 1 Myxophyceae, Phaeophyceae, Heterokontae and Chlorophyceae exclusive of the desmidiaceae. Wisconsin Geological and Natural History Survey bulletin No 57 Scientific series No 12.

(1921) The phytoplankton of the Muskoka region Ontario, Canada. Trans. of the Wisconsin Acad. of Science, Arts, and letters. Vol. XX

(1924) Desmidiaceae. Bulletin of Uni. of Wis. serial No 1270 general serial No 1048.

(1944) A Comparative study of the species of <u>Volvox</u>. Trans. Amer. Microscop. Soc. Vol. LXIII (4) p. 265-310. SPENCER W.I. (1882) On the fresh water algae of NZ. Trans. Proc. Roy. Soc. NZ <u>14</u>.

STEIN J.R. & GERRATH J.F. (1968) Fresh water algae of Columbia(Canada): desmids of bog lakes in Coast Mountain area. Syesis <u>1</u> p. 187-197. SULEK (1969) Taxonomissche Ubersicht Der Gattung <u>Pediastrum</u> Meyen IN Studies in Phycology edited by B. Fott. Publishing House of the Acad. of Sci.

Teiling E. (1942) Schwedische Planktonalgen 3 Neue oder wening bekannte Formen Bot. Notiser 1942.

(1946) Zur Phytoplankton Flora Schwedens Botaniska Not. 46 p;61-88 TEILING E. (1948) <u>Staurodesmid</u>, genus novum. Botaniska Notiser Halfe 1 Lund.

(1967) The desmid genus <u>Staurastrum</u>. A Taxonomic study. Arkiv for Botnik Serie 2 Band 6 nr. 11 THOMASSON K.(1957) Notes on the phytoplankton of lake Bangweulu.

Nova Acta Regiae Soc. Sci. Upsaliensis ser IV 17 (3).

(1960) Notes on the phytoplankton of Lake Bangweulu Part 2 Nova Acta Soc. Sci. Upsaliensis ser IV <u>17</u> (12)

(1963) Araucanian Lakes Plankton Studies in North Patagonia with notes on Terrestrial vegetation. Acta Phytogeograph. Suecica <u>47</u>.

(1965) Notes on Algal vegetation of Lake Kariba. Nova Acta Regiae Soc. Sci. Upsaliensis ser IV vol. 19 (1).

(1967) Phytoplankton from some lakes on Mt. Wilhelm East New Guinea. Blumea <u>15</u>.

(1972) Some planktic <u>Staurastra</u> from NZ (2) . Sven. Bot, Tidsk. Bd. 66 p. 257-274.

(1973) Notes on the plankton of some Sydney Reservoirs, with description of two intersting desmids. Contribution from the N.S.W. National Herbarium vol. 4 (6).

(1973a) <u>Actinotaenium</u>. <u>Cosmarium</u>. <u>& Staurastrum</u> in the plankton of Rotorua Lakes. Sven. Bot. Tidsk. Bd. 67 p. 127-141.

(1974) Some planktic <u>Staurastra</u> from NZ (3). Sven. Bot. Tidsk. Bd 68 p. 33-50.

THOMASSON K. & TYLER P.A. (1971) Taxonomy of Australian fresh water algae (2) Some planktic <u>Staurastra</u> from Tasmania. Nova Hedwigia Band XXI p. 287-319.

TYLER P.A. (1970) Taxonomy of Australian fresh water algae I The genus <u>Micrasterias</u> in South Eastern Australia. Br. Phycol. Jour. <u>5</u> (2) p. 211-234. TRAINOR F.R. & HILTON R.L. (1963) Identification of species of <u>Scenedesmus</u>. Nature 200 (4908) p. 800.

TURNER W.B. (1892) The fresh water algae (principlely Desmidiaceae) of East India. Kogl. Sven. Veten.-Akad. Handl. Bd. 25 (5). WOOD E.J.F. (1960) Studies on Australian & NZ diatoms, part VI description of further sedentary species. Trans. Roy. Soc. NZ 88 p. 669.

WEST W. & WEST G.S. (1902) A Contribution to the fresh water of Ceylon. Trans. Linn. Soc. Lond. IV (3) p. 123-215. WEST G.S. (1909) The algae of Yan Yean Reservoir, Victoria, a biological and oecological study. J. Linn. Soc. (Bot.) <u>39</u> p. 1-88. WEST W. & WEST G.S. (1904, 1905, 1908, 1912) A monograph of the British Desmidiaceae Vol. 1-4 respectively. Ray Soc. Lond. WEST G.S. & CARTER N. (1923) A monograph of the British Desmidiaceae Vol. 5.